

Relations between connected and self-avoiding walks in a digraph

Thibault Espinasse* and Paul Rochet†

Abstract

Walks in a directed graph can be given a partially ordered structure that extends to possibly unconnected objects, called hikes. Studying the incidence algebra on this poset reveals unsuspected relations between walks and self-avoiding hikes. These relations are derived by considering truncated versions of the characteristic polynomial of the weighted adjacency matrix, resulting in a collection of matrices whose entries enumerate the self-avoiding hikes of length ℓ from one vertex to another.

Keywords: Directed graph; poset; characteristic polynomial; weighted adjacency matrix; incidence algebra.

MSC: 05C22, 05C30, 05C38

1 Introduction

A directed graph $G = (V, E)$ is defined by a vertex set $V = \{v_1, \dots, v_N\}$ and set E of ordered pairs of vertices representing the directed edges of G . Directed graphs, or digraphs, have been extensively used in the literature as mathematical models to describe actual phenomena such as social interactions [5], road traffic [9], physical processes [4] or random walks [3] among many others. In most models, it is convenient to allocate weights to each edge of the graph in order to incorporate some additional information. For example, a weight ω_{ij} between two nodes v_i, v_j may serve defining the transition probability of a random walk, the speed limit or the type of road in the traffic network and so on. A finite graph on N vertices is then characterized by its *weighted adjacency matrix* $W = (\omega_{ij})_{i,j=1,\dots,N}$, which accounts for the level of interaction between vertices. In this way, every edge of G is identified with a weight ω_{ij} .

An oriented walk on the digraph G is defined as a succession of contiguous edges ω_{ij} . Seeing the weights ω_{ij} as formal variables, a walk w of length ℓ from v_i to v_j can be viewed as a degree ℓ monomial $w = \omega_{i i_1} \omega_{i_1 i_2} \dots \omega_{i_{\ell-1} j}$. The weighted adjacency matrix then provides a practical tool to handle walks on the graph, as they can be derived from analytical transformations of W . For

*Université Lyon 1 Claude Bernard, laboratoire de mathématiques Camille Jordan

†Université de Nantes, laboratoire de mathématiques Jean Leray, paul.rochet@univ-nantes.fr

instance, the (i, j) entry of W^2 , given by $W_{ij}^2 = \sum_k \omega_{ik}\omega_{kj}$, enumerates all walks of length 2 from v_i to v_j . The introduction of the weighted adjacency matrix W to describe the walks on a graph goes back to the 60's. In [10] and [17], the spectral properties of a graph are investigated via the determinant and characteristic polynomial of W . Digraphs also provide a useful tool to compute the determinant and minors of sparse matrices, as discussed in [15]. For general results on spectral graph theory, we refer to [7, 8].

In [17], the author shows that the coefficients of the characteristic polynomial of W can be interpreted in term of the self-avoiding cycles in G . In this paper, we derive a similar result concerning self-avoiding hikes, defined as a generalization of walks to possibly unconnected sequences of edges. We construct a collection of polynomials of W whose entries enumerate the self-avoiding hikes of a given length from one vertex to another. The polynomials are obtained as Cauchy products of the characteristic polynomial coefficients with the sequence of successive powers of W .

The analytical expression linking the self-avoiding hikes and the walks on the graph hides a deeper connection when considering each hike individually. Precisely, the relation can be investigated in the partially ordered set formed by the hikes. In this context, combinatorial properties arise when studying functions of the hikes in the reduced incidence algebra of this poset. In particular, we show that the number of different ways to travel a closed hike can be expressed in term of its self-avoiding divisors via a Mobius-like inversion on this poset. Another result on the decomposition of a walk into self-avoiding components is then derived.

The paper is organized as follows. Definitions are introduced in Section 2 as well as the preliminary result. Section 3 is devoted to the study of the different relations between self-avoiding hikes and walks, where many combinatorial properties are investigated. The results are verified on specific examples in Section 4.

2 Notations and preliminary results

Let $G = (V, E)$ be a labeled directed graph, or *digraph*, with finite vertex set $V = \{v_1, \dots, v_N\}$ and edge set E which may contain loops. The adjacency matrix of G is defined as the $N \times N$ matrix A with entries a_{ij} equal to one if v_i is connected to v_j and zero otherwise. Because G is directed, A may not be symmetric (v_i can be connected to v_j without v_j being connected to v_i). In this paper, an edge always refers to a directed edge.

The adjacency matrix can be used to derive numerous properties of a graph. For instance, the (i, j) entry of A^ℓ gives the number of walks of length ℓ from v_i to v_j . When one is interested in each walk specifically, a useful tool is to allocate a weight, or variable, to each non-zero entry of A . In this way, the digraph G is characterized by its weighted adjacency matrix

$$W = (\omega_{ij})_{i,j=1,\dots,N}$$

where the ω_{ij} 's are real variables, setting $\omega_{ij} = 0$ whenever there is no edge from v_i to v_j . An edge of G can then be identified with a non-zero variable ω_{ij} and a walk w from v_i to v_j with the product $w = \omega_{ii_1}\omega_{i_1i_2}\dots\omega_{i_{\ell-1}j}$ of the edges composing it (two walks are thus considered equal if

they are composed of the same edges, counted with multiplicity, regardless of their order). The walk w is closed (a cycle) if $i = j$ and open (a path) otherwise. Moreover, w is simple if it does not cross the same vertex twice, that is, if the indices $i, i_1, \dots, i_{\ell-1}, j$ are mutually different, with the possible exception $i = j$ if w is closed. Loops ω_{ii} and backtracks $\omega_{ij}\omega_{ji}$ are considered cycles of length 1 and 2 respectively.

The representation as monomials in the formal variables ω_{ij} provides a simple multiplicative structure on walks. The cycle-erasing procedure of Lawler [13] shows that a walk from v_i to v_j can always be decomposed as the product of a simple walk from v_i to v_j and cycles, as illustrated in Figure 1. However, the reverse is not true in general as the product of a simple walk and simple cycles might not be connected. In this paper, we define a new object, called hike, which extends the definition of a walk by relaxing the connectedness condition.

Definition 2.1 *A hike from v_i to v_j is a monomial $h = \omega_{i_1 j_1} \dots \omega_{i_\ell j_\ell}$ that can be decomposed into the product of a simple walk from v_i to v_j and simple cycles.*

Properties of walks naturally extend to hikes. A hike is closed if it is a product of simple cycles and open otherwise. By convention, the trivial cycle 1 is considered a closed hike of length 0. Similarly as for walks, a hike is self-avoiding if it does not cross the same vertex twice. The connected components of a self-avoiding hike are simple and vertex-disjoint. In this setting, a walk can be viewed as a connected hike.

In the sequel, the length of a hike h (its degree) will be denoted by $\ell(h)$ and its number of connected components by $n(h)$. A walk is a connected hike, that is, a hike h such that $n(h) = 1$. Moreover, we denote by $V(h)$ the set of vertices crossed by h and $|V(h)|$ its cardinal. Remark that an open hike h is self-avoiding if, and only if, $|V(h)| = \ell(h) + 1$ while for a closed hike, self-avoiding is equivalent to $|V(h)| = \ell(h)$.




Figure 1: The walk $w = \omega_{12}\omega_{23}\omega_{34}\omega_{42}\omega_{25}$ from v_1 to v_5 is the product of the simple walk $\omega_{12}\omega_{25}$ and simple cycle $\omega_{23}\omega_{34}\omega_{42}$.




Figure 2: The self-avoiding hike $h = \omega_{12}\omega_{34}\omega_{45}\omega_{53}$ from v_1 to v_2 contains $n(h) = 2$ connected components: the path ω_{12} and the cycle $\omega_{34}\omega_{45}\omega_{53}$.

The following notations are used throughout the paper.

- \mathcal{W} denotes the set of walks.
- \mathcal{H} denotes the set of hikes (open and closed).

- \mathcal{C} denotes the set of closed hikes.
- \mathcal{S} denotes the set of self-avoiding hikes.

For these sets, we may specify the end-vertices in index and/or their length in exponent, e.g. \mathcal{H}_{ij}^ℓ is the set of hikes of length ℓ from v_i to v_j . Remark that \mathcal{W} , \mathcal{C} and \mathcal{S} are subsets of \mathcal{H} , while \mathcal{S}_{ii} and \mathcal{W}_{ii} are subsets of \mathcal{C} for all $i = 1, \dots, N$. The set of self-avoiding closed hikes is written as $\mathcal{C} \cap \mathcal{S}$.

It is known that the ℓ -th power of \mathbf{W} enumerates with multiplicity the walks of length ℓ on the graph. The (i, j) entry of \mathbf{W}^ℓ is a homogenous degree ℓ polynomial attributing to each walk w of length ℓ , a coefficient $f_{ij}(w)$ counting the number of ways to write w as a succession of contiguous edges starting from v_i and ending at v_j (the function f_{ij} is computed on some examples in Figure 3). The formal series associated to the functions f_{ij} follows by the identity

$$(\mathbf{I} - \mathbf{W})_{ij}^{-1} = (1 + \mathbf{W} + \mathbf{W}^2 + \dots)_{ij} = \sum_{w \in \mathcal{W}} f_{ij}(w)w, \quad (1)$$

which holds whenever $\|\mathbf{W}\| := \sup_{v \in \mathbb{R}^N, \|v\|=1} \|\mathbf{W}v\| < 1$, with $\|\cdot\|$ the Euclidean norm in \mathbb{R}^N . Remark that if w is an open walk, there is at most one couple (i, j) for which $f_{ij}(w)$ is non-zero. This property is no longer verified for a closed walk c in which case $f_{ii}(c)$ may take different positive values for different nodes $v_i \in V(c)$, as illustrated in Figure 3.




Figure 3: The walk $w = \omega_{19}\omega_{93}\omega_{32}\omega_{29}\omega_{95}\omega_{54}\omega_{49}\omega_{97}\omega_{76}\omega_{69}\omega_{98}$ (gray and black edges) has $f_{18}(w) = 6$ contiguous representations starting from v_1 and ending in v_8 . Each representation relies on an ordering of the simple cycles $\omega_{93}\omega_{32}\omega_{29}$, $\omega_{95}\omega_{54}\omega_{49}$ and $\omega_{97}\omega_{76}\omega_{69}$ to travel w . The cycle $c = \omega_{95}\omega_{54}\omega_{49}\omega_{97}\omega_{76}\omega_{69}$ (black edges) has $f_{99}(c) = 2$ contiguous representations starting from v_9 and $f_{ii}(c) = 1$ contiguous representation for any other vertex $v_i \in V(c) \setminus \{v_9\}$.

Closed hikes also play an essential part in analytical graph theory. A simple reason is that a self-avoiding closed hike can be associated to a permutation σ on a subset of $\{1, \dots, N\}$ by the relation $c_\sigma = \prod_{v_i \in V(c_\sigma)} \omega_{i\sigma(i)}$. Using that the number of connected components $n(c_\sigma)$ is linked to the signature of σ through the identity $\text{sgn}(\sigma) = (-1)^{N-n(c_\sigma)}$, we obtain an expression of the

determinant of W by

$$\det(W) = \sum_{\sigma} \operatorname{sgn}(\sigma) \omega_{1\sigma(1)} \dots \omega_{N\sigma(N)} = (-1)^N \sum_{c \in \mathcal{C} \cap \mathcal{S}^N} (-1)^{n(c)} c$$

where the first sum is taken over all permutations over $\{1, \dots, N\}$ and the second sum over all self-avoiding closed hikes of length N . A more general formula, given in Theorem 1 in [17], links the coefficients ψ_k of the characteristic polynomial

$$\chi(\lambda) = \det(\lambda I - W) = \sum_{k=0}^N \psi_k \lambda^{N-k}$$

with the self-avoiding closed hikes of length k by

$$\psi_0 = 1, \quad \psi_k = \sum_{c \in \mathcal{C} \cap \mathcal{S}^k} (-1)^{n(c)} c, \quad k = 1, 2, \dots \quad (2)$$

the coefficient ψ_k being trivially zero for $k > N$.

The coefficient $(-1)^{n(c)}$ is reminiscent of a Mobius function. In fact, the function μ defined over \mathcal{C} by $\mu(1) = 1$ and

$$\mu(c) := \begin{cases} (-1)^{n(c)} & \text{if } c \text{ is self-avoiding} \\ 0 & \text{otherwise,} \end{cases} \quad (3)$$

is the Mobius function of the trace monoid of a partially commutative version of closed hikes, studied in [6] under the name circuit. The particular value of the characteristic polynomial

$$\chi(1) = \det(I - W) = \sum_{k=0}^N \psi_k = \sum_{c \in \mathcal{C}} \mu(c) c \quad (4)$$

will turn out to be particularly important for our purposes, as it gives the formal series associated to μ . Although the definition of μ is restricted to closed hikes, some of its properties have direct repercussions on open hikes. This is due to the fact that an open hike h between two different vertices v_i, v_j can be expressed as a closed hike to which the edge ω_{ji} has been removed. Actually, one verifies easily the following equivalence for $i \neq j$

$$h \in \mathcal{H}_{ij} \iff h\omega_{ji} \in \mathcal{C} \quad (5)$$

where we recall that \mathcal{H}_{ij} is the set of hikes from v_i to v_j . This property provides a natural extension of μ to open hikes. For all directed pairs of vertices (i, j) , define

$$\mu_{ij}(h) := -\mu(h\omega_{ji}) = \begin{cases} (-1)^{n(h)+1} & \text{if } i \neq j \text{ and } h \in \mathcal{S}_{ij} \\ (-1)^{n(h)} & \text{if } i = j, h \in \mathcal{C} \cap \mathcal{S} \text{ and } v_i \notin V(h), \\ 0 & \text{otherwise.} \end{cases} \quad (6)$$

If $i \neq j$, the function μ_{ij} only takes non-zero values for self-avoiding hikes from v_i to v_j . In particular, $\mu_{ij}(1) = 0$ and $\mu_{ij}(h) = 1$ if h is a simple path. On the other hand, $\mu_{ii}(h)$ is

non-zero only if h is closed and does not cross v_i . We have for instance $\mu_{ii}(1) = \mu(\omega_{ii}) = 1$ (the convention $\mu_{ii}(1) = 1$ instead of -1 justifies the minus sign in the definition of μ_{ij}). Remark moreover that $\mu_{ij}(h)$ is null for all hikes h of length $\ell(h) \geq N$ since a self-avoiding hike on the graph has maximal length N .

The main result of this paper (Theorem 3.2) exhibits a duality between the functions f_{ij} counting the number of connected representations of a hike and the functions μ_{ij} whose supports are restricted to self-avoiding hikes. The duality is actually a consequence of Lemma 2.2, which expresses the matrix formal series $\mathbf{M} = (m_{ij})_{i,j=1,\dots,N}$ of μ_{ij} , defined by

$$m_{ij} := \sum_{h \in \mathcal{H}} \mu_{ij}(h)h, \quad i, j = 1, \dots, N, \quad (7)$$

in function of the powers of \mathbf{W} .

Lemma 2.2 *If $\|\mathbf{W}\| < 1$,*

$$\mathbf{M} = \sum_{\ell \geq 0} \sum_{k=0}^{\ell} \psi_k \mathbf{W}^{\ell-k}. \quad (8)$$

where the $\psi_k, k = 0, 1, \dots$ are the coefficients of the characteristic polynomial defined in (2).

Proof. The proof relies on the adjugate identity $\text{adj}(\mathbf{B}) = \det(\mathbf{B})\mathbf{B}^{-1}$ applied to $\mathbf{B} = \mathbf{I} - \mathbf{W}$. Recall that for a matrix $\mathbf{B} = (b_{ij})_{i,j=1,\dots,N}$, the adjugate of \mathbf{B} is defined as

$$\text{adj}(\mathbf{B}) = (\det(\mathbf{B}^{(ji)}))_{i,j=1,\dots,N}, \quad (9)$$

where $\mathbf{B}^{(ji)}$ is the matrix obtained by setting $b_{ji} = 1$, $b_{ki} = 0$ for $k \neq j$ and $b_{jk} = 0$ for $k \neq i$ in \mathbf{B} . Combining Equation (4) with the identity $\text{adj}(\mathbf{I} - \mathbf{W}) = \det(\mathbf{I} - \mathbf{W}) (\mathbf{I} - \mathbf{W})^{-1}$ gives for $\|\mathbf{W}\| < 1$

$$\text{adj}(\mathbf{I} - \mathbf{W}) = \sum_{k=0}^N \psi_k \times \sum_{k \geq 0} \mathbf{W}^k = \sum_{\ell \geq 0} \sum_{k=0}^{\ell} \psi_k \mathbf{W}^{\ell-k} \quad (10)$$

with $\psi_k = 0$ for $k > N$. It remains to show that $\mathbf{M} = \text{adj}(\mathbf{I} - \mathbf{W})$. We proceed entry-wise, first considering the case $i = j$. By construction of the adjugate matrix in (9) for $\mathbf{B} = \mathbf{I} - \mathbf{W}$, the conditions $b_{ii} = 1$ and $b_{ki} = b_{ik} = 0$ for $k \neq i$ correspond to setting $\omega_{ki} = \omega_{ik} = 0$ for all k . Plugging these values into $\det(\mathbf{I} - \mathbf{W}) = \sum_{c \in \mathcal{C}} \mu(c)c$ sets to zero every cycle crossing v_i , yielding

$$(\text{adj}(\mathbf{I} - \mathbf{W}))_{ii} = \sum_{\substack{c \in \mathcal{C} \\ i \notin V(c)}} \mu(c)c = \sum_{h \in \mathcal{H}} \mu_{ii}(h)h. \quad (11)$$

Now consider the case $i \neq j$. Going back to Equation (10), we see that the (i, j) entry of $\text{adj}(\mathbf{I} - \mathbf{W})$ satisfies

$$(\text{adj}(\mathbf{I} - \mathbf{W}))_{ij} = \sum_{k=0}^N \psi_k \times \sum_{k \geq 0} \mathbf{W}_{ij}^k = \sum_{c \in \mathcal{C}} \mu(c)c \times \sum_{w \in \mathcal{W}} f_{ij}(w)w = \sum_{(c,w) \in \mathcal{C} \times \mathcal{W}} \mu(c)f_{ij}(w)cw.$$

The right-hand side is a sum over hikes of the form $h = cw$ with c a self-avoiding closed hike and w a walk from v_i to v_j . By (9), the left-hand side is obtained by plugging the values $\omega_{ji} = -1$, $\omega_{ki} = \omega_{jk} = 0$ for $k \neq i, j$ and $\omega_{ii} = \omega_{jj} = 1$ into $\det(\mathbf{I} - \mathbf{W}) = \sum_{c \in \mathcal{C}} \mu(c)c$. Since only hikes from v_i to v_j remain (by identification with the right-hand side), we deduce in view of (5),

$$(\operatorname{adj}(\mathbf{I} - \mathbf{W}))_{ij} = - \sum_{h \in \mathcal{H}_{ij}} \mu(h\omega_{ji})h = \sum_{h \in \mathcal{H}} \mu_{ij}(h)h. \quad (12)$$

Thus, $\operatorname{adj}(\mathbf{I} - \mathbf{W}) = \mathbf{M}$. □

The restriction $\mathbf{M}^{(\ell)}$ of \mathbf{M} to hikes of length ℓ satisfies, by identifying the terms of equal degrees in Lemma 2.2,

$$\mathbf{M}^{(\ell)} = \sum_{k=0}^{\ell} \psi_k \mathbf{W}^{\ell-k} = \left(\sum_{h \in \mathcal{H}^{\ell}} \mu_{ij}(h)h \right)_{i,j=1,\dots,N}. \quad (13)$$

In particular, $\mathbf{M}^{(\ell)}$ commutes with \mathbf{W} for all $\ell \geq 0$. If the digraph contains few self-avoiding hikes of length ℓ , the matrix $\mathbf{M}^{(\ell)}$ may have many zero entries. The construction gives in this case a non-trivial sparse matrix in the commutant of \mathbf{W} . This kind of problems has some applications in practice, for instance in the study of random processes. In [2], the authors investigate conditions under which the transition kernel of a finite state Markov chain observed at random times can be estimated consistently. They show that sparsity conditions, arising from particular state transitions known to be impossible, suffice to recover the transition kernel when it commutes with a certain matrix for which an estimator is available. The identifiability conditions in this model rely on the existence of a sparse matrix in the commutant. Similar problems and applications are studied in [12, 14, 16].

From the definition of μ_{ij} , one verifies easily that $\tilde{\mathbf{M}}^{(\ell)} := \psi_{\ell} \mathbf{I} - \mathbf{M}^{(\ell)}$ satisfies

$$\tilde{\mathbf{M}}_{ij}^{(\ell)} = - \sum_{k=0}^{\ell-1} \psi_k \mathbf{W}_{ij}^{\ell-k} = \sum_{h \in \mathcal{S}_{ij}^{\ell}} (-1)^{n(h)} h \quad (14)$$

for all $i, j = 1, \dots, N$. This shows that the matrix enumerating, up to the coefficient $(-1)^{n(h)}$, the self-avoiding hikes of length ℓ for all pairs of vertices can be obtained as a polynomial of \mathbf{W} . Because $\mu_{ij}(h)$ is trivially zero when $\ell(h) = N$, one recovers Cayley-Hamilton's theorem by setting $\ell = N$ in Equation (13). The general case also gives a direct proof of the identity

$$\psi_{\ell} = -\frac{1}{\ell} \sum_{k=0}^{\ell-1} \psi_k \operatorname{tr}(\mathbf{W}^{\ell-k}), \quad (15)$$

for which different proofs can be found in [19] and [11]. To prove it using Lemma 2.2, observe that (14) yields

$$\operatorname{tr}(\tilde{\mathbf{M}}^{(\ell)}) = \sum_{i=1}^N \sum_{c \in \mathcal{C} \cap \mathcal{S}_{ii}^{\ell}} (-1)^{n(c)} c = \ell \sum_{c \in \mathcal{C} \cap \mathcal{S}^{\ell}} (-1)^{n(c)} c = \ell \psi_{\ell},$$

noticing that each self-avoiding closed hike c appears exactly $\ell = \ell(c)$ times when summing over i . Hence, Equation (15) follows directly by computing the trace on both sides of the equality

$$\tilde{M}^{(\ell)} = - \sum_{k=0}^{\ell-1} \psi_k W^{\ell-k}.$$

3 Incidence algebra on hikes

Because \mathcal{C} is stable by multiplication, it forms a monoid with the empty cycle 1 as identity element. A natural partial order on \mathcal{C} arises from division: $d \in \mathcal{C}$ divides c , denoted by $d|c$, if there exists $c' \in \mathcal{C}$ such that $c = dc'$. The closed hikes ordered by division form a locally finite partially ordered set, or *poset*.

Definition 3.1 *The reduced incidence algebra of closed hike is the algebra of real valued functions on \mathcal{C} , endowed with the Dirichlet convolution, defined for $f, g : \mathcal{C} \rightarrow \mathbb{R}$ by*

$$f * g(c) = \sum_{d|c} f(d)g\left(\frac{c}{d}\right), \quad c \in \mathcal{C}.$$

In this definition, the sum is taken over all divisors $d \in \mathcal{C}$ of c , including the trivial cycle 1 and c itself. The reduced incidence algebra is isomorphic to the algebra of formal series, endowed with multiplication. Indeed, for two functions $f, g : \mathcal{C} \rightarrow \mathbb{R}$

$$\left(\sum_{c \in \mathcal{C}} f(c)c \right) \times \left(\sum_{c \in \mathcal{C}} g(c)c \right) = \sum_{c \in \mathcal{C}} f * g(c)c.$$

It follows that the Dirichlet convolution is associative, commutative and distributive over addition. The function δ defined on \mathcal{C} by $\delta(1) = 1$ and $\delta(c) = 0$ for all $c \neq 1$ is the identity element for this operation as we have, for any function f on \mathcal{C} , $f * \delta = \delta * f = f$. We refer to [18] for a more comprehensive study on this subject.

Extending this structure of open hikes is slightly more complicated. The set of hikes \mathcal{H} is not stable by multiplication and for this reason, defining a division over \mathcal{H} leads to some difficulties. However, hikes are stable by multiplication with a closed hike so that (\mathcal{H}, \cdot) forms an act over (\mathcal{C}, \cdot) . Thus, the division by a closed hike can be extended to open hikes: for $h \in \mathcal{H}$, d divides h if $d \in \mathcal{C}$ and $h = dh'$ for some $h' \in \mathcal{H}$. Similarly, the Dirichlet convolution can be extended to $f : \mathcal{C} \rightarrow \mathbb{R}$ and $g : \mathcal{H} \rightarrow \mathbb{R}$ by

$$f * g(h) = \sum_{d|h} f(d)g\left(\frac{h}{d}\right), \quad h \in \mathcal{H}.$$

Here again, the sum is taken only over closed divisors of h . Remark that f and g cannot be permuted in this expression, unless $h \in \mathcal{C}$. Interesting combinatorial properties arise from the poset structure of hikes, by considering each hike individually in Equation (8). The duality between walks and self-avoiding hikes becomes apparent in the next theorem. We extend the definition of f_{ij} to \mathcal{H} by setting $f_{ij}(h) = 0$ whenever h is not connected.

Theorem 3.2 For all $i, j = 1, \dots, N$, $\mu_{ij} = \mu * f_{ij}$.

Proof. From Lemma 2.2, we know that whenever $\|W\| < 1$,

$$M = \det(I - W) (I - W)^{-1}.$$

From (1) and (6) and using that $\det(I - W) = \sum_{c \in \mathcal{C}} \mu(c)c$, we obtain for all $i, j = 1, \dots, N$,

$$\sum_{h \in \mathcal{H}} \mu_{ij}(h)h = \sum_{c \in \mathcal{C}} \mu(c)c \times \sum_{h \in \mathcal{H}} f_{ij}(h)h = \sum_{h \in \mathcal{H}} \mu * f_{ij}(h) h.$$

The result follows by identification. \square

This theorem reveals a somewhat unexpected relation between the function μ_{ij} which only takes non zero values for self-avoiding hikes and f_{ij} whose support contains only walks. Taking a closer look, the result is not surprising if h is self-avoiding. Indeed, a self-avoiding hike h from v_i to v_j has exactly one divisor $d \in \mathcal{C}$ such that h/d is a walk from v_i to v_j . Thus, the convolution $\mu * f_{ij}$ is calculated over only one non-zero element and the equality is easily verified in this case. The result is actually more interesting if h is not self-avoiding as it yields in this case the non-trivial identity

$$\forall h \in \mathcal{H} \setminus \mathcal{S}, \sum_{d|h} \mu(d)f_{ij}\left(\frac{h}{d}\right) = 0.$$

Clearly, the function μ is a key feature to understand the combinatorial properties of this poset. The fact that $\mu(1) = 1 \neq 0$ makes it invertible through the Dirichlet convolution, and its inverse $\beta : \mathcal{C} \rightarrow \mathbb{R}$ is the unique function characterized by $\mu * \beta = \beta * \mu = \delta$. A reversed relation, expressing f_{ij} in function of μ_{ij} can then be derived easily, noticing that

$$f_{ij} = (\beta * \mu) * f_{ij} = \beta * (\mu * f_{ij}) = \beta * \mu_{ij}.$$

This relation turns out to be particularly important for our purposes, as we show that β satisfies interesting properties. In particular, we establish in the next proposition an expression of $\beta(c)$ that involves the number of appearances of each edge and vertex in c . Let $\tau_{ij}(c)$ denote the multiplicity of ω_{ij} in c and $\tau_i(c) = \sum_{j=1}^N \tau_{ij}(c)$ the number of edges starting from v_i in c (counted with multiplicity).

Theorem 3.3 The function β satisfies for all $c \in \mathcal{C}$,

$$\beta(c) = \prod_{i=1}^N \frac{\tau_i(c)!}{\tau_{i1}(c)! \times \dots \times \tau_{iN}(c)!}.$$

Proof. We will prove that the function β as defined in the theorem is the inverse of μ through the Dirichlet convolution. The case $c = 1$ being trivial, we take $c \neq 1$. Since $\mu(d) = 0$ if d is not self-avoiding, we have

$$\mu * \beta(c) = \sum_{d|c} \mu(d)\beta\left(\frac{c}{d}\right) = \sum_{\substack{d|c \\ d \in \mathcal{C} \setminus \mathcal{S}}} \mu(d)\beta\left(\frac{c}{d}\right).$$

For $d \neq 1$ a self-avoiding divisor of c , denote by σ_d the permutation over $V(d)$ associated to d , i.e. such that $d = \prod_{v_i \in V(d)} \omega_{i\sigma_d(i)}$. Clearly, $\tau_{ij}(c/d) = \tau_{ij}(c)$ if $v_i \notin V(d)$, while for $v_i \in V(d)$, we have $\tau_i(c/d) = \tau_i(c) - 1$ and

$$\tau_{ij}\left(\frac{c}{d}\right) = \begin{cases} \tau_{ij}(c) - 1 & \text{if } j = \sigma_d(i) \\ \tau_{ij}(c) & \text{otherwise.} \end{cases}$$

It follows

$$\beta\left(\frac{c}{d}\right) = \prod_{i=1}^N \frac{\tau_i(c/d)!}{\tau_{i1}(c/d)! \times \dots \times \tau_{iN}(c/d)!} = \prod_{v_i \in V(d)} \frac{\tau_{i\sigma_d(i)}(c)}{\tau_i(c)} \times \beta(c).$$

We get, including the case $d = 1$

$$\sum_{d|c} \mu(d) \beta\left(\frac{c}{d}\right) = \beta(c) \left(1 + \sum_{\substack{d|c \\ d \in \mathcal{C} \cap \mathcal{S} \setminus \{1\}}} \mu(d) \prod_{v_i \in V(d)} \frac{\tau_{i\sigma_d(i)}(c)}{\tau_i(c)} \right).$$

Recall that for $d \in \mathcal{C} \cap \mathcal{S}^k$, $\mu(d) = (-1)^k \text{sgn}(\sigma_d)$. Let $\ell = |V(c)|$ (the number of different vertices in c), the previous equality becomes, regrouping the divisors with equal lengths,

$$\sum_{d|c} \mu(d) \beta\left(\frac{c}{d}\right) = \beta(c) \left(1 + \sum_{k=1}^{\ell} (-1)^k \sum_{\substack{d|c \\ d \in \mathcal{C} \cap \mathcal{S}^k}} \text{sgn}(\sigma_d) \prod_{v_i \in V(d)} \frac{\tau_{i\sigma_d(i)}(c)}{\tau_i(c)} \right).$$

Now consider the $\ell \times \ell$ matrix $\mathbf{B}(c)$ with entries $\tau_{ij}(c)/\tau_i(c)$ for $v_i, v_j \in V(c)$. By identifying each self-avoiding divisor d of c with its corresponding permutation σ_d , we recognize in the above expression the characteristic polynomial of $\mathbf{B}(c)$ taken at $\lambda = 1$,

$$1 + \sum_{k=1}^{\ell} (-1)^k \sum_{\substack{d|c \\ d \in \mathcal{C} \cap \mathcal{S}^k}} \text{sgn}(\sigma_d) \prod_{i \in d} \frac{\tau_{i\sigma_d(i)}(c)}{\tau_i(c)} = \det(\mathbf{I} - \mathbf{B}(c)).$$

Since $\mathbf{B}(c)$ is a stochastic matrix, $\det(\mathbf{I} - \mathbf{B}(c)) = 0$ which ends the proof. \square

The coefficient $\beta(c)$ corresponds to the number of arrangements of the edges in c , regrouped by their initiating vertex. Indeed, the multinomial coefficient

$$\frac{\tau_i(c)!}{\tau_{i1}(c)! \times \dots \times \tau_{iN}(c)!}$$

counts the ways of ordering the edges initiating from v_i in c , accounting for their multiplicity $\tau_{ij}(c)$. Considering all configurations for each vertex in c recovers the coefficient $\beta(c)$. So, β enumerates the different ways to travel along a closed hike.

This result points out some interesting properties of β , most of which are not straightforward from its initial definition as the inverse of μ . The first immediate consequence is that β is positive. Secondly, $\beta(c)$ is equal to one if c is a self-avoiding closed hike. This condition is sufficient but not necessary, as we have for instance $\beta(c^2) = 1$ as soon as $\beta(c) = 1$. A third consequence is that β is non-decreasing with respect to multiplication, which can be stated formally as: $\forall c_1, c_2 \in \mathcal{C}$, $\beta(c_1 c_2) \geq \max\{\beta(c_1), \beta(c_2)\}$. Finally, β is multiplicative over decompositions on disjoint closed hikes. Indeed, if c can be written as the product of say $p \geq 2$ mutually vertex-disjoint components $c_1, \dots, c_p \in \mathcal{C}$, then $\beta(c) = \beta(c_1) \dots \beta(c_p)$. This property is reminiscent of the multiplicity of arithmetic functions over coprime integers (see for instance [1]). In this framework, two closed hikes c_1, c_2 can be considered coprime if they share no common vertex. The multiplicity property of β is then inherited from the multiplicity of its inverse μ .

Remark. The function μ is the Mobius function (i.e. the inverse of the constant function equal to 1) of the trace monoid of circuits described in [6]. Circuits correspond to partially commutative versions of closed hike so that, in our fully commutative framework, circuits composed of the same edges are seen as the same object. Thus, $\beta(c)$ counts the number of circuits composed of the same edges as c .

A different expression for β can be derived from the inverse relation in Lemma 2.2, writing W^ℓ in terms of the $M^{(k)}$, $k = 0, 1, \dots, \ell$. This result is given as a corollary.

Corollary 3.4 *If $\|W\| < 1$, then for $\ell \in \mathbb{N}$,*

$$W^\ell = \sum_{k=0}^{\ell} \phi_k M^{(\ell-k)}, \quad (16)$$

where the coefficients $\phi_0, \phi_1, \dots, \phi_N$ are defined by

$$\phi_0 = 1, \quad \phi_k = \sum_{k_1 + \dots + k_p = k} (-1)^p \psi_{k_1} \dots \psi_{k_p}, \quad k = 1, \dots, N. \quad (17)$$

Before proving the result, let us clarify that the ϕ_k 's are defined by taking the sum over all compositions of k , that is, all positive tuples (k_1, \dots, k_p) such that $k_1 + \dots + k_p = k$, for all $p = 1, \dots, k$ (two tuples composed of the same integers k_1, \dots, k_p but in different orders are to be counted twice).

Proof. This holds if W is the null matrix with the convention $W^0 = I$ in this case. For $W \neq 0$, Equation (10) yields

$$M = \sum_{\ell \geq 0} M^{(\ell)} = \sum_{k=0}^N \psi_k \times \sum_{k \geq 0} W^k \iff \sum_{\ell \geq 0} W^\ell = \frac{1}{\sum_{k=0}^N \psi_k} \sum_{k \geq 0} M^{(k)}.$$

We use the formal series expansion

$$\frac{1}{\sum_{k=0}^N \psi_k} = \frac{1}{1 + \sum_{k=1}^N \psi_k} = \sum_{p \geq 0} (-1)^p \left(\sum_{k=1}^N \psi_k \right)^p.$$

By regrouping the terms of equal degree, we get

$$\frac{1}{\sum_{k=0}^N \psi_k} = 1 + \sum_{k \geq 1} \sum_{k_1 + \dots + k_p = k} (-1)^p \psi_{k_1} \dots \psi_{k_p} = \sum_{k \geq 0} \phi_k. \quad (18)$$

Hence,

$$\sum_{\ell \geq 0} \mathbb{W}^\ell = \sum_{k \geq 0} \phi_k \times \sum_{k \geq 0} \mathbb{M}^{(k)} = \sum_{\ell \geq 0} \sum_{k=0}^{\ell} \phi_k \mathbb{M}^{(\ell-k)},$$

and the result follows by identification. \square

Like the ψ_k 's, the coefficients ϕ_k are homogenous polynomials of degree k in the ω_{ij} 's. While ψ_k only involves the self-avoiding closed hikes, ϕ_k depends on all the closed hikes of length k on the digraph. The formal series inversion in Equation (18) actually corresponds to the inversion of the Dirichlet convolution when identifying each closed hike. This means in particular that the coefficient ϕ_k can be expressed as

$$\phi_k = \sum_{c \in \mathcal{C}^k} \beta(c)c. \quad (19)$$

One can verify this formula directly from the formal series multiplication

$$1 = \sum_{k \geq 0} \phi_k \times \sum_{k \geq 0} \psi_k = \sum_{c \in \mathcal{C}} \beta(c)c \times \sum_{c \in \mathcal{C}} \mu(c)c = \sum_{c \in \mathcal{C}} \beta * \mu(c)c$$

recovering exactly the formal series version of the equality $\delta = \beta * \mu$. By combining Equations (17) and (19), we deduce a new expression of $\beta(c)$ for $c \neq 1$:

$$\beta(c) = \sum_{p \geq 1} \sum_{c_1 \dots c_p = c} (-1)^p \mu(c_1) \dots \mu(c_p) = \sum_{s_1 \dots s_p = c} (-1)^{n'(s_1) + \dots + n'(s_p)} \quad (20)$$

setting $n'(\cdot) = n(\cdot) + 1$, where the final sum is taken over all p -tuples (s_1, \dots, s_p) of non-empty self-avoiding closed hikes such that $s_1 \dots s_p = c$. This equality provides an expression of $\beta(c)$ involving the different decompositions of c into self-avoiding closed hikes. While this expression is presumably less practical than the previous one, it induces nevertheless interesting consequences from a combinatorial point of view, which are discussed in Section 4.

We now come to our final result, which expresses the multiplicity of an open walk in terms of its decompositions into self-avoiding components. This result will be illustrated on some examples in Section 4.

Theorem 3.5 *Let h be a non-empty hike from v_i to v_j ,*

$$f_{ij}(h) = \sum_{s_1 \dots s_k p = h} (-1)^{n'(s_1) + \dots + n'(s_k) + n'(p)}$$

setting $n'(\cdot) = n(\cdot) + 1$, where the sum is taken over all self-avoiding decompositions of h , i.e., all $(k+1)$ -tuples $(s_1, \dots, s_k, p) \in (\mathcal{C} \cap \mathcal{S} \setminus \{1\})^k \times \mathcal{S}_{ij}$ with $k \geq 0$ such that $s_1 \dots s_k p = h$.

Proof. Plug the expression of β in Equation (20) into $f_{ij}(h) = \sum_{d|h} \beta(d) \mu_{ij}(h/d)$. For $i \neq j$, the fact that $\mu_{ij}(h/d) = 0$ for $h/d \notin \mathcal{S}_{ij}$ simplifies into

$$f_{ij}(h) = \sum_{\substack{d=s_1 \dots s_k | h \\ h/d \in \mathcal{S}_{ij}}} (-1)^{n'(s_1) + \dots + n'(s_k)} \times (-1)^{n'(h/d)} = \sum_{s_1 \dots s_k p = h} (-1)^{n'(s_1) + \dots + n'(s_k) + n'(p)}$$

for $p = h/d$, thus recovering the result. For $i = j$, we use that $\mu_{ii}(c) = \mu(c) \mathbf{1}\{v_i \notin V(c)\}$,

$$f_{ii}(h) = \sum_{d|h} \beta(d) \mu_{ii}\left(\frac{h}{d}\right) = \sum_{d|h} \beta(d) \mu\left(\frac{h}{d}\right) - \sum_{\substack{d|h \\ v_i \in V(h/d)}} \beta(d) \mu\left(\frac{h}{d}\right).$$

The first term of the right-hand side is $-\beta * \mu(h)$ which is zero for all $h \neq 1$. The result follows by using the expression of β given in Equation (20), similarly as in the case $i \neq j$. \square

4 Examples

In this section, the functions f_{ij}, μ_{ij}, β and μ are computed on some examples. For ease of comprehension, we start with explicit simple cases before considering more general structures in the final examples.

Example 1. Let us begin with the graph represented in Figure 4 which contains only two disjoint simple cycles.




Figure 4: Disjoint simple cycles.

The hike covering the whole graph is the closed hike $h_1 = \omega_{12}\omega_{23}\omega_{34}\omega_{41}\omega_{56}\omega_{67}\omega_{75}$ (recall that the order is not important). We obtain directly $f_{ij}(h_1) = 0$ (because h_1 is not connected) and $\mu_{ij}(h_1) = 0$ (because h_1 crosses every vertex) for all $i, j = 1, \dots, 7$. Moreover, the definitions of μ and β give in this case $\mu(h_1) = \beta(h_1) = 1$. To check the equalities $\mu * f_{ii} = \mu_{ii}$ and $\beta * \mu_{ii} = f_{ii}$, the calculations are straightforward, since the only closed divisors of h_1 are $h_1, c_1 = \omega_{12}\omega_{23}\omega_{34}\omega_{41}, c_2 = \omega_{56}\omega_{67}\omega_{75}$ and the void cycle 1. We get for instance,

$$\mu * f_{11}(h_1) = \mu(h_1) f_{11}(1) + \mu(c_2) f_{11}(c_1) = 0 = \mu_{11}(h_1),$$

using that $f_{11}(c_1) = 1, f_{11}(c_2) = 0$ and $\mu(c_2) = -1$. From $\mu_{11}(c_1) = 0$, we also verify

$$\beta * \mu_{11}(h_1) = \beta(h_1) \mu_{11}(1) + \beta(c_1) \mu_{11}(c_2) = 0 = f_{11}(h_1).$$

Example 2. We now consider the graph given in Figure 5, composed of two simple cycles sharing one vertex and the covering hike $h_2 = \omega_{12}\omega_{23}\omega_{31}\omega_{24}\omega_{45}\omega_{52}$.




Figure 5: Two simple cycles with one common vertex.

Since h_2 is closed, $f_{ij}(h_2)$ is null for all $i \neq j$. Moreover, there are two ways to travel across h_2 starting from v_2 , depending on which side is visited first, and one way for every other vertex. We deduce $f_{22}(h_2) = 2$ and $f_{ii}(h_2) = 1$ for $i = 1, 3, 4, 5$. Since h_2 is not self-avoiding $\mu(h_2) = 0$ and Theorem 3.3 gives $\beta(h_2) = 2$. To check the formulas, we now consider all the decompositions of h_2 into a product of closed hikes. The two non-trivial divisors of h_2 are $c_1 = \omega_{12}\omega_{23}\omega_{31}$ and $c_2 = \omega_{24}\omega_{45}\omega_{52}$. We verify for instance,

$$\begin{aligned}\mu * f_{22}(h_2) &= \mu(1)f_{22}(h_2) + \mu(c_2)f_{22}(c_1) + \mu(c_1)f_{22}(c_2) = 0 = \mu_{22}(h_2) \\ \beta * \mu_{11}(h_2) &= \beta(h_2)\mu_{11}(1) + \beta(c_1)\mu_{11}(c_2) = 1 = f_{11}(h_2)\end{aligned}$$

Example 3. This example deals with the walk $h_3 = \omega_{12}\omega_{23}\omega_{34}\omega_{41}\omega_{62}\omega_{25}\omega_{54}\omega_{46}$ composed of two simple cycles sharing two vertices, represented in Figure 6. The non-trivial divisors of h_3 are detailed in Figure 7.




Figure 6: Two simple cycles with two common vertices.

Clearly, $f_{ij}(h_3) = 0$ for $i \neq j$. We compute the values $f_{ii}(h_3)$ by enumerating all ways to travel across h_3 starting from v_i . For instance, the two ways from v_1 are

$$\begin{aligned}1 &\rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 6 \rightarrow 2 \rightarrow 5 \rightarrow 4 \rightarrow 1 \\ 1 &\rightarrow 2 \rightarrow 5 \rightarrow 4 \rightarrow 6 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 1\end{aligned}$$

We find $f_{11}(h_3) = f_{33}(h_3) = f_{55}(h_3) = f_{66}(h_3) = 2$ and $f_{22}(h_3) = f_{44}(h_3) = 4$. Here again, $\mu_{ij}(h_3)$ is null for all $i, j = 1, \dots, 6$ as well as $\mu(h_3)$, while $\beta(h_3) = 4$. We verify the convolution

equalities $\mu_{ii}(h_3) = \mu * f_{ii}(h_3)$ and $f_{ii}(h_3) = \beta * \mu_{ii}(h_3)$ for arbitrary vertices, e.g.

$$\begin{aligned} \mu * f_{11}(h_3) &= \mu(1)f_{11}(h_3) + \mu(c_2)f_{11}(c_1) + \mu(c_3)f_{11}(c_4) = 0 = \mu_{11}(h_3) \\ \beta * \mu_{22}(h_3) &= \beta(h_3)\mu_{22}(1) = 4 = f_{22}(h_3) \end{aligned}$$




Figure 7: Simple divisors of h_3 : $c_1 = \omega_{12}\omega_{23}\omega_{34}\omega_{41}$ (top-left), $c_2 = \omega_{25}\omega_{54}\omega_{46}\omega_{62}$ (top-right), $c_3 = \omega_{12}\omega_{25}\omega_{54}\omega_{41}$ (bottom-left) and $c_4 = \omega_{23}\omega_{34}\omega_{46}\omega_{62}$ (bottom-right).

Example 4. We consider the closed hike $h_4 = \omega_{12}\omega_{23}\omega_{35}\omega_{56}\omega_{64}\omega_{41}\omega_{25}\omega_{54}\omega_{42}$, illustrated in Figure 8, composed of 2 cycles sharing 3 vertices. In this case, note that the orientation of the two cycles has an impact on the values of f_{ii} .




Figure 8: Two simple cycles with three common vertices.

The walk h_4 has 8 non-trivial divisors, detailed in Figure 9.




Figure 9: Simple divisors of h_4 . From top-left to bottom-right: $c_1 = \omega_{25}\omega_{54}\omega_{42}$, $c_2 = \omega_{12}\omega_{23}\omega_{35}\omega_{56}\omega_{64}\omega_{41}$, $c_3 = \omega_{12}\omega_{25}\omega_{54}\omega_{41}$, $c_4 = \omega_{23}\omega_{35}\omega_{56}\omega_{64}\omega_{42}$, $c_5 = \omega_{25}\omega_{56}\omega_{64}\omega_{42}$, $c_6 = \omega_{12}\omega_{23}\omega_{35}\omega_{54}\omega_{41}$, $c_7 = \omega_{23}\omega_{35}\omega_{54}\omega_{42}$ and $c_8 = \omega_{12}\omega_{25}\omega_{56}\omega_{64}\omega_{41}$.

We know that $f_{ij}(h_4) = 0$ for $i \neq j$, $\mu_{ij}(h_4) = 0$ for all $i, j = 1, \dots, 6$, $\mu(h_4) = 0$ and $\beta(h_4) = 8$. Counting the connected paths starting from each vertex gives $f_{11}(h_4) = f_{33}(h_4) = f_{66}(h_4) = 4$ and $f_{22}(h_4) = f_{44}(h_4) = f_{55}(h_4) = 8$. We recover the correct values from the identities $\mu_{ii}(h_4) = \mu * f_{ii}(h_4)$ and $f_{ii}(h_4) = \beta * \mu_{ii}(h_4)$. Keeping only the non-zero values in the convolution gives,

for instance

$$\begin{aligned}
\mu_{22}(h_4) &= \mu(1)f_{22}(h_4) + \mu(c_2)f_{22}(c_1) + \mu(c_1)f_{22}(c_2) + \mu(c_4)f_{22}(c_3) + \mu(c_3)f_{22}(c_4) \\
&\quad + \mu(c_6)f_{22}(c_5) + \mu(c_5)f_{22}(c_6) + \mu(c_8)f_{22}(c_7) + \mu(c_7)f_{22}(c_8) \\
&= 8 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 = 0 \\
f_{11}(h_4) &= \beta(h_4)\mu_{11}(1) + \beta(c_2)\mu_{11}(c_1) + \beta(c_3)\mu_{11}(c_4) + \beta(c_6)\mu_{11}(c_5) + \beta(c_8)\mu_{11}(c_7) \\
&= 8 - 1 - 1 - 1 - 1 = 4.
\end{aligned}$$

Example 5. Consider a closed hike h_5 composed of two simple cycles of opposite directions crossing n times. This example can be represented as n cycles placed one after the other. As we observed in the previous examples, the length of these cycles does not impact the values of the functions f_{ii}, μ_{ii}, μ and β so that we can take cycles of length 2 without loss of generality, considering for instance the closed hike $\omega_{12}\omega_{21}\omega_{23}\omega_{32}\cdots\omega_{n1}\omega_{1n}$ illustrated in Figure 10.




Figure 10: Two simple cycles with $n = 8$ common vertices.

We find $f_{ij}(h_5) = 0$ for $i \neq j$, $f_{ii}(h_5) = 2n$, $\mu_{ij}(h_5) = 0$ for all $i, j = 1, \dots, n$, $\mu(h_5) = 0$ and $\beta(h_5) = 2^n$. Let $a_k = \omega_{kk+1}\omega_{k+1k}$, $k = 1, \dots, n-1$, $a_n = \omega_{n1}\omega_{1n}$ be the length 2 divisors of h_5 and $d_0 = \omega_{12}\omega_{23}\cdots\omega_{n-1n}\omega_{n1}$, $d_1 = \omega_{21}\omega_{32}\cdots\omega_{nn-1}\omega_{1n}$ the outside and inside cycles composing h_5 . The non-trivial divisors of h_5 are d_0, d_1 and every product $a_{k_1}\cdots a_{k_p}$ obtained for a subset $\{k_1, \dots, k_p\} \subset \{1, \dots, n\}$. Using that f_{11} is non-zero only for walks passing through v_1 and μ vanishes for non self-avoiding closed hikes, we obtain by keeping only the non-zero terms in the Dirichlet convolution

$$\mu * f_{11}(h_5) = \mu(1)f_{11}(h_5) + \mu(d_1)f_{11}(d_0) + \mu(d_0)f_{11}(d_1) + \sum_{k=2}^n \mu(a_k)f_{11}\left(\frac{h_5}{a_k}\right)$$

which recovers ultimately $\mu * f_{11}(h_5) = 2n - 2 - 2(n-1) = 0 = \mu_{11}(h_5)$. The reverse relation $f_{11}(h_5) = \beta * \mu_{11}(h_5)$ is less trivial. To compute it, we have to enumerate for any p , the sets $\{k_1, \dots, k_p\} \subset \{1, \dots, n\}$ such that $\mu_{11}(a_{k_1}\cdots a_{k_p}) \neq 0$, i.e., such that $a_{k_1}\cdots a_{k_p}$ is a self-avoiding closed hike that does not cross v_1 . For each such closed hike, the complement $h_5/(a_{k_1}\cdots a_{k_p})$ is composed of p disjoint connected components, one of which contains a_1a_n . This component

can be divided into two connected components by separating a_1 and a_n . Thus, each closed hike $a_{k_1} \cdots a_{k_p}$ such that $\mu_{11}(a_{k_1} \cdots a_{k_p}) \neq 0$ can be associated with a composition of $n - p$ containing $p + 1$ elements, which there are $\binom{n-p-1}{p}$ of them. For each $a_{k_1} \cdots a_{k_p}$, the coefficient β of the complement equals 2^{n-2p} and one recovers the formula

$$\beta * \mu_{11}(h_5) = \sum_{p=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-p-1}{p} (-1)^p 2^{n-2p} = 2n = f_{11}(h_5).$$

Example 6. Consider a self-avoiding closed hike h_6 composed of $n \geq 2$ simple connected components a_1, \dots, a_n (we may assume without loss of generality that each connected component is a loop as illustrated in Figure 11).




Figure 11: Illustration of the hikes considered in Examples 6 (left) and 7 (right) for $n = 8$.

From the first expression of β given in Theorem 3.3, it is clear that $\beta(h_6) = 1$. On the other hand, the number of ways to decompose h_6 into a product of $k \leq n$ non-empty self-avoiding closed hikes writes as the sum of the multinomial coefficients over all positive compositions n_1, \dots, n_k of n . Since for any self-avoiding decomposition s_1, \dots, s_k , the product $\mu(s_1) \dots \mu(s_k)$ always equals $\mu(h_6) = (-1)^n$, combining the two expressions of $\beta(h_6)$ yields

$$\beta(h_6) = 1 = (-1)^n \sum_{k=1}^n (-1)^k \sum_{n_1 + \dots + n_k = n} \frac{n!}{n_1! \dots n_k!}.$$

Alternatively, this equality can be obtained by identifying the coefficient of x^k in the power series expansions of $e^{-x} = 1/e^x$. Since h_6 is not connected, we know that $f_{ii}(h_6) = 0$ for all i . To compute the expression of $f_{ii}(h_6)$ from Theorem 3.5, we consider the self-avoiding decompositions of the form $h_6 = s_1 \dots s_k p$ with $s_1, \dots, s_k \in \mathcal{C} \cap \mathcal{S} \setminus \{1\}$ and $p \in \mathcal{S}_{ii}$. To verify that this expression gives $f_{ii}(h_6) = 0$ in this case simply observe that any self-avoiding decomposition s_1, \dots, s_k, p such that $p \neq \omega_{ii}$ cancels out with the decomposition $s_1, \dots, s_k, p/\omega_{ii}, \omega_{ii}$ in view of

$$(-1)^{n'(s_1) + \dots + n'(s_k) + n'(w)} = -(-1)^{n'(s_1) + \dots + n'(s_k) + n'(p/\omega_{ii}) + n'(\omega_{ii})}.$$

Thus, summing over all self-avoiding decompositions recovers $f_{ii}(h_6) = 0$.

Example 7. We now consider the closed hike h_7 constructed from the previous example with an extra cycle c_0 passing through each vertex: $h_7 = h_6 \times c_0$ (e.g. in the right graph in Figure 11 where $c_0 = \omega_{12}\omega_{23}\omega_{34}\omega_{45}\omega_{56}\omega_{67}\omega_{78}\omega_{81}$). In this example, the cycle c_0 is isolated in every self-avoiding decomposition since it shares a common node with all the other divisors of h_7 . Thus, the different ways to express h_7 as a product of self-avoiding closed hikes can be obtained from the previous example, inserting the cycle c_0 wherever possible. Precisely, for a decomposition $h_6 = s_1 \dots s_k$ of h_6 into $k \leq n$ non-empty self-avoiding closed hikes, there are exactly $k + 1$ possibilities to insert c_0 . Moreover, remark that $\mu(s_1) \dots \mu(s_k) \mu(c_0) = \mu(h_6) \mu(c_0) = (-1)^{n+1}$ is constant over all self-avoiding decompositions. Combining the two expressions of $\beta(h_7)$ thus recovers the formula

$$\sum_{k=1}^n (-1)^{k+1} (k+1) \sum_{n_1 + \dots + n_k = n} (-1)^{n+1} \frac{n!}{n_1! \dots n_k!} = 2^n.$$

In this example, there are two ways of visiting the whole walk h_7 from one vertex v_i to itself, depending on whether the loop at v_i is traveled at the start or at the end. Thus, $f_{ii}(h_7) = 2$ for all v_i . In a self-avoiding decomposition with $s_1, \dots, s_k \in \mathcal{C} \cap \mathcal{S} \setminus \{1\}$ and $p \in \mathcal{S}_{ii}$ we can distinguish the cases $p = c_0$, $p = \omega_{ii}$ and $p \neq \omega_{ii}, c_0$. Clearly, the sum over all self-avoiding decompositions $h_7 = s_1 \dots s_k p$ such that $p = c_0$ yields $\beta(h_6)$ since $(-1)^{n'(c_0)} = 1$. Moreover, the sum over all self-avoiding decompositions with $p = \omega_{ii}$ recovers $\beta(h_7/\omega_{ii}) = 2^{n-1}$. Finally, for a self-avoiding decomposition $h_6 = s_1 \dots s_k p$ of h_6 with $p \neq \omega_{ii}$, there are k possibilities to insert c_0 , yielding

$$f_{ii}(h_7) = \beta(h_6) + \beta\left(\frac{h_7}{\omega_{ii}}\right) + \sum_{k=1}^{n-1} (-1)^{k+1} k \sum_{n_1 + \dots + n_k = n-1} (-1)^{n+1} \frac{(n-1)!}{n_1! \dots n_k!} = 1 + 2^{n-1} - 2^{n-1} + 1 = 2.$$

Acknowledgments

The authors are grateful to Pierre-Louis Giscard for his explanations on the poset structure of hikes, and to an anonymous referee for its helpful comments which helped improve this paper.

References

- [1] Tom M Apostol. Modular functions and Dirichlet series in number theory. *AMC*, 10:12, 1990.
- [2] Flavia Barsotti, Yohann De Castro, Thibault Espinasse, and Paul Rochet. Estimating the transition matrix of a Markov chain observed at random times. *Statistics & Probability Letters*, 94:98–105, 2014.
- [3] Philippe Blanchard and Dimitri Volchenkov. *Random Walks and Diffusions on Graphs and Databases: An Introduction*, volume 10. Springer Science & Business Media, 2011.

- [4] Raffaella Burioni and Davide Cassi. Random walks on graphs: ideas, techniques and results. *Journal of Physics A: Mathematical and General*, 38(8):R45, 2005.
- [5] Peter J Carrington, John Scott, and Stanley Wasserman. *Models and methods in social network analysis*, volume 28. Cambridge university press, 2005.
- [6] Pierre Cartier and Dominique Foata. Problemes combinatoires de commutation et réarrangements. *Lecture notes in mathematics*, 85, 1969.
- [7] Fan RK Chung. *Spectral graph theory*, volume 92. American Mathematical Soc., 1997.
- [8] Dragoš M Cvetković. *Eigenspaces of graphs*. Number 66. Cambridge University Press, 1997.
- [9] Xingguang Han, Yangzhou Chen, Jianjun Shi, and Zhonghe He. An extended cell transmission model based on digraph for urban traffic road network. In *Intelligent Transportation Systems (ITSC), 2012 15th International IEEE Conference on*, pages 558–563. IEEE, 2012.
- [10] Frank Harary. The determinant of the adjacency matrix of a graph. *Siam Review*, 4(3):202–210, 1962.
- [11] Zhang Hong-Hao, Yan Wen-Bin, and Li Xue-Song. Trace formulae of characteristic polynomial and cayley–hamilton’s theorem, and applications to chiral perturbation theory and general relativity. *Communications in Theoretical Physics*, 49(4):801, 2008.
- [12] Yasunari Inamura. Estimating continuous time transition matrices from discretely observed data. *Bank of Japan Working Paper Series*, (06), 2006.
- [13] Gregory F Lawler. Loop-erased self-avoiding random walk and the laplacian random walk. *Journal of Physics A: Mathematical and General*, 20(13):4565, 1987.
- [14] Elizabeth Chase MacRae. Estimation of time-varying Markov processes with aggregate data. *Econometrica*, 45(1):183–198, 1977.
- [15] John S Maybee, DD Olesky, Driessche P van den, and G Wiener. Matrices, digraphs, and determinants. *SIAM Journal on Matrix Analysis and Applications*, 10(4):500–519, 1989.
- [16] A. O. Pittenger. Time changes of Markov chains. *Stochastic Process. Appl.*, 13(2):189–199, 1982.
- [17] J Ponstein. Self-avoiding paths and the adjacency matrix of a graph. *SIAM Journal on Applied Mathematics*, 14(3):600–609, 1966.
- [18] Gian-Carlo Rota. On the foundations of combinatorial theory. In *Classic Papers in Combinatorics*, pages 332–360. Springer, 1987.
- [19] Lotfi Asker Zadeh and Charles A Deoser. *Linear system theory*. Robert E. Krieger Publishing Company, 1976.