

CONCENTRATION AND NON-CONCENTRATION FOR THE SCHRÖDINGER EVOLUTION ON ZOLL MANIFOLDS

FABRICIO MACIÀ AND GABRIEL RIVIÈRE

ABSTRACT. We study the long time dynamics of the Schrödinger equation on Zoll manifolds. We establish criteria under which the solutions of the Schrödinger equation can or cannot concentrate on a given closed geodesic. As an application, we derive some results on the set of semiclassical measures for eigenfunctions of Schrödinger operators: we prove that adding a potential $V \in C^\infty(\mathbb{S}^d)$ to the Laplacian on the sphere results on the existence of geodesics γ such that the uniform measure supported on γ cannot be obtained as a weak- \star accumulation point of the densities $(|\psi_n|^2 \text{vol}_{\mathbb{S}^d})$ for any sequence of eigenfunctions (ψ_n) of $\Delta_{\mathbb{S}^d} - V$. We also show that the same phenomenon occurs for the free Laplacian on certain Zoll surfaces.

1. INTRODUCTION

In this article we are interested in understanding the dynamics of Schrödinger equations and the structure of eigenfunctions of Schrödinger operators on Zoll manifolds. Recall that a *Zoll manifold* is a smooth, connected, compact, Riemannian manifold without boundary (M, g) such that *all its geodesics are closed*. This means that, for every x in M , all the geodesics issued from x come back to x . Thanks to Theorem of Wadsley (see [4] – section 7.B), the geodesic flow φ^s acting on the unit cotangent bundle S^*M of such a manifold is periodic, meaning that all its trajectories have a minimal common period $l > 0$.¹ Using the terminology of [4] – chapter 7, we will say that the metric g is a P_l -metric, or that (M, g) is a P_l -manifold. Similarly, in the case where all the geodesics have the same length l , we will say that g is a C_l -metric or that (M, g) is a C_l -manifold. The main examples of C_l -manifolds are the compact rank one symmetric spaces (that we will sometimes abbreviate by CROSS) and certain surfaces of revolution – see chapters 3 and 4 in [4]. It is also known that the set of C_l -metrics on \mathbb{S}^2 that can be obtained as smooth deformations of the canonical metric is infinite dimensional and contains many metrics that are not of revolution. The characterization of such deformations is a remarkable result by Guillemin [22].

FM takes part into the visiting faculty program of ICMAT and is partially supported by grants ERC Starting Grant 277778 and MTM2013-41780-P (MEC).

GR is partially supported by the Agence Nationale de la Recherche through the Labex CEMPI (ANR-11-LABX-0007-01) and the ANR project GeRaSic (ANR-13-BS01- 0007-01).

¹Note that this does not mean that all the geodesics have length equal to l . There may exist exceptional geodesics whose length is strictly smaller than l , the Lens spaces (quotients of \mathbb{S}^{2m-1} by certain finite cyclic groups of isometries) provide an example of this – see Ex. 2.43 in [27].

Our goal here is to understand the long time dynamics of the following Schrödinger equation:

$$(1) \quad i\partial_t v(t, x) = \left(-\frac{1}{2}\Delta_g + V(x) \right) v(t, x), \quad v|_{t=0} = u \in L^2(M),$$

or the behavior of eigenfunctions:

$$(2) \quad \left(-\frac{1}{2}\Delta_g + V(x) \right) u(x) = \lambda^2 u(x), \quad \|u\|_{L^2(M)} = 1,$$

in the high-frequency limit $\lambda \rightarrow \infty$. As usual, Δ_g is the Laplace Beltrami operator induced by the Riemannian metric g on M , and we shall assume that V is in $\mathcal{C}^\infty(M, \mathbb{R})$.

The study of the spectral properties of the operator $-\frac{1}{2}\Delta_g + V$ in this geometric context is a problem which has a long history in microlocal analysis starting with the works of Duistermaat-Guillemin [14, 23, 24], Weinstein [51] and Colin de Verdière [9]. Many other important results on the fine structure of the spectrum of Zoll manifolds were obtained both in the microlocal framework [25, 47, 48, 53, 54], and in the semiclassical setting [8, 28, 26] – see also [30, 31] in the nonselfadjoint setting.

In this this article we use similar techniques to those that were originally developed in order to study of the spectrum of Schrödinger operators to actually provide some results on the long-time dynamics of the Schrödinger evolution (1) and the structure of high-frequency eigenfunctions (2).

1.1. Concentration and non-concentration of eigenfunctions. Let us start by describing our results in the context of eigenfunctions of Schrödinger operators, as they are somewhat simpler to state. We are mainly interested in analyzing how the mass $|u|^2$ of a high-frequency eigenfunction satisfying (2) distributes over M . More precisely, consider the set $\mathcal{N}(\infty)$ of probability measures in M that are obtained as follows. A probability measure ν belongs to $\mathcal{N}(\infty)$ provided there exist a sequence of eigenfunctions (u_n) :

$$-\frac{1}{2}\Delta_g u_n + V u_n = \lambda_n^2 u_n, \quad \|u_n\|_{L^2(M)} = 1,$$

with eigenvalues satisfying $\lambda_n \rightarrow \infty$ such that

$$\lim_{n \rightarrow \infty} \int_M a(x) |u_n|^2(x) \text{vol}_g(dx) = \int_M a(x) \nu(dx), \quad \text{for every } a \in \mathcal{C}(M).$$

Measures in $\mathcal{N}(\infty)$ therefore describe the asymptotic mass distribution sequences of eigenfunctions (u_n) whose corresponding eigenvalues tend to infinity. The problem of characterizing the probability measures in $\mathcal{N}(\infty)$ has attracted a lot of attention in the last forty years.

In the case of Zoll manifolds, it is well known that $\mathcal{N}(\infty)$ is contained in \mathcal{N}_g which is, by definition, the closed convex hull (with respect to the weak- \star topology) of the set of probability measures δ_γ , where γ is a geodesic of (M, g) . Recall that

$$\int_M a(x) \delta_\gamma(dx) = \frac{1}{l} \int_0^l a(\gamma(s)) ds,$$

where l denotes the length of γ and the parametrization $\gamma(s)$ has unit speed.

In the case of \mathbb{S}^2 endowed with its canonical metric, it was proved that a “generic” orthonormal basis of eigenfunctions satisfies the quantum unique ergodicity property [52, 49]. In particular, in this case, the normalized Riemannian volume is the only accumulation point of the sequence of eigenfunctions under consideration. Note that quantum ergodicity properties were also proved recently for sequences of eigenfunctions on \mathbb{S}^d satisfying certain symmetry assumptions [36, 7]. Yet, there are in fact some situations for which one has

$$\mathcal{N}(\infty) = \mathcal{N}_g,$$

which is in a certain sense the opposite situation to quantum ergodicity. Jakobson and Zelditch proved in [35] that this holds when (M, g) is the sphere \mathbb{S}^d equipped with its canonical metric $g = \text{can}$ and the potential V vanishes identically. This was also shown to hold for general Compact Rank-One Symmetric Spaces (see [37]) and for any manifold of positive constant curvature (see [3] – this analysis relies on the study of eigenfunctions of the canonical Laplacian on the sphere that are invariant by certain groups of isometries); in both cases one has to assume that $V = 0$. To the authors’ knowledge, the question of whether this is always the case when the potential V does not vanish identically or when (M, g) is a Zoll manifold that is not isometric to a CROSS remains open.

Here we answer these questions by the negative. It turns out that these regimes are somewhat intermediate between the quantum ergodicity results and the results on a CROSS. Let us first introduce some notations. Denote by \mathring{T}^*M the cotangent bundle of M with the zero section removed. The Radon transform of a functions $b \in \mathcal{C}^\infty(T^*M)$ is defined as

$$(3) \quad \mathcal{I}_g(b)(x, \xi) := \frac{\|\xi\|_x}{l} \int_0^{\frac{l}{\|\xi\|_x}} b \circ \varphi^\tau(x, \xi) d\tau, \quad (x, \xi) \in \mathring{T}^*M.$$

Above, φ^τ is the geodesic flow of (M, g) acting on the cotangent bundle, and l is the minimal common period of the trajectories of $\varphi^\tau|_{S^*M}$. With a slight abuse of notation, we shall still denote by $\mathcal{I}_g(V)$ the Radon transform of $V \circ \pi$, where $\pi : T^*M \rightarrow M$ stands for the canonical projection. Clearly, $\mathcal{I}_g(V) \in \mathcal{C}^\infty(\mathring{T}^*M)$ is zero-homogeneous with respect to the variable ξ . We shall consider the projection onto M of the set of critical points of the Radon transform of V :

$$\mathcal{C}(V) = \{x \in M : d_{(x, \xi)} \mathcal{I}_g(V) = 0 \text{ for some } \xi \in T_x^*M \setminus \{0\}\}.$$

The set $\mathcal{C}(V)$ is a union of geodesics of (M, g) and if the critical points of $\mathcal{I}_g(V)$ are non degenerate in an appropriate sense, $\mathcal{C}(V)$ consists in a finite number of geodesics (see Section 4). Note in particular that if $d_{(x, \xi)} \mathcal{I}_g(V) = 0$, then the geodesic issued from (x, ξ) is included in $\mathcal{C}(V)$. As an application of Theorem 1.9 below, one has

Theorem 1.1. *Suppose that (M, g) is a Compact Rank-One Symmetric Space and that $\mathcal{C}(V) \neq M$. Then there exist infinitely many geodesics γ of (M, g) such that $\nu(\gamma) = 0$ for every $\nu \in \mathcal{N}(\infty)$. In particular, $\delta_\gamma \notin \mathcal{N}(\infty)$, and*

$$\mathcal{N}(\infty) \neq \mathcal{N}_g.$$

When $d = 2$ and M is orientable, we are able to obtain a more precise result.

Theorem 1.2. *Suppose that $(M, g) = (\mathbb{S}^2, \text{can})$. Then every $\nu \in \mathcal{N}(\infty)$ can be written as:*

$$\nu = f \text{vol}_g + \nu_{\text{sing}},$$

where $f \in L^1(\mathbb{S}^2)$ and ν_{sing} is a nonnegative measure supported on $\mathcal{C}(V)$. When $\mathcal{C}(V)$ consists in a finite number of geodesics $\gamma_1, \dots, \gamma_n$ then one has:

$$\nu = f \text{vol}_g + \sum_{j=1}^n c_j \delta_{\gamma_j},$$

for some $c_j \geq 0$.

Remark 1.3. Note that the condition on $\mathcal{C}(V)$ in Theorems 1.1 and 1.2 is non-empty as soon as the Radon transforms $\mathcal{I}_g(V)$ is not constant, which is the case generically if, for instance, (M, g) is the 2-sphere endowed with its canonical metric – see e.g. appendix A of [22].

Our third result on eigenfunctions deals with Zoll surfaces that are not isometric to $(\mathbb{S}^2, \text{can})$. Recall from chapter 4 in [4] that the $C_{2\pi}$ -metrics of revolution on \mathbb{S}^2 are precisely those that can be written in spherical coordinates as:

$$g_\sigma = (1 + \sigma(\cos \theta))^2 d\theta^2 + \sin^2 \theta d\phi^2,$$

where σ is a smooth odd function on $[-1, 1]$ satisfying $\sigma(1) = 0$; such surfaces are also called Tannery surfaces. Combining our methods to some earlier results by Zelditch [53, 54], one can prove the following result for $C_{2\pi}$ -surfaces of revolution on \mathbb{S}^2 :

Theorem 1.4. *Let g_σ be a $C_{2\pi}$ -metric on \mathbb{S}^2 such that $\sigma'(0) \neq 0$ and suppose that $V \equiv 0$. Then there exist infinitely many geodesics γ of (\mathbb{S}^2, g_σ) such that $\nu(\gamma) = 0$ for every ν in $\mathcal{N}(\infty)$. In particular, $\delta_\gamma \notin \mathcal{N}(\infty)$, and*

$$\mathcal{N}(\infty) \neq \mathcal{N}_{g_\sigma}.$$

This result still holds if $V \neq 0$, provided the Radon transform of the potential does not differ from a constant from a certain function depending only on g_σ that is presented in the proof, which is given in paragraph 3.4. Theorems 1.1 and 1.2 will also be proved in Section 3; they are obtained as a consequence of our analysis of the time-dependent Schrödinger equation.

1.2. Long-time dynamics for the semiclassical Schrödinger equation. Our analysis of the time-dependent Schrödinger evolution fits in a natural way in the semiclassical framework. There will be three scales involved: the characteristic scale of oscillation of the solution \hbar , the size of the perturbation ϵ^2 and the time-scale τ . Our goal is to describe how these three scales affect the dynamics of the Schrödinger equation.

We shall assume that $\hbar \in (0, 1]$, and define the semiclassical the Schrödinger operator:

$$(4) \quad P_\epsilon(\hbar) := -\frac{\hbar^2 \Delta_g}{2} + \epsilon_\hbar^2 V,$$

where, as before, V belongs to $\mathcal{C}^\infty(M, \mathbb{R})$. We shall be interested in the regime in which the strength $\epsilon := (\epsilon_h)_{0 < h \leq 1}$ of the perturbation satisfies

$$\epsilon_h \rightarrow 0 \text{ as } \hbar \rightarrow 0^+.$$

The corresponding semiclassical Schrödinger equation is:

$$(5) \quad i\hbar \partial_t v_h = P_\epsilon(\hbar) v_h, \quad v_h|_{t=0} = u_h \in L^2(M).$$

We assume that the sequence of initial data $(u_h)_{h>0}$ is normalized in $L^2(M)$ and satisfies the following oscillation properties:

$$(6) \quad \limsup_{\hbar \rightarrow 0} \left\| \mathbf{1}_{[R, \infty)} (-\hbar^2 \Delta_g) u_h \right\|_{L^2(M)} \rightarrow 0, \quad \text{as } R \rightarrow \infty,$$

and

$$(7) \quad \limsup_{\hbar \rightarrow 0} \left\| \mathbf{1}_{[0, \delta]} (-\hbar^2 \Delta_g) u_h \right\|_{L^2(M)} \rightarrow 0, \quad \text{as } \delta \rightarrow 0^+.$$

Remark 1.5. Note that any normalized sequence $(u_n)_{n \in \mathbb{N}}$ satisfies (6) for some (\hbar_n) ; this is also the case with (7), provided (u_n) weakly converges to zero (the sequence (\hbar_n) here may be different from the one in (6)). Yet, in general, it is possible to construct normalized sequences (u_n) such that (6), (7) never hold simultaneously (see [18]). Note also that any sequence of normalized eigenfunctions of $-\frac{1}{2}\Delta_g + V$ satisfies (6), (7) with $\hbar_n = \lambda_n^{-1}$ where λ_n^2 is the corresponding eigenvalue.

We are interested in understanding the dynamics of the sequence of solutions (v_h) at time scales $\tau := (\tau_h)_{0 < h \leq 1}$ where

$$\lim_{\hbar \rightarrow 0^+} \tau_h = +\infty.$$

More precisely, the main object in our study is the sequence of time-scaled position densities:

$$\nu_h : (t, x) \mapsto |v_h(\tau_h t, x)|^2 = \left| e^{-i \frac{\tau_h}{\hbar} t P_\epsilon(\hbar)} u_h \right|^2(x).$$

The sequence (ν_h) is bounded in $\mathcal{D}'(\mathbb{R} \times M)$ and satisfies:

$$|\langle \nu_h, b \rangle_{\mathcal{D}' \times \mathcal{C}^\infty}| \leq \int_{\mathbb{R}} \|b(t, \cdot)\|_{L^\infty(M)} dt, \quad \text{for every } b \in \mathcal{C}_c^\infty(\mathbb{R} \times M).$$

Therefore, every accumulation point of the sequence (ν_h) is an element in $L^\infty(\mathbb{R}, \mathcal{P}(M))$, where $\mathcal{P}(M)$ denotes the set of probability measures on M . We shall denote by $\mathcal{N}(\tau, \epsilon)$ the set of such accumulation points obtained as the sequence of initial data (u_h) varies among normalized sequences satisfying (6) and (7).

Our goal is to understand the structure of the elements in $\mathcal{N}(\tau, \epsilon)$, in particular how this set depends on τ and ϵ as well as on the geometry of the manifold and the form of the potential. Again any element $\nu(t)$ in $\mathcal{N}(\tau, \epsilon)$ satisfies $\nu(t) \in \mathcal{N}_g$ for a.e. t in \mathbb{R} .

Remark 1.6. When $\epsilon_h = \hbar$ and $\tau_h = \hbar^{-1}$, one has

$$e^{-i \frac{\tau_h}{\hbar} t P_\epsilon(\hbar)} = e^{it(\frac{1}{2}\Delta_g - V)}.$$

Therefore this particular scaling corresponds to studying solutions to the non-semiclassical Schrödinger equation (1).

Remark 1.7. If u is an eigenfunction of the operator (2) then, by the previous remark,

$$|e^{-i\frac{\tau\hbar}{\hbar}tP_h(\hbar)}u|^2 = |e^{i\hbar\tau t(\frac{1}{2}\Delta_g - V)}u|^2 = |u|^2.$$

This implies that $\mathcal{N}(\infty) \subset \mathcal{N}(\tau, \hbar)$ for any time scale τ .

Remark 1.8. Most of our analysis can be extended to perturbations of the form $\epsilon_\hbar^2 Q_\hbar$, where Q_\hbar is a selfadjoint operator in $\Psi^{0,0}(M)$. Taking $Q_\hbar = V$ makes the exposition slightly simpler as we can use homogeneity properties of V .

The set $\mathcal{N}(\tau, \epsilon)$ was characterized in [38] in the case of short times ($\tau_\hbar \ll \hbar^{-2}$) and small perturbations ($\epsilon_\hbar \leq \hbar$) in the same geometric setting as in this article. It turns out that in that case, the elements in $\mathcal{N}(\tau, \epsilon)$ do not depend in t and one has

$$\mathcal{N}(\tau, \epsilon) = \mathcal{N}_g,$$

that is, $\mathcal{N}(\tau, \epsilon)$ gets in some sense as big as it can be.

In the context of negatively curved surfaces, some equidistribution properties of the elements of $\mathcal{N}(\tau, \epsilon)$ were obtained for strong enough perturbations [16, 44].

This problem has also been studied in great detail for the case $M = \mathbb{T}^d$ endowed with its natural metric. The articles [1, 2] describe this set in the case of small perturbations ($\epsilon_\hbar \leq \hbar$), whereas the case of stronger perturbations will be studied in more detail in [39]. In particular, in [1, 2] it is shown that when $(M, g) = (\mathbb{T}^d, \text{can})$ and $\epsilon_\hbar = \hbar$, the time scale $\tau_\hbar = \hbar^{-1}$ is critical for this problem. When $\tau_\hbar \ll \hbar^{-1}$ the set $\mathcal{N}(\tau, \epsilon)$ contains measures that are singular with respect to Lebesgue measure (in particular, it contains all δ_γ corresponding to closed geodesics); whereas if $\tau_\hbar \geq \hbar^{-1}$ then $\mathcal{N}(\tau, \epsilon)$ consists only of measures that are absolutely continuous with respect to the Lebesgue measure. Moreover, when $\tau_\hbar = \hbar^{-1}$ a precise description of the dependence on t of those measures is given.

In this article we show that a similar phenomenon takes place for Zoll manifolds. It turns out that the critical time scale for this problem is $\tau_\hbar = \epsilon_\hbar^{-2}$ in the case of strong perturbation ($\epsilon_\hbar \geq \hbar$) or $\tau_\hbar = \hbar^{-2}$ for small perturbations ($\epsilon_\hbar \ll \hbar$):

- Below the critical time-scale, one has $\mathcal{N}(\tau, \epsilon) = \mathcal{N}_g$, and we provide a formula to compute ν from the sequence of initial data.
- At the critical time-scale, measures in $\mathcal{N}(\tau, \epsilon)$ may depend in a non trivial way on t . We give an explicit propagation law that involves the Radon transform of the potential and the sequence of initial data.
- Above the critical time-scale, the restriction of the measures in $\mathcal{N}(\tau, \epsilon)$ to the complement of the critical set $\mathcal{C}(V)$ has some additional regularity. When $(M, g) = (\mathbb{S}^2, \text{can})$, this restriction is in fact absolutely continuous with respect to the Riemannian measure.

The precise statement of our results is given in Propositions 2.2, 2.3 and 2.4; they are formulated in terms of semiclassical measures and are presented in Section 2.2. In Section 2.3 we apply these results to the study of the quantum Loschmidt echo.

1.3. **Some results on the structure of $\mathcal{N}(\tau, \epsilon)$.** Let us now present some direct consequences of our results.

Theorem 1.9. *Suppose (M, g) is a P_l -manifold and that*

$$\tau_h \epsilon_h^2 \rightarrow +\infty.$$

Suppose one the following condition holds:

- (1) $\epsilon_h \hbar^{-1} \rightarrow +\infty$;
- (2) (M, g) is a compact rank one symmetric space.

Then, for every ν in $\mathcal{N}(\tau, \epsilon)$ and for every geodesic γ_0 that is not contained in $\mathcal{C}(V)$, one has:

$$\nu(t)(\gamma_0) = 0, \text{ for a.e. } t \in \mathbb{R}.$$

We note that for general P_l manifolds we are still able to obtain a similar result even if the condition $\epsilon_h \gg \hbar$ does not hold – see section 2 for more details. It is however more complicated to state as it involves the structure of a remainder term appearing in the natural decomposition of $\sqrt{-\Delta_g}$ on Zoll manifolds [9, 14]. In the case of a general P_l -metric on $M = \mathbb{S}^2$, an explicit expression of this term involving curvature terms and Jacobi fields was obtained by Zelditch in [53, 54] (see Paragraph 3.4 for more details on this issue).

This Theorem is a direct consequence of Propositions 2.2 and 2.3 and it will be proved in Paragraph 3.3. It tells us in particular that, for large enough times, solutions of the Schrödinger equation cannot be concentrated on closed geodesics corresponding to regular points of $\mathcal{I}_g(V)$. For instance, if one considers a sequence $(u_h)_{0 < h \leq 1}$ of coherent states that is microlocalized at a certain point $\rho \in \mathring{T}^*M$, then the corresponding solution will not be concentrated along the corresponding closed geodesic (for large enough times) provided that the geodesic consists of regular points.

Theorem 1.9 admits the following reformulation in terms of quasimodes for the Schrödinger operator $P_\epsilon(\hbar)$:

Corollary 1.10. *Suppose the hypotheses of Theorem 1.9 hold. Let $(\psi_h)_{0 < h \leq 1}$ be a normalized sequence in $L^2(M)$ satisfying*

$$\left(-\frac{\hbar^2 \Delta_g}{2} + \epsilon_h^2 V \right) \psi_h = E(\hbar) \psi_h + o(\hbar \epsilon_h^2),$$

with $E(\hbar) \rightarrow E \neq 0$ as $\hbar \rightarrow 0$. Then, for every weak- \star accumulation point ν_0 of the sequence

$$\nu_h := |\psi_h|^2 \text{vol}_g,$$

and for every geodesic γ_0 that is not contained in $\mathcal{C}(V)$, one has

$$\nu_0(\gamma_0) = 0.$$

When $V = 0$ and $d = 2$, Zelditch [55] has recently given criteria that ensure that every measure of the form δ_γ , γ being a geodesic of (M, g) , can be realized by a sequence of Gaussian beam-type quasimodes of order $\mathcal{O}(\hbar^2)$ (see Remark 3.6 in paragraph 3.4 for a more precise statement). We would like to stress the fact that we are not requiring

$\epsilon_{\hbar} \ll \sqrt{\hbar}$ for our results to hold. One often makes this assumption when studying the spectral properties of semiclassical operators with periodic bicharacteristics (see [41, 26] for instance) in order to keep the nice cluster structure of the spectrum of the operator $-\Delta_g$, see [9].

For the 2-sphere endowed with its canonical metric $g = \text{can}$, we are able to be more precise regarding the regularity of the elements in $\mathcal{N}(\tau, \epsilon)$:

Theorem 1.11. *Let V be a smooth function on $(\mathbb{S}^2, \text{can})$ such that $\mathcal{C}(V)$ consists in a finite number of geodesics $\gamma_1, \dots, \gamma_r$. Suppose*

$$\tau_{\hbar} \epsilon_{\hbar}^2 \rightarrow +\infty.$$

Then, any $\nu \in \mathcal{N}(\tau, \epsilon)$ is of the following form

$$\nu(t, \cdot) = f(t, \cdot) \text{vol}_{\text{can}} + \sum_{j=1}^r c_j(t) \delta_{\gamma_j},$$

where for a.e. t in \mathbb{R} , $f(t, \cdot) \in L^1(\mathbb{S}^2)$ and c_1, \dots, c_r are non-negative functions in $L^\infty(\mathbb{R})$.

Remark 1.12. This Theorem is a direct consequence of Proposition 2.3 and Corollary 4.4.

Remark 1.13. Recall that the space of geodesics on \mathbb{S}^2 can be identified with \mathbb{S}^2 [4] (remark 2.10) and that \mathcal{I}_{can} induces an isomorphism from $\mathcal{C}_{\text{even}}^\infty(\mathbb{S}^2)$ onto $\mathcal{C}_{\text{even}}^\infty(\mathbb{S}^2)$ [22]. Moreover, $\mathcal{I}_{\text{can}}(V) = 0$ for any odd function on \mathbb{S}^2 . In particular, for a generic choice of V , the assumption of the Theorem is satisfied.

1.4. An application to spectral theory. We mention the following Proposition on the spectral properties of $P_\epsilon(\hbar)$ which can also be obtained using the tools developed in the present article:

Proposition 1.14. *Suppose (M, g) is a P_l -manifold. Let $E_0 > 0$ and let $0 < 2\delta_0 \leq E_0$. Let $(E_j(\hbar))_{j=1}^{N(\hbar)}$ be the distinct eigenvalues of $P_\epsilon(\hbar)$ in the interval $[E_0 - \delta_0, E_0 + \delta_0]$.*

If $\mathcal{C}(V) \neq M$, then there exists some constant $c_0 > 0$ such that, for $\hbar > 0$ small enough, one has

$$s_{E_0, \delta_0}(\hbar) := \inf\{|E_j(\hbar) - E_k(\hbar)| : 1 \leq j \neq k \leq N(\hbar)\} \leq c_0 \hbar \epsilon_{\hbar}^2,$$

provided one of the following conditions holds:

- (1) $\epsilon_{\hbar} \hbar^{-1} \rightarrow +\infty$;
- (2) (M, g) is a compact rank-one symmetric space.

The proof of this result will be given in Section 6 and is based on Proposition 6.1 in that section, that shows that if the spectral separation condition in the conclusion of Proposition 1.14 is violated, then necessarily all uniform measures supported on geodesics can be realized as accumulation points of moduli squares of eigenfunctions. In the case where $d = 2$ and $\hbar^N \ll \epsilon_{\hbar} \ll \sqrt{\hbar}$ (where N is some positive exponent related to the clustering of the unperturbed operator), a much stronger result was for instance obtained in [26]. In fact, it was proved in this reference that, near regular values of $\mathcal{I}_g(V)$, one can obtain an asymptotic expansion of the eigenvalues with a level spacing which is exactly

of order $\hbar\epsilon_h^2$. Compared with this result, the above Proposition only provides a simple criterium for the existence of distinct eigenvalues which are asymptotically at distance less than $\hbar\epsilon_h^2$. On the other hand, it is valid in any dimension and even for strong perturbation, meaning $\epsilon_h \geq \sqrt{\hbar}$.

Acknowledgments. Much of this work was done while the first author was visiting Laboratoire Paul Painlevé at Université de Lille 1 in October 2013 in the framework of the CEMPI program (ANR-11-LABX-0007-01). He wishes to thank this institution for its warm hospitality and support. The authors thank San Vũ Ngọc for useful discussions on integrable systems.

2. MAIN RESULTS: PROPAGATION OF SEMICLASSICAL MEASURES

We now describe our main results, that are formulated using the notion of semiclassical or Wigner measure.

2.1. Semiclassical measures. In order to study the asymptotic properties of the solutions of (5), we will make use of the so-called semiclassical measures [17], or more precisely of their time dependent version [38] – see also appendix B for a brief reminder. We now recall their construction. For a given t in \mathbb{R} , we denote the Wigner distribution at time $t\tau_h$ by

$$(8) \quad w_h(t\tau_h)(a) := \langle v_h(t\tau_h), \text{Op}_h(a)v_h(t\tau_h) \rangle,$$

where $\text{Op}_h(a)$ is a \hbar -pseudodifferential operator with principal symbol $a \in \mathcal{C}_c^\infty(T^*M)$ – see appendix A. This quantity represents the distribution of the solution of (5) in the phase space T^*M .

Recall now that we can extract a subsequence $\hbar_n \rightarrow 0^+$ as $n \rightarrow +\infty$ such that, for every a in $\mathcal{C}_c^\infty(T^*M)$ and for every θ in $L^1(\mathbb{R})$,

$$\lim_{\hbar_n \rightarrow 0^+} \int_{\mathbb{R} \times T^*M} \theta(t)a(x, \xi)w_{\hbar_n}(t\tau_{\hbar_n}, dx, d\xi)dt = \int_{\mathbb{R} \times T^*M} \theta(t)a(x, \xi)\mu(t, dx, d\xi)dt,$$

where $(t, x, \xi) \mapsto \mu(t, x, \xi)$ belongs to $L^\infty(\mathbb{R}, \mathcal{M}(T^*M))$, with $\mathcal{M}(T^*M)$ the set of finite complex measures carried on T^*M . Recall also that, for a.e. $t \in \mathbb{R}$, $\mu(t, \cdot)$ is in fact a *probability measure* which is carried on \mathring{T}^*M and which is *invariant by the geodesic flow* φ^s on T^*M . For instance, $\mu(t, \cdot)$ can be the normalized Lebesgue measure along closed orbit of the geodesic flow. We refer to appendix B for a brief reminder of these results from [38].

We will denote by $\mathcal{M}(\tau, \epsilon)$ the set of accumulation points of the sequences

$$\mu_h : (t, x, \xi) \longmapsto w_h(t\tau_h, x, \xi)$$

as (u_h) varies among normalized sequences satisfying (6) and (7).

Remark 2.1. Thanks to the frequency assumption (6), one can also verify that $\mathcal{N}(\tau, \epsilon)$ corresponds in fact to the projections on M of the elements of $\mathcal{M}(\tau, \epsilon)$.

2.2. Propagation at different time scales. The Zoll structure on M allows to prove that every element in $\mathcal{M}(\tau, \epsilon)$ satisfies an additional invariance or transport property, depending on the relative size of τ and ϵ .

We denote by φ_V^t the Hamiltonian flow associated to the Radon transform of the potential $\mathcal{I}_g(V)$. Note that φ_V^t commutes with the geodesic flow φ^s .

Let us start presenting our results in the particular case when (M, g) is a Compact Rank-One Symmetric Space, since they are somewhat simpler to describe. It turns out, as it was already stated in the introduction, that the time scale $\tau_h = \epsilon_h^{-2}$ is critical for this problem. More precisely, the following result holds:

Proposition 2.2. *Suppose (M, g) is a Compact Rank-One Symmetric Space. Let $\mu \in \mathcal{M}(\tau, \epsilon)$ and denote by μ_0 the semiclassical measure of the sequence of initial data used to generate μ . The following results hold.*

i) *If $\tau_h \epsilon_h^2 \rightarrow 0^+$ then μ is continuous with respect to t and, for every $b \in \mathcal{C}_c^\infty(T^*M)$ and every $t \in \mathbb{R}$:*

$$\mu(t)(b) = \mu_0(\mathcal{I}_g(b)).$$

ii) *If $\tau_h \epsilon_h^2 = 1$ then μ is continuous with respect to t and, for every $b \in \mathcal{C}_c^\infty(T^*M)$ and every $t \in \mathbb{R}$:*

$$\mu(t)(b) = \mu_0(\mathcal{I}_g(b) \circ \varphi_V^t).$$

iii) *If $\tau_h \epsilon_h^2 \rightarrow +\infty$ then μ has an additional invariance property. For almost every $t \in \mathbb{R}$ and every $s \in \mathbb{R}$:*

$$(\varphi_V^s)_* \mu(t) = \mu(t).$$

This result can be obtained by similar arguments as the proof of Propositions 2.3, 2.4 presented below, which describe additional properties satisfied by the elements of $\mathcal{M}(\tau, \epsilon)$ in the more general case of P_l -manifolds. In this case, one must take into account a certain (φ^s) -invariant function q_0 on \mathring{T}^*M that depends on the metric g . Let $\varphi_{q_0}^t$ and $\varphi_{V+q_0}^t$ denote respectively the Hamiltonian flows on \mathring{T}^*M generated by q_0 and $\mathcal{I}_g(V) + q_0$.

The first of these result is concerned with the “big perturbation” regime, when the square root of the size of the perturbation dominates the characteristic length scale of oscillations.

Proposition 2.3. *Suppose M is endowed with a P_l -metric g and that:*

$$\epsilon_h \geq \hbar.$$

*There exists a smooth, 0-homogeneous and (φ^s) -invariant function q_0 defined on \mathring{T}^*M (and depending only on (M, g)) such that, for every $\mu \in \mathcal{M}(\tau, \epsilon)$ associated to a sequence of initial data with an unique semiclassical measure μ_0 , the following holds:*

i) *If $\tau_h \epsilon_h^2 \rightarrow 0^+$, then, μ is continuous in the t variable and, for every $b \in \mathcal{C}_c^\infty(T^*M)$ and every $t \in \mathbb{R}$:*

$$(9) \quad \mu(t)(b) = \mu_0(\mathcal{I}_g(b)).$$

ii) If $\tau_{\hbar}\epsilon_{\hbar}^2 = 1$ then μ is continuous in the t variable and, for every $b \in \mathcal{C}_c^\infty(T^*M)$ and every $t \in \mathbb{R}$:

$$(10) \quad \mu(t)(b) = \mu_0(\mathcal{I}_g(b) \circ \varphi_{V+q_0}^t), \text{ if } \epsilon_{\hbar} = \hbar,$$

$$(11) \quad \mu(t)(b) = \mu_0(\mathcal{I}_g(b) \circ \varphi_V^t), \text{ if } \epsilon_{\hbar}\hbar^{-1} \rightarrow +\infty.$$

iii) If $\tau_{\hbar}\epsilon_{\hbar}^2 \rightarrow +\infty$, then μ has an additional invariance property. For almost every $t \in \mathbb{R}$ and every $s \in \mathbb{R}$:

$$(12) \quad (\varphi_{V+q_0}^s)_*\mu(t) = \mu(t), \text{ if } \epsilon_{\hbar} = \hbar,$$

$$(13) \quad (\varphi_V^s)_*\mu(t) = \mu(t), \text{ if } \epsilon_{\hbar}\hbar^{-1} \rightarrow +\infty.$$

The next result deals with the ‘‘small perturbation’’ regime, when \hbar dominates ϵ_{\hbar} (and therefore, the time scale must be compared to \hbar^2 instead of ϵ_{\hbar}^2). As it can be expected, the effect of the perturbation in this case is negligible. We however present the precise statement for the sake of completeness.

Proposition 2.4. *Suppose M is endowed with a P_1 -metric g and let q_0 be the function introduced in Proposition 2.3. Suppose that:*

$$\epsilon_{\hbar}\hbar^{-1} \rightarrow 0^+.$$

Then, for every $\mu \in \mathcal{M}(\tau, \epsilon)$ associated to a sequence of initial data with an unique semi-classical measure μ_0 , the following holds:

i) *If $\tau_{\hbar}\hbar^2 \rightarrow 0^+$, then, μ is continuous in the t variable and, for every $b \in \mathcal{C}_c^\infty(T^*M)$ and every $t \in \mathbb{R}$:*

$$(14) \quad \mu(t)(b) = \mu_0(\mathcal{I}_g(b)).$$

ii) *If $\tau_{\hbar}\hbar^2 = 1$ then μ is continuous in the t variable and, for every $b \in \mathcal{C}_c^\infty(T^*M)$ and every $t \in \mathbb{R}$:*

$$(15) \quad \mu(t)(b) = \mu_0(\mathcal{I}_g(b) \circ \varphi_{q_0}^t).$$

iii) *If $\tau_{\hbar}\hbar^2 \rightarrow +\infty$, then μ has an additional invariance property. For almost every $t \in \mathbb{R}$ and every $s \in \mathbb{R}$:*

$$(16) \quad (\varphi_{q_0}^s)_*\mu(t) = \mu(t).$$

We note that the statements of these Propositions remain valid for $V \equiv 0$ (in fact, the results of Proposition 2.3 are contained in those of Proposition 2.4 in this case).

As it will become clear from our proof, the symbol q_0 is related to the ‘‘natural’’ decomposition of the operator $\sqrt{-\Delta_g}$ on Zoll manifolds – see for instance Theorem 1.1 of [9]. As stated in the Proposition, this symbol depends only on the choice of the metric. We emphasize yet that it is given by an hardly explicit formula – see Remark 3.5 or [53, 54].

However, when (M, g) is a Compact Rank-One Symmetric Space, one can take $q_0 = 0$ and therefore it is possible to derive the result in Proposition 2.2 without making any assumption regarding the relative size of \hbar and ϵ_{\hbar} .

Remark 2.5. When $\mathcal{I}_g(V)$ is constant, the Hamiltonian flow φ_V^t acts trivially, and some of the above statements are thus empty. This is the case, for instance, when $(M, g) = (\mathbb{S}^d, \text{can})$ and $V = V(x)$ does not depend on ξ and is an odd function plus a constant (see [29], Theorems 1.17, 1.23).

2.3. Application to the study of the quantum Loschmidt echo. As another application of the methods developed to prove Proposition 2.2, 2.3 and 2.4, we derive some properties on the so-called quantum Loschmidt echo. This quantity is defined as follows

$$\mathcal{E}_h(t\tau_h) := |\langle v_h(t\tau_h), v_h^0(t\tau_h) \rangle|^2,$$

where

- $v_h(t\tau_h)$ is the solution of (5);
- $v_h^0(t\tau_h)$ is the solution of (5) when we pick $V \equiv 0$.

This quantity was introduced by Peres in [43] and it allows to measure the sensitivity of a quantum system to perturbations of the Hamiltonian. Peres predicted that this quantity should typically goes to 0 and that the decay rate indicates the chaotic or integrable nature of the underlying classical system. Since this seminal work of Peres [43], many progresses have been made in the physics literature regarding the asymptotic properties of this quantity, especially in the context of chaotic systems. We refer the reader to [19, 34, 20] for recent surveys on these questions. Our approach allows to study a slightly related quantity:

$$F_h(t\tau_h) := \langle v_h(t\tau_h), v_h^0(t\tau_h) \rangle.$$

Up to an extraction $\hbar_n \rightarrow 0$, one can suppose that there exists $F(t)$ in $\mathcal{D}'(\mathbb{R})$ such that, for every θ in $\mathcal{C}_c^\infty(\mathbb{R})$,

$$\lim_{\hbar_n \rightarrow 0} \int_{\mathbb{R}} \theta(t) F_{\hbar_n}(t\tau_{\hbar_n}) dt = \int_{\mathbb{R}} \theta(t) F(t) dt.$$

Our last result gives a description of the limit distribution $F(t)$ in the context of P_l -manifolds:

Proposition 2.6. *Suppose (M, g) is a P_l -manifold and that*

$$\lim_{\hbar \rightarrow 0^+} \epsilon_h \hbar^{-\frac{1}{2}} = 0.$$

Suppose also that one the following condition holds:

- (1) $\epsilon_h \hbar^{-\frac{3}{2}} \rightarrow +\infty$,
- (2) (M, g) is a compact rank one symmetric space.

Then, for every $t \mapsto F(t)$ associated to a sequence of initial data with an unique semiclassical measure μ_0 , the following holds

i) If $\lim_{\hbar \rightarrow 0^+} \frac{\tau_h \epsilon_h^2}{\hbar} = 0$, then, for every t in \mathbb{R} :

$$F(t) = 1.$$

ii) If $\tau_h = \frac{\hbar}{\epsilon_h^2}$, then, $F(t)$ is continuous in the t variable and, for every t in \mathbb{R} :

$$F(t) = \mu_0 \left(e^{it\mathcal{I}_g(V)} \right).$$

iii) If $\lim_{\hbar \rightarrow 0^+} \frac{\tau_{\hbar} \epsilon_{\hbar}^2}{\hbar} = +\infty$ and if $\{\mathcal{I}_g(V) = 0\} = \emptyset$, then, for every t in \mathbb{R} ,

$$F(t) = 0.$$

This Theorem will follow from Proposition 5.2 and its proof will make use of similar tools as the ones used to prove Proposition 2.2, 2.3 and 2.4. Note that other mathematical studies of the long time properties of the quantum Loschmidt echo appeared recently in various geometric settings: 1-dimensional systems [6, 10], negatively curved surfaces [44].

3. AVERAGING, TRANSPORT AND INVARIANCE

In this section, we will prove Propositions 2.2, 2.3 and 2.4. It is organized as follows. First, we recall an averaging procedure due to Weinstein [51] which we formulate in a semiclassical language following [8, 28, 12]. Then, in paragraph 3.2, we deduce the above Propositions from this averaging Lemma, and, in paragraph 3.3, we derive the proof of a slightly stronger version of Theorem 1.9 from these Propositions.

3.1. Semiclassical averaging Lemma. The following result is a quantum analogue of the averaging method for classical dynamical systems.

Theorem 3.1. *Let (M, g) be a P_l -manifold. Then, for every $b \in C_c^\infty(\mathring{T}^*M)$, there exists an operator $\langle B_{\hbar} \rangle \in \Psi^{0,0}(M)$ whose principal symbol is the classical average:*

$$(17) \quad \mathcal{I}_g(b)(x, \xi) := \frac{\|\xi\|_x}{l} \int_0^{\frac{l}{\|\xi\|_x}} b \circ \varphi^s(x, \xi) ds,$$

and which satisfies:

$$(18) \quad [\langle B_{\hbar} \rangle, P_0(\hbar)] = \mathcal{O}_{L^2 \rightarrow L^2}(\hbar^3).$$

In addition, if (M, g) is a Compact Rank-One Symmetric Space, $\langle B_{\hbar} \rangle$ can be chosen such that the above formula is exact, that is:

$$(19) \quad [\langle B_{\hbar} \rangle, P_0(\hbar)] = 0.$$

This type of result is rather well-known and goes back to [51, 14, 9]. We will give here a semiclassical version of the argument presented in those references following the presentation of [12] (chapter 15) – see also [8, 28] for a semiclassical treatment.

The proof of Theorem 3.1 will be done in three steps.

3.1.1. Reparametrization of the classical Hamiltonian. We have made the assumption that all the geodesics of M have a common least period $l > 0$. This means that, for every (x, ξ) in \mathring{T}^*M , one has $\varphi^{\frac{l}{\sqrt{2E}}}(x, \xi) = (x, \xi)$, where $E = \frac{\|\xi\|_x^2}{2}$. Following [12], the first step will be to “reparametrize” the Hamiltonian, both at the quantum and at the classical level, in order to have a common period 2π for the flow on \mathring{T}^*M . For that purpose, we set

$$\tilde{P}_0(\hbar) := \frac{l}{2\pi} \sqrt{-\hbar^2 \Delta_g}.$$

The Hamiltonian corresponding to this operator is given by $\tilde{p}_0(x, \xi) := \|\xi\|_x$, and the Hamiltonian vector fields of p_0 and \tilde{p}_0 are related in the following way:

$$X_{\tilde{p}_0}(x, \xi) = \frac{l}{2\pi\sqrt{2E}} X_{p_0}(x, \xi),$$

where $E = \frac{\|\xi\|_x^2}{2}$. If we denote by ψ^t the Hamiltonian flow of \tilde{p}_0 , we find that, for every (x, ξ) in \mathring{T}^*M and for every s in \mathbb{R} , one has

$$\psi^s(x, \xi) = \varphi^{\frac{l}{2\pi\|\xi\|_x}s}(x, \xi), \text{ and } \psi^{2\pi}(x, \xi) = (x, \xi).$$

3.1.2. Periodicity of the quantum propagator. We recall that the Fourier integral operator associated to the Hamiltonian vector field satisfies a certain periodicity property. This follows from the periodicity of the classical flow. We will denote by $\alpha \in \mathbb{Z}$ the common Maslov index of the closed trajectories of ψ^t on the energy layers $\tilde{p}_0^{-1}((0, +\infty))$. According to Lemma 29.2.1 in [33] – see also [9] or [28], one knows that there exists a polyhomogeneous pseudodifferential operator A of order 1 and a polyhomogeneous pseudodifferential operator Q of order -1 (see Ch. 18 in [32] for the precise definitions) such that the following holds:

- (1) $\frac{l}{2\pi}\sqrt{-\Delta_g} = A + \frac{\alpha}{4} + Q$;
- (2) $[\sqrt{-\Delta_g}, Q] = 0$;
- (3) $\text{Sp}(A) \subset \mathbb{N}$.

Translated in our semiclassical framework, we get

$$\tilde{P}_0(\hbar) = \hbar A + \frac{\alpha\hbar}{4} + \hbar Q,$$

and, in particular,

$$e^{-\frac{2i\pi}{\hbar}(\tilde{P}_0(\hbar) - \frac{\hbar\alpha}{4} - \hbar Q)} = \text{Id}_{L^2(M)}.$$

Remark 3.2. In the case of a compact rank one symmetric space, the situation is slightly simpler using the explicit description of the spectrum – see paragraph 8.8 in [4]. In fact, we can write that

$$-\tilde{\Delta}_g := -\Delta_g + \left(\frac{\pi\alpha}{2l}\right)^2 \text{Id} = \left(\frac{2\pi}{l}\right)^2 \left(\tilde{A} + \frac{\alpha}{4}\right)^2,$$

where \tilde{A} is a polyhomogeneous pseudodifferential operator of order one such that $\text{Sp}(\tilde{A}) \subset \mathbb{N}$, and then

$$(20) \quad e^{-\frac{i2\pi}{\hbar}\left(\frac{l}{2\pi}\sqrt{-\hbar^2\tilde{\Delta}_g - \frac{\hbar\alpha}{4}}\right)} = e^{-i2\pi\tilde{A}} = \text{Id}_{L^2(M)}.$$

3.1.3. Averaging procedure. We will now verify that the operator $\tilde{P}_0(\hbar)$ satisfies the commutation property (18) where the operator $\langle B_\hbar \rangle$ is defined as the quantum average of the pseudodifferential operator $\text{Op}_\hbar(b)$ by the Fourier integral operator associated to $\hbar A$. This kind of averaging procedure is standard in this context and it seems that it first appeared in Weinstein's article [51].

Let $\delta < \frac{1}{2} < R$. For every symbol $b \in C_c^\infty(T^*M)$ which is supported in $p_0^{-1}([\delta, R])$, we define the following averaged operator:

$$\langle B_\hbar \rangle := \frac{1}{2\pi} \int_0^{2\pi} e^{itA} \text{Op}_\hbar(b) e^{-itA} dt.$$

According to Egorov's Theorem, $\langle B_\hbar \rangle$ is an element in $\Psi^{0,0}(M)$ whose principal symbol is equal to $\mathcal{I}_g(b)$, which is a smooth function since b is supported in $p_0^{-1}([\delta, R])$ and therefore:

$$\mathcal{I}_g(b)(x, \xi) = \frac{1}{2\pi} \int_0^{2\pi} b \circ \psi^s(x, \xi) ds.$$

Using the fact that $\text{Sp}(A) \subset \mathbb{N}$, we start noticing that the following commutation relation holds:

$$(21) \quad [\langle B_\hbar \rangle, A] = 0.$$

Let $0 \leq \chi \leq 1$ be a smooth cut-off function in $C_c^\infty\left(\left[\frac{l}{2\pi}\sqrt{\delta}, \frac{l}{2\pi}\sqrt{4R}\right]\right)$ satisfying $\chi \equiv 1$ in a small neighborhood of $\left[\frac{l}{2\pi}\sqrt{2\delta}, \frac{l}{2\pi}\sqrt{2R}\right]$. We can now use property (21) in order to derive an expression for the commutator $[\langle B_\hbar \rangle, P_0(\hbar)]$. First note that:

$$[\langle B_\hbar \rangle, P_0(\hbar)] = [\langle B_\hbar \rangle, \chi(\tilde{P}_0(\hbar))P_0(\hbar)] + \mathcal{O}(\hbar^\infty).$$

Recall now that $P_0(\hbar) = \frac{(2\pi)^2}{l^2} \tilde{P}_0(\hbar)^2$. We use the definition of A in order to write:

$$\tilde{P}_0^2(\hbar) = \hbar^2 A^2 + \frac{\hbar^2 \alpha^2}{16} + \frac{\hbar^2 \alpha}{2} A + \hbar^2 Q_0,$$

where

$$(22) \quad Q_0 := (AQ + QA) + \frac{\alpha}{2} Q + Q^2,$$

which also defines an operator commuting with $\tilde{P}_0(\hbar)$. Therefore, taking into account identity (21), we finally obtain:

$$(23) \quad [\langle B_\hbar \rangle, P_0(\hbar)] = \frac{2\pi^2 \hbar^2}{l^2} [\langle B_\hbar \rangle, \chi(\tilde{P}_0(\hbar))Q_0] + \mathcal{O}(\hbar^\infty).$$

Using pseudodifferential calculus rules, one knows that $\frac{2\pi^2 \hbar^2}{l^2} \chi(\tilde{P}_0(\hbar))Q_0$ is an element in $\Psi^{-\infty,0}(M)$ with principal symbol equal to $\chi(\|\xi\|l/(2\pi))q_0(x, \xi)$ where q_0 is the principal symbol of the polyhomogeneous pseudodifferential operator $(AQ + QA)(2\pi)^2/l^2$ of degree 0. Recall that it defines a smooth function on \hat{T}^*M which is 0-homogeneous. Note that this function depends only on (M, g) . Thus, using the composition formula for pseudodifferential operators and the Calderón-Vaillancourt Theorem [56], we conclude that:

$$[\langle B_\hbar \rangle, P_0(\hbar)] = \mathcal{O}_{L^2 \rightarrow L^2}(\hbar^3),$$

as we wanted to prove.

Remark 3.3. In the case that (M, g) is a Compact Rank-One Symmetric Space, we can use Remark 3.2 and mimick the above proof in order to obtain an exact commutation formula:

$$[\langle B_{\hbar} \rangle, P_0(\hbar)] = 0,$$

where

$$\langle B_{\hbar} \rangle := \frac{1}{2\pi} \int_0^{2\pi} e^{\frac{it}{\hbar} \left(\frac{t}{2\pi} \sqrt{-\hbar^2 \tilde{\Delta}_g - \frac{\hbar\alpha}{4}} \right)} \text{Op}_{\hbar}(b) e^{-\frac{it}{\hbar} \left(\frac{t}{2\pi} \sqrt{-\hbar^2 \tilde{\Delta}_g - \frac{\hbar\alpha}{4}} \right)} dt.$$

3.2. Invariance and transport: proof of Propositions 2.2, 2.3, and 2.4. We will now use Theorem 3.1 to prove additional invariance properties satisfied by every element μ in $\mathcal{M}(\tau, \epsilon)$. Recall that, for almost every t in \mathbb{R} , $\mu(t, \cdot)$ is a probability measure which is invariant by the geodesic flow φ^s . We also emphasize that, for almost every t in \mathbb{R} , one has $\mu(t)(\mathring{T}^*M) = 1$ thanks to (6) and (7). These conditions imply that for every $b \in \mathcal{C}_c^\infty(\mathring{T}^*M)$ and almost every t one has:

$$(24) \quad \mu(t)(b) = \mu(t)(\mathcal{I}_g(b)).$$

Let φ_V^t , $\varphi_{q_0}^t$ and $\varphi_{V+q_0}^t$ denote respectively the Hamiltonian flows on \mathring{T}^*M generated by $\mathcal{I}_g(V)$, q_0 and $\mathcal{I}_g(V) + q_0$. In the case of a Compact Rank One Symmetric Space, we use the convention $q_0 \equiv 0$. Note that φ_V^t , $\varphi_{q_0}^t$ and $\varphi_{V+q_0}^t$ commute with the geodesic flow, since $\{p_0, \mathcal{I}_g(V)\} = \{p_0, q_0\} = 0$.

We will now give the proof of Propositions 2.2, 2.3, and 2.4. Let $\mu \in \mathcal{M}(\tau, \epsilon)$. This means that there exists a sequence $(u_{\hbar})_{0 < \hbar \leq 1}$ of normalized states in $L^2(M)$ satisfying the frequency assumptions (6) and (7) such that the Wigner distributions $\mu_{\hbar} : t \mapsto w_{\hbar}(t\tau_{\hbar})$ corresponding to the solutions $e^{-\frac{it\tau_{\hbar}P_{\epsilon}(\hbar)}{\hbar}} u_{\hbar}$ converge to μ – see appendix B.

Assume moreover that $(u_{\hbar})_{0 < \hbar \leq 1}$ has a semiclassical measure μ_0 , meaning that the Wigner distributions $w_{\hbar}(0)$ weak- \star converge towards μ_0 as $\hbar \rightarrow 0^+$. Let b be a smooth compactly supported function on \mathring{T}^*M .

Remark 3.4. We note that, as $\mu(t)(M \times \{0\}) = 0$ a.e., we can restrict ourselves to proving the invariance properties for such functions b .

Let $\langle B_{\hbar} \rangle$ be the operator obtained from b using Theorem 3.1. One clearly has:

$$(25) \quad \frac{d}{dt} \left\langle u_{\hbar}, e^{\frac{it\tau_{\hbar}P_{\epsilon}(\hbar)}{\hbar}} \langle B_{\hbar} \rangle e^{-\frac{it\tau_{\hbar}P_{\epsilon}(\hbar)}{\hbar}} u_{\hbar} \right\rangle = \frac{i\tau_{\hbar}}{\hbar} \left\langle u_{\hbar}, e^{\frac{it\tau_{\hbar}P_{\epsilon}(\hbar)}{\hbar}} [P_{\epsilon}(\hbar), \langle B_{\hbar} \rangle] e^{-\frac{it\tau_{\hbar}P_{\epsilon}(\hbar)}{\hbar}} u_{\hbar} \right\rangle.$$

Recall that, by Theorem 3.1 and by the composition rules for pseudodifferential operators, one has:

$$\frac{\tau_{\hbar}}{\hbar} [P_{\epsilon}(\hbar), \langle B_{\hbar} \rangle] = \mathcal{O}_{L^2 \rightarrow L^2}(\tau_{\hbar} \hbar^2 + \tau_{\hbar} \epsilon_{\hbar}^2).$$

In the case of a CROSS, we also note that the remainder is in fact of order $\mathcal{O}_{L^2 \rightarrow L^2}(\tau_{\hbar} \epsilon_{\hbar}^2)$. If $\tau_{\hbar} \epsilon_{\hbar}^2 \rightarrow 0^+$ (and $\tau_{\hbar} \hbar^2 \rightarrow 0^+$ when (M, g) is not a CROSS), after integrating both sides of identity (25) on the interval $t \in [0, \tau]$ and taking limits as $\hbar \rightarrow 0^+$, one has for almost every $\tau \in \mathbb{R}$:

$$\mu(\tau)(b) = \mu(\tau)(\mathcal{I}_g(b)) = \mu_0(\mathcal{I}_g(b)),$$

which, in view of (24) concludes the proof of i) of the three Propositions.

We now turn to the proof of ii) of these different Propositions. Equation (23) implies that the commutator takes the form:

$$\frac{\tau_h}{\hbar}[P_\epsilon(\hbar), \langle B_\hbar \rangle] = \tau_h \hbar [\text{Op}_\hbar(q_0), \langle B_\hbar \rangle] + \mathcal{O}_{L^2 \rightarrow L^2}(\tau_h \hbar^3) + \frac{\tau_h \epsilon_h^2}{\hbar}[V, \langle B_\hbar \rangle],$$

where q_0 was defined in paragraph 3.1.3 – see also remark 3.5 below. Integrating again (25) with respect to t , letting $\hbar \rightarrow 0^+$ and using the composition formula for pseudodifferential operators gives that, for every $\tau \in \mathbb{R}$ the following holds

$$(26) \quad \mu(\tau)(\mathcal{I}_g(b)) - \mu_0(\mathcal{I}_g(b)) = \int_0^\tau \mu(t)(\{L, \mathcal{I}_g(b)\}) dt,$$

where

$$L := \begin{cases} q_0 & \text{if } \epsilon_h \hbar^{-1} \rightarrow 0^+, \\ q_0 + V & \text{if } \epsilon_h = \hbar, \\ V & \text{if } \epsilon_h \hbar^{-1} \rightarrow +\infty, \end{cases}$$

In the case of a CROSS, we let $L := V$ without any assumptions on the size of the perturbations. The geodesic flow preserves the symplectic form on T^*M , therefore:

$$(27) \quad \{\mathcal{I}_g(L), \mathcal{I}_g(b)\} = \mathcal{I}_g(\{\mathcal{I}_g(L), b\}) = \mathcal{I}_g(\{L, \mathcal{I}_g(b)\}).$$

In particular, combining this equality to (24), we can rewrite (26) as

$$\forall \tau \in \mathbb{R}, \quad \mu(\tau)(\mathcal{I}_g(b)) - \mu_0(\mathcal{I}_g(b)) = \int_0^\tau \mu(t)(\{\mathcal{I}_g(L), \mathcal{I}_g(b)\}) dt,$$

Therefore, using again (27), one has $\frac{d}{d\tau} \mu(\tau)(\mathcal{I}_g(b \circ \varphi_L^{-\tau})) = 0$, and thus

$$\mu(\tau)(\mathcal{I}_g(b)) = \mu_0(\mathcal{I}_g(b \circ \varphi_L^\tau)) = \mu_0(\mathcal{I}_g(b) \circ \varphi_L^\tau),$$

where φ_L^τ is the Hamiltonian flow of $\mathcal{I}_g(L)$. This completes the proof of ii) of Propositions 2.2, 2.3, and 2.4.

We consider now the large time regime, meaning $\tau_h \epsilon_h^2 \rightarrow +\infty$ (in the case of Propositions 2.2 and 2.3) or $\tau_h \hbar^2 \rightarrow +\infty$ (in the case of Proposition 2.4). Let θ be an element in $\mathcal{C}_c^1(\mathbb{R})$. We use an integration by parts on (25) to derive:

$$\begin{aligned} \int_{\mathbb{R}} \theta'(t) \left\langle u_\hbar, e^{\frac{it\tau_h P_\epsilon(\hbar)}{\hbar}} \langle B_\hbar \rangle e^{-\frac{it\tau_h P_\epsilon(\hbar)}{\hbar}} u_\hbar \right\rangle dt = \\ - \frac{i\tau_h}{\hbar} \int_{\mathbb{R}} \theta(t) \left\langle u_\hbar, e^{\frac{it\tau_h P_\epsilon(\hbar)}{\hbar}} [P_\epsilon(\hbar), \langle B_\hbar \rangle] e^{-\frac{it\tau_h P_\epsilon(\hbar)}{\hbar}} u_\hbar \right\rangle dt. \end{aligned}$$

Thanks to relation (23), we deduce that

$$\frac{i}{\hbar} \int_{\mathbb{R}} \theta(t) \left\langle u_\hbar, e^{\frac{it\tau_h P_\epsilon(\hbar)}{\hbar}} [\hbar^2 \text{Op}_\hbar(q_0) + \epsilon_h^2 V, \langle B_\hbar \rangle] e^{-\frac{it\tau_h P_\epsilon(\hbar)}{\hbar}} u_\hbar \right\rangle dt = \mathcal{O}(\tau_h^{-1}) + \mathcal{O}(\hbar^3).$$

Again, in the case of a CROSS, the remainder is only $\mathcal{O}(\tau_h^{-1})$. Using the composition formula for pseudodifferential operators, we find that

$$(28) \quad \int_{\mathbb{R}} \theta(t) \langle w_\hbar(t\tau_h), \{\hbar^2 q_0 + \epsilon_h^2 V, \mathcal{I}_g(b)\} \rangle dt = \mathcal{O}(\tau_h^{-1}) + \mathcal{O}(\hbar^3).$$

Taking limits as $\hbar \rightarrow 0^+$ and using identities (24), (27) shows that

$$\int_{\mathbb{R}} \theta(t) \mu(t) (\{\mathcal{I}_g(L), \mathcal{I}_g(b)\}) dt = 0.$$

This concludes the proof of part iii) in each case.

Remark 3.5. Again, we would like to stress the fact that the symbol q_0 is only related to the choice of the metric on M . In order to get an (implicit) expression for q_0 , we have to write that

$$0 = [\langle B_{\hbar} \rangle, \hbar^2 A^2] = [\langle B_{\hbar} \rangle, \tilde{P}_0(\hbar)^2 - \hbar^2 \text{Op}_{\hbar}(q_0)] + \mathcal{O}_{L^2 \rightarrow L^2}(\hbar^4).$$

Using energy cut-offs, this can be in fact rewritten as

$$[\langle B_{\hbar} \rangle, \text{Op}_{\hbar}(q_0)] = \frac{l^2}{2\pi^2 \hbar^2} [\langle B_{\hbar} \rangle, P_0(\hbar)] + \mathcal{O}_{L^2 \rightarrow L^2}(\hbar^2).$$

From this expression, we can get an expression for $\{\mathcal{I}_g(b), q_0\}$ by identifying the principal symbol of the right hand side which turns out to be a delicate task [53, 54]. We will discuss this in more details in paragraph 3.4 in the case of metrics of revolution on \mathbb{S}^2 .

3.3. Proof of Theorem 1.9. Theorem 1.9 is now an easy consequence of Proposition 2.3. We will in fact prove something slightly more precise that works for any size of ϵ_{\hbar} . We use the above convention for defining the function L . Recall that L is a smooth function on \dot{T}^*M which is 0-homogeneous and that, in the case of a compact rank-one symmetric space, one has always $L = V \circ \pi$.

Let $\mu \in \mathcal{M}(\tau, \epsilon)$. Suppose $\tau_{\hbar} \epsilon_{\hbar}^2 \rightarrow +\infty$, or $\tau_{\hbar} \hbar^2 \rightarrow +\infty$ when $\epsilon_{\hbar} \ll \hbar$ (if (M, g) is not a CROSS). Then, Propositions 2.2, 2.3 and 2.4 assert that $\mu(t, \cdot)$ is invariant with respect to the flow φ_L^{τ} (the Hamiltonian flow of $\mathcal{I}_g(L)$) for a.e. $t \in \mathbb{R}$.

First, we consider $\Gamma_0 \subset S^*M$ a closed orbit of φ^s that is not included in

$$\text{Crit}(L) := \left\{ \rho \in \dot{T}^*M : d_{\rho} \mathcal{I}_g(L) = 0 \right\}.$$

The fact that $\Gamma_0 \not\subset \text{Crit}(L)$ is equivalent to the fact that the set Γ_0 is not invariant by φ_L^{τ} – see Lemma 4.1 for a proof of this fact. For a given closed orbit $\Gamma_0 \subset S^*M$ and for a fixed $\lambda > 0$, we now define

$$\Gamma_0(\lambda) := \{(x, \lambda \xi) : (x, \xi) \in \Gamma_0\}.$$

We fix a compact interval $[c_1, c_2] \subset (0, +\infty)$ and $\lambda \in [c_1, c_2]$. Since φ_L^{τ} commutes with the geodesic flow, it follows that all of the sets $\varphi_L^{\tau}(\Gamma_0(\lambda))$ are distinct orbits of φ^{τ} (at least for τ small enough). On the other hand, the invariance of $\mu(t, \cdot)$ implies that, for a.e. t in \mathbb{R} ,

$$\mu(t) \left(\bigcup_{\lambda \in [c_1, c_2]} \varphi_L^{\tau}(\Gamma_0(\lambda)) \right) = \mu(t) \left(\bigcup_{\lambda \in [c_1, c_2]} \Gamma_0(\lambda) \right)$$

for every τ in some interval with non-empty interior. Since $\mu(t, \cdot)$ is finite, one must have

$$(29) \quad \mu(t) \left(\bigcup_{\lambda \in [c_1, c_2]} \Gamma_0(\lambda) \right) = 0,$$

for a.e. t in \mathbb{R} .

Consider now the projection $\nu(t, \cdot)$ of $\mu(t, \cdot)$ on M . Suppose $\Gamma_0 \subset S^*M$ is a closed orbit of φ^s and denote by $\gamma_0 := \pi(\Gamma_0)$ where $\pi : T^*M \rightarrow M$ is the canonical projection. Suppose that Γ_0 is not contained in $\text{Crit}(L)$. Using (29), we note that

$$\nu(t)(\gamma_0) = \mu(t) \left(\bigcup_{\lambda \in (0, \infty)} \{(x, \xi) \in \pi^{-1}(\gamma_0) : \|\xi\| = \lambda \text{ and } (x, \xi) \notin \Gamma_0(\lambda)\} \right).$$

Using invariance by the geodesic flow and the fact that the measure is finite, a similar argument as above allows to conclude that this quantity is in fact equal to 0.

3.4. Explicit expression of q_0 on \mathbb{S}^2 . In this last section, we briefly recall a result due to Zelditch which gives a more or less explicit expression for q_0 – see Theorem 3 in [54] for more details. We suppose that g is a $C_{2\pi}$ -metric on \mathbb{S}^2 ; in this case, every geodesic is a simple curve, see [21]. We fix Γ_0 a closed geodesic on S^*M issued from the point (x_0, ξ_0) , i.e.

$$\Gamma_0 := \{\varphi^s(x_0, \xi_0) = (x(s), \xi(s)) : s \in \mathbb{R}\}.$$

We denote by $\tilde{f}_{x_0, \xi_0}(t)$ the unique solution of the ordinary differential equation

$$f''(t) + K(x(t))f(t) = 0, \quad f(0) = 1, \quad f'(0) = 0,$$

and by $f_{x_0, \xi_0}(t)$ the unique solution of the ordinary differential equation

$$f''(t) + K(x(t))f(t) = 0, \quad f(0) = 0, \quad f'(0) = 1,$$

where K is the scalar curvature. Using the conventions of Appendix C, we also define

$$\mathcal{K}(x, \xi) := g_x^*(d_x K, \xi^\perp).$$

The formula obtained by Zelditch can then be expressed as follows:

$$q_0(x_0, \xi_0) := \frac{1}{8\pi} \int_0^{2\pi} K(x(t)) dt + \frac{1}{24\pi} \int_0^{2\pi} \mathcal{K}(x(t), \xi(t)) \tilde{f}_{x_0, \xi_0}(t)^2 R_{x_0, \xi_0}(t) dt.$$

where

$$\begin{aligned} R_{x_0, \xi_0}(t) &:= \tilde{f}_{x_0, \xi_0}(t) \int_0^t \mathcal{K}(x(s), \xi(s)) f_{x_0, \xi_0}(s)^3 ds \\ &\quad - 3f_{x_0, \xi_0}(t) \int_0^t \tilde{f}_{x_0, \xi_0}(s) \mathcal{K}(x(s), \xi(s)) f_{x_0, \xi_0}(s)^2 ds. \end{aligned}$$

We proceed as in [54] and specialize this expression to the case of $C_{2\pi}$ -metrics of revolution on \mathbb{S}^2 . According to [4] – Theorem 4.13, such a metric can be written in spherical coordinates as:

$$g = (1 + \sigma(\cos \theta))^2 d\theta^2 + \sin^2 \theta d\phi^2,$$

where σ is a smooth odd function on $[-1, 1]$ satisfying $\sigma(1) = 0$. Recall from Paragraph 4.17 in [4] that the sectional curvature can be expressed as follows:

$$K(\theta) = \frac{1}{(1 + \sigma(\cos \theta))^3} (1 + \sigma(\cos \theta) - \cos \theta \sigma'(\cos \theta)).$$

Let $\pi(\Gamma_0)$ be the closed geodesic corresponding to $\theta = \frac{\pi}{2}$. We note that the curvature is constant and equal to 1 on this closed geodesic. For every (x_0, ξ_0) in Γ_0 , we find then

$$q_0(x_0, \xi_0) := \frac{1}{4} + \frac{3^2 \sigma'(0)^3}{4\pi} \int_0^\pi \left(\frac{\cos^6 \phi}{3} - \cos^4 \phi - \cos^2 \phi \sin^4 \phi \right) d\phi = \frac{1}{4} - \frac{3\sigma'(0)^3}{4},$$

In particular, q_0 is not equal to $\frac{1}{4}$ on this closed geodesic if $\sigma'(0) \neq 0$. This condition is of course generic among the possible choice of σ . Using the symmetry of revolution and the Gauss-Bonnet formula, one verifies that, for any (x_0, ξ_0) belonging to a geodesic orthogonal to the geodesic $\theta = \frac{\pi}{2}$, one has $q_0(x_0, \xi_0) = \frac{1}{4}$. In particular, q_0 is not a constant function on $T^*\mathbb{S}^2$. Combining this observation to the results of paragraph 3.3, we deduce Theorem 1.4 – see Remark 1.7.

Remark 3.6. It is interesting to compare the results in this section with the results recently obtained by Zelditch in [55]. In that reference, a necessary and sufficient condition was given in order that, for any geodesic γ , there exists a sequence of Gaussian-beams (ψ_h) satisfying

$$(-\hbar^2 \Delta_g - \hbar^2 Q_0) \psi_h = \psi_h + \mathcal{O}(\hbar^4),$$

and such that $|\psi_h|^2 \text{vol}_g$ tends to δ_γ . Here, Proposition 2.3 implies that the following holds whenever $\pi^{-1}(\gamma) \not\subset \text{Crit}(q_0)$: for any sequence of quasimodes satisfying

$$-\hbar^2 \Delta_g \psi_h = \psi_h + o(\hbar^3),$$

and such that $|\psi_h|^2 \text{vol}_g$ converges to ν , one must have $\nu(\gamma) = 0$.

4. MEASURES WITH TWO INVARIANCE PROPERTIES

This section is of more geometric flavor; it is devoted to analyzing the structure of measures on the cotangent bundle that are invariant by two flows that commute.

4.1. Measures that are invariant by two Hamiltonians that commute. As before, we write $p_0(x, \xi) := \frac{1}{2} \|\xi\|_x^2$ and φ^s denotes the geodesic flow acting on T^*M .

Let L be a smooth function on \mathring{T}^*M satisfying:

- L is 0-homogeneous in the ξ variable, i.e. $L(x, \xi) = L(x, \lambda \xi)$ for every $\lambda > 0$.
- The one-form dL does not vanish identically.
- L Poisson-commutes with p_0 : $\{L, p_0\} = 0$.

We shall denote by X_0 and X_L the respective Hamiltonian vector fields of p_0 and L ; let φ_L^τ be the flow of X_L . Then the flows φ_L^τ , and φ^s commute.

The purpose of this section is to describe the set of probability measures on \mathring{T}^*M that are simultaneously invariant by the the flow φ_L^τ and by the geodesic flow φ^s . We are especially interested in the case when $d = 2$ and M is orientable,

We introduce the set $\text{Crit}(L)$ of critical points of L in \mathring{T}^*M :

$$\text{Crit}(L) := \left\{ \rho \in \mathring{T}^*M : d_\rho L = 0 \right\}.$$

This is a closed set in \mathring{T}^*M that is invariant by both flows $\varphi^s, \varphi_L^\tau$. More precisely, the following holds.

Lemma 4.1. *Under the above assumptions, $\text{Crit}(L)$ is formed by those orbits of the geodesic flow that are invariant by φ_L^τ .*

Proof. Since φ^t and φ_L^s commute, $\text{Crit}(L)$ is formed by orbits of the geodesic flow that are fixed by φ_L^s . On the other hand, if the orbit of the geodesic flow issued from some ρ_0 is invariant by φ_L^s then necessarily $X_L(\rho_0) = \alpha X_0(\rho_0)$ for some $\alpha \in \mathbb{R}$. This forces $X_L(\rho) = \alpha X_0(\rho)$ for every ρ in the orbit. Since L is zero-homogeneous with respect to ξ it turns out that X_L and X_0 are orthogonal with respect to the Sasaki metric (see Appendix C). Therefore, $\alpha = 0$, as we wanted to prove. \square

Consider now the set:

$$\mathcal{R}(L) := \mathring{T}^*M \setminus \text{Crit}(L).$$

This is an open subset of \mathring{T}^*M that is invariant by φ^s and φ_L^τ . The map

$$\phi : \mathbb{R}^2 \times \mathcal{R}(L) \ni (s, t, \rho) \longmapsto \varphi^s \circ \varphi_L^t(\rho) \in \mathcal{R}(L),$$

is a group action of \mathbb{R}^2 on $\mathcal{R}(L)$. Moreover, for any $\rho_0 \in \mathcal{R}(L)$ the map:

$$\phi_{\rho_0} : \mathbb{R}^2 \ni (s, t) \longmapsto \varphi^s \circ \varphi_L^t(\rho_0) \in \mathcal{R}(L),$$

is an immersion. Therefore, the stabilizer group G_{ρ_0} of ρ_0 under ϕ is discrete. This proves that the orbits of the action ϕ are either diffeomorphic to the torus \mathbb{T}^2 , to the cylinder $\mathbb{T} \times \mathbb{R}$ or to \mathbb{R}^2 .

The moment map:

$$\Phi : \mathcal{R}(L) \ni (x, \xi) \longmapsto (p_0(x, \xi), L(x, \xi)) \in \mathbb{R}^2,$$

is a submersion, and for every $(E, J) \in \Phi(\mathcal{R}(L))$ the level set

$$\Lambda_{(E, J)} := \Phi^{-1}(E, J),$$

is a smooth submanifold of $\mathcal{R}(L)$ of codimension two. Note that the 0-homogeneity of L implies that $\Lambda_{(\alpha E, J)} = h_\alpha(\Lambda_{(E, J)})$ for every $\alpha > 0$, and $h_\alpha : \mathring{T}^*M \rightarrow \mathring{T}^*M$ being the homothety of ratio α on the fibers.

When $d = 2$ the couple p_0, L forms a completely integrable system on $\mathcal{R}(L)$; and the map ϕ_{ρ_0} induces a diffeomorphism:

$$\phi_{\rho_0} : \mathbb{R}^2 / G_{\rho_0} \longrightarrow \Lambda_{(E_0, J_0)}^{\rho_0}, \quad \text{for } (E_0, J_0) := \Phi(\rho_0);$$

Above, $\Lambda_{(E_0, J_0)}^{\rho_0}$ denotes the connected component of $\Lambda_{(E_0, J_0)}$ that contains ρ_0 . Therefore, if $\Lambda_{(E_0, J_0)}^{\rho_0}$ is compact then it is an embedded Lagrangian torus in T^*M . In that case, we shall write $\mathbb{T}_{\rho_0}^2 := \mathbb{R}^2 / G_{\rho_0}$. See [13, 40] for more detailed proofs.

Our next result clarifies the structure of the set of probability measures on $\mathcal{R}(L)$ that are invariant by the geodesic flow and the flow φ_L^τ . Let us introduce first some notation: $\mathcal{R}_c(L)$ will be the set formed by those $\rho \in \mathcal{R}(L)$ such that $\Lambda_{\Phi(\rho)}^\rho$ is compact.

Proposition 4.2. *Suppose that $d = 2$. Let μ be a probability measure on $\mathcal{R}(L)$ that is invariant by φ^s and φ_L^τ and set $\bar{\mu} := \Phi_*\mu$. Then, for every $b \in \mathcal{C}_c(\mathcal{R}(L))$, one has*

$$\int_{\mathcal{R}(L)} b(x, \xi) \mu(dx, d\xi) = \int_{\Phi(\mathcal{R}(L))} \int_{\Lambda_{(E,J)}} b(x, \xi) \lambda_{E,J}(dx, d\xi) \bar{\mu}(dE, dJ),$$

where, for $(E, J) \in \Phi(\mathcal{R}(L))$, the measure $\lambda_{E,J}$ is a convex combination of the (normalized) Haar measures on the tori $\Lambda_{(E,J)}^\rho$ for $\rho \in \Lambda_{(E,J)} \cap \mathcal{R}_c(L)$ (see equation (30)).

Proof. The disintegration Theorem (see e.g. Th. 5.14 in [15]) gives, for $\bar{\mu}$ -a.e. $(E, J) \in \mathcal{R}(L)$, the existence of a family of probability measures $\mu_{E,J}$ concentrated on $\Lambda_{(E,J)}$ such that:

$$\int_{\mathcal{R}(L)} b(x, \xi) \mu(dx, d\xi) = \int_{\Phi(\mathcal{R}(L))} \int_{\Lambda_{(E,J)}} b(x, \xi) \mu_{E,J}(dx, d\xi) \bar{\mu}(dE, dJ),$$

for every $b \in \mathcal{C}_c(\mathcal{R}(L))$. The measures $\mu_{E,J}$ inherit the same invariance properties as μ . In particular,

$$\int_{\Lambda_{(E,J)}} b(x, \xi) \mu_{E,J}(dx, d\xi) = \int_{\Lambda_{(E,J)}} b(\phi(s, t, x, \xi)) \mu_{E,J}(dx, d\xi)$$

for every $(s, t) \in \mathbb{R}^2$. Each connected component of the manifold $\Lambda_{(E,J)}$ has a group structure induced by the map ϕ_ρ , for any $\rho \in \Lambda_{(E,J)}$. The invariance property of $\mu_{E,J}|_{\Lambda_{(E,J)}^\rho}$ is equivalent to stating that it is invariant by translations in the group. Therefore, $\mu_{E,J}|_{\Lambda_{(E,J)}^\rho}$ must coincide with a multiple of the Haar measure on $\Lambda_{(E,J)}^\rho$ for every $\rho \in \Lambda_{(E,J)}$. If $\Lambda_{(E,J)}^\rho$ is non compact, this measure is infinite and $\mu_{E,J}(\Lambda_{(H,J)}^\rho) = 0$. \square

An explicit formula for the restriction of the measure $\lambda_{E,J}$ to a connected component $\Lambda_{(E,J)}^\rho$ with $\rho \in \mathcal{R}_c(L) \cap \Lambda_{(E,J)}$ is the following:

$$(30) \quad \int_{\Lambda_{(E,J)}^\rho} b(x, \xi) \lambda_{E,J}(dx, d\xi) = c \int_{\mathbb{T}_{\rho_0}^2} b(\phi_\rho(s, t)) ds dt,$$

for some constant $c \in [0, 1]$. Proposition 4.2 merely states that bi-invariant measures are superpositions of measures of this form.

Note that so far we have not used the fact that the geodesic flow is periodic. When this is the case it turns out that for every $\rho_0 \in \mathcal{R}_c(L)$ the stabilizer group G_{ρ_0} of ρ_0 contains an element of the form $(q, 0)$ for some $q \in \mathbb{R} \setminus \{0\}$.

4.2. Structure of the projection onto the base. Our next goal is to study the regularity properties of the projection of a bi-invariant measure μ onto the base manifold M . We are going to prove that they decompose as a measure that is absolutely continuous with respect to the Riemannian volume plus a singular measure supported on the projection of the critical set $\text{Crit}(L)$.

Theorem 4.3. *Suppose that M is an orientable P_l -surface. Let μ be a probability measure on $\mathcal{R}(L)$ that is invariant by φ^s and φ_L^τ . Then $\nu := \pi_*\mu$ is a probability measure on M that is absolutely continuous with respect to the Riemannian measure.*

Denote by $\mathcal{N}(L)$ the convex closure of the set of measures $\delta_{\pi\circ\gamma}$ where $\gamma \subset \mathring{T}^*M$ ranges over the orbits of the geodesic flow that are contained in $\text{Crit}(L)$. The following is a direct consequence of Theorem 4.3.

Corollary 4.4. *Suppose that M is an orientable P_l -surface. The projection $\nu := \pi_*\mu$ of a probability measure μ on \mathring{T}^*M that is invariant by φ^s and φ_L^τ can be decomposed as:*

$$\nu = f \text{vol}_g + \alpha \nu_{\text{sing}}$$

where $f \in L^1(M)$, $\alpha \in [0, 1]$ and $\nu_{\text{sing}} \in \mathcal{N}(L)$.

Remark 4.5. For a “generic” function L , the set $\text{Crit}(L)$ is formed by critical points that are non degenerate with respect to directions that are orthogonal to the geodesic flow and to the vertical vector field that generates the dilations on the fibers. In this case, for every $E > 0$ the set $\text{Crit}(L) \cap p_0^{-1}(E)$ consists in a finite number of orbits of the geodesic flow that project onto a finite family of geodesics $\{\gamma_1, \dots, \gamma_r\}$ in M . The measures ν described in Corollary 4.4 then take the simple form:

$$\nu = f \text{vol}_g + \sum_{j=1}^r c_j \delta_{\gamma_j},$$

where $c_j \geq 0$ for $j = 1, \dots, r$ and $\sum_{j=1}^r c_j \leq 1$.

Theorem 4.3 essentially follows from a structure result for the projections of the Lagrangian tori $\Lambda_{(E,J)}^{\rho_0}$ onto M : the projection $\pi|_{\Lambda_{(E,J)}^{\rho_0}}$ is a local diffeomorphism onto its image except at points lying on a finite family of smooth curves (called sometimes a *caustic*). See also Lemma 2.1 in [5].

Lemma 4.6. *Let $\rho_0 \in \mathcal{R}_c(L)$; then there exist a finite family $\{\Gamma_{\rho_0}^j\}_{j=1}^n$ of smooth disjoint closed curves contained in $\Lambda_{(E,J)}^{\rho_0}$ that are transversal to the geodesic flow and such that $\Gamma_{\rho_0}^{\text{caus}} := \bigsqcup_{j=1}^n \Gamma_{\rho_0}^j$ is exactly the set of singular points of $\pi|_{\Lambda_{(E,J)}^{\rho_0}} : \Lambda_{(E,J)}^{\rho_0} \rightarrow M$.*

Proof. Let $\{X_0, W, Y_0, U\}$ be the orthogonal frame on \mathring{T}^*M defined in Appendix C. Since L is 0-homogeneous with respect to ξ , the Hamiltonian vector field X_L has no component in X_0 , that is:

$$g_\rho^S(X_L(\rho), X_0(\rho)) = g_{\pi(\rho)}(d_\rho \pi(X_L(\rho)), d_\rho \pi(X_0(\rho))) = 0.$$

The tangent space of $\Lambda_{(E,J)}^{\rho_0}$ at a point ρ is spanned by $\{X_0(\rho), X_L(\rho)\}$. The points $\rho \in \Lambda_{(E,J)}^{\rho_0}$ close to which $\pi|_{\Lambda_{(E,J)}^{\rho_0}}$ is not a local diffeomorphism are precisely those at which $d_\rho \pi(X_L(\rho)) = 0$. Define:

$$F(t, s) := g_{\pi(\phi_{\rho_0}(t,s))}(d_{\phi_{\rho_0}(t,s)} \pi(X_L(\phi_{\rho_0}(t, s))), d_{\phi_{\rho_0}(t,s)} \pi(W(\phi_{\rho_0}(t, s)))).$$

One has:

$$F(t, s) = 0 \iff X_L(\phi_{\rho_0}(t, s)) \in \text{Ker}(d_{\phi_{\rho_0}(t, s)}\pi).$$

We are going to apply the implicit function Theorem to the equation $F(t, s) = 0$ in order to prove our claim. Note that $J_s(t) := d_{\phi_{\rho_0}(t, s)}\pi(X_L(\phi_{\rho_0}(t, s))) = \partial_s(\pi \circ \phi_{\rho_0})(t, s)$ is a Jacobi field along the geodesic $\gamma_s(t) := \pi(\varphi^t(\varphi_L^s(\rho_0)))$. Write $Z_s(t) := d_{\phi_{\rho_0}(t, s)}\pi(W(\phi_{\rho_0}(t, s)))$. One has:

$$\partial_t F(t, s) = g_{\pi(\phi_{\rho_0}(t, s))} \left(\frac{DJ_s}{dt}(t), Z_s(t) \right) + g_{\pi(\phi_{\rho_0}(t, s))} \left(J_s(t), \frac{DZ_s(t)}{dt} \right),$$

where $\frac{D}{dt}$ is the covariant derivative along the curve $\gamma_s(t)$. If $F(t_0, s_0) = 0$ then $J_{s_0}(t_0) = 0$ but:

$$g_{\pi(\phi_{\rho_0}(t_0, s_0))} \left(\frac{DJ_{s_0}}{dt}(t_0), Z_{s_0}(t_0) \right) \neq 0;$$

the reason for this is that if that term were to vanish, this would imply that $\frac{D}{dt}J_{s_0}(t_0)$ is proportional to $\gamma'(t_0)$. Therefore, the Jacobi field $J_{s_0}(t)$ should be proportional to $t\gamma'(t)$ which forces J_{s_0} to vanish identically. In other words, $X_L(\phi_{\rho_0}(t, s_0)) \in \text{Ker}(d_{\phi_{\rho_0}(t, s_0)}\pi)$ for every t . But this is a contradiction, since the values of t for which that property holds turn out to form a discrete set in \mathbb{R} , as we show next.

This fact is a consequence of the twist property of the vertical subbundle as stated in [42]: recall that $\Lambda_{(E, J)}^{\rho_0}$ is a Lagrangian submanifold of T^*M ; applying Proposition 2.11 of [42] to the Lagrangian subspaces $T_\rho\Lambda_{(E, J)}^{\rho_0}$ with $\rho \in \Lambda_{(E, J)}^{\rho_0}$, we deduce that, for every s_0 the set of the $t \in \mathbb{R}$ such that:

$$\text{Ker}(d_{\phi_{\rho_0}(t, s_0)}\pi) \cap T_{\phi_{\rho_0}(t, s_0)}\Lambda_{(L, J)}^{\rho_0} = \text{Ker}(d_{\phi_{\rho_0}(t, s_0)}\pi) \cap d_{\varphi_L^{s_0}(\rho_0)}\varphi_0^t(T_{\varphi_L^{s_0}(\rho_0)}\Lambda_{(E, J)}^{\rho_0}) \neq \{0\},$$

is discrete.

If the geodesic $\varphi^t(\rho_0)$ is q -periodic, this set is in fact finite modulo $q\mathbb{Z}$. Set $s_0 = 0$ and let $\{t_1, \dots, t_r\} \subset \mathbb{R}/q\mathbb{Z}$ be the set at which $F(t, 0) = 0$. We can apply the implicit function Theorem to conclude the existence of a unique family of smooth functions $t_j(s)$ with $s \in \mathbb{R}/r(J)\mathbb{Z}$ satisfying $t_j(0) = t_j$ and $F(t_j(s), s) = 0$. Taking into account that $\partial_t F(t, s) \neq 0$ as soon as $F(t, s) = 0$, we conclude that two curves $(t_j(s), s)$ and $(t_i(s), s)$ are either disjoint or they coincide. Let us relabel these curves in order to have $(t_j(s), s)$ with $j = 1, \dots, n$ are disjoint. Since these curves are the unique solutions to the equation $F(t, s) = 0$, the trajectories $(t_j(s), s)$ must be closed. Therefore, the smooth curves $\Gamma_{\rho_0}^j := \{\phi_{\rho_0}(t_j(s), s) : s \in \mathbb{R}/r(J)\mathbb{Z}\}$ with $j = 1, \dots, n$ are disjoint and closed. \square

To conclude the proof of Theorem 4.3 let $B \subset M$ be of zero Riemannian measure. By construction,

$$\nu(B) = \int_{\Phi(\mathcal{R})(L)} \lambda_{E, J}(\pi^{-1}(B) \cap \Lambda_{(E, J)}) \bar{\mu}(dE, dJ).$$

Fix a connected component $\Lambda_{(E, J)}^{\rho_0}$ in $\Lambda_{(E, J)}$, let Ω_{ρ_0} be any connected component of $\Lambda_{(E, J)}^{\rho_0} \setminus \Gamma_{\rho_0}^{\text{caus}}$. Then $\lambda_{E, J}(\Gamma_{\rho_0}^{\text{caus}}) = 0$ by Lemma 4.6 and $\pi|_{\Omega_{\rho_0}}$ is a local diffeomorphism; therefore, $\pi^{-1}(B) \cap \Omega_{\rho_0}$ has $\lambda_{E, J}$ -measure equal to zero. Hence,

$$\lambda_{E, J}(\pi^{-1}(B) \cap \Lambda_{(H, J)}^{\rho_0}) = 0.$$

This shows that $\nu(E) = 0$ and therefore, ν is absolutely continuous with respect to the Riemannian measure.

5. QUANTUM LOSCHMIDT ECHO ON ZOLL MANIFOLDS

In this section, we revisit the proof of section 3 in order to study a quantity which is related to the so called quantum Loschmidt echo defined in the section 2. Precisely, we prove Proposition 2.6. For every b in $\mathcal{C}_c^\infty(T^*M)$ and every t in \mathbb{R} , we define

$$M_{\hbar}(t\tau_{\hbar})(b) := \langle v_{\hbar}(t\tau_{\hbar}), \text{Op}_{\hbar}(b)v_{\hbar}^0(t\tau_{\hbar}) \rangle,$$

where

- $(\tau_{\hbar})_{0 < \hbar \leq 1}$ is a scale of times satisfying $\lim_{\hbar \rightarrow 0} \tau_{\hbar} = +\infty$;
- $v_{\hbar}(t\tau_{\hbar})$ is the solution of (5);
- $v_{\hbar}^0(t\tau_{\hbar})$ is the solution of (5) when we pick $V \equiv 0$.

If we suppose that the sequence of initial data $(u_{\hbar})_{0 < \hbar \leq 1}$ is normalized and that it satisfies (6) and (7), then, proceeding as in appendix (B), we can extract a subsequence $\hbar_n \rightarrow 0$ such that, for every $\theta \in \mathcal{C}_c^\infty(\mathbb{R})$ and for every a in $\mathcal{C}_c^\infty(T^*M)$, one has

$$\lim_{n \rightarrow +\infty} \int_{\mathbb{R}} \theta(t) M_{\hbar_n}(t\tau_{\hbar_n})(b) dt = \int_{\mathbb{R}} \theta(t) \left(\int_{T^*M} b(x, \xi) M(t, dx, d\xi) \right) dt,$$

where $t \mapsto M(t)$ belongs to $L^\infty(\mathbb{R}, \mathcal{M}(T^*M))$. Thanks to the frequency assumptions, the support of $M(t)$ is included in $T^*M \setminus (M \times \{0\})$. Moreover, in the case where $\epsilon_{\hbar} \ll \sqrt{\hbar}$, we also observe that, for a.e. t in \mathbb{R} and for every s in \mathbb{R} , one has

$$(31) \quad \int_{T^*M} b(x, \xi) M(t, dx, d\xi) = \int_{T^*M} b \circ \varphi^s(x, \xi) M(t, dx, d\xi).$$

Remark 5.1. In order to make the link with Proposition 2.6, we note that, thanks to (6) and (7), one has

$$\lim_{n \rightarrow +\infty} \int_{\mathbb{R}} \theta(t) M_{\hbar_n}(t\tau_{\hbar_n})(1) dt = \int_{\mathbb{R}} \theta(t) \left(\int_{T^*M} M(t, dx, d\xi) \right) dt.$$

Proposition 2.6 follows then from the previous remark and the following Proposition.

Proposition 5.2. *Suppose (M, g) is a P_l -manifold and that*

$$\lim_{\hbar \rightarrow 0^+} \epsilon_{\hbar} \hbar^{-\frac{1}{2}} = 0.$$

Suppose also that one the following condition holds:

- (1) $\epsilon_{\hbar} \hbar^{-\frac{3}{2}} \rightarrow +\infty$,
- (2) (M, g) is a Compact Rank One Symmetric Space.

Then, for every $t \mapsto F(t)$ associated to a sequence of initial data with an unique semiclassical measure μ_0 , the following holds:

*i) If $\lim_{\hbar \rightarrow 0^+} \frac{\tau_{\hbar} \epsilon_{\hbar}^2}{\hbar} = 0$, then, for every t in \mathbb{R} and for every b in $\mathcal{C}_c^\infty(T^*M)$,*

$$M(t)(b) = \mu_0(\mathcal{I}_g(b)).$$

ii) If $\tau_h = \frac{h}{\epsilon_h^2}$, then, for every t in \mathbb{R} and for every b in $\mathcal{C}_c^\infty(T^*M)$,

$$M(t)(b) = \mu_0(\mathcal{I}_g(b)e^{it\mathcal{I}_g(V)}).$$

iii) If $\lim_{h \rightarrow 0^+} \frac{\tau_h \epsilon_h^2}{h} = +\infty$, then, for a.e. every t in \mathbb{R} and for every b in $\mathcal{C}_c^\infty(T^*M)$,

$$M(t)(b\mathcal{I}_g(V)) = 0.$$

Remark 5.3. In the case (iii), we emphasize that we can deduce that

$$\text{supp}M(t) \subset \{\mathcal{I}_g(V) = 0\}.$$

We will now give the proof of this Proposition which follows the same lines as the proofs in section 3.

Proof. Let b be a smooth compactly supported function on T^*M . Using (31), one has that

$$M(t)(b) = M(t)(\mathcal{I}_g(b)).$$

We start our proof by computing the following derivative

$$\frac{d}{dt} (\langle v_h(t\tau_h), \langle B_h \rangle v_h^0(t\tau_h) \rangle),$$

where $\langle B_h \rangle$ is the operator from Theorem 3.1. Using Theorem 3.1, we obtain in fact

$$(32) \quad \frac{d}{dt} (\langle v_h(t\tau_h), \langle B_h \rangle v_h^0(t\tau_h) \rangle) = -\frac{i\epsilon_h^2 \tau_h}{h} \langle v_h(t\tau_h), V \langle B_h \rangle v_h^0(t\tau_h) \rangle + \mathcal{O}(\tau_h h^2).$$

We observe that, if (M, g) is a compact rank-one symmetric space, then the remainder is in fact equal to 0. The case (i) follows immediatly by integrating (32) between 0 and t . In the case where $\tau_h = \frac{h}{\epsilon_h^2}$, one finds that, for every θ in $\mathcal{C}_c^\infty(\mathbb{R})$, one has

$$\int_{\mathbb{R}} \theta'(t) M(t)(\mathcal{I}_g(b)) dt = i \int_{\mathbb{R}} \theta(t) M(t)(\mathcal{I}_g(b)V) dt = i \int_{\mathbb{R}} \theta(t) M(t)(\mathcal{I}_g(b)\mathcal{I}_g(V)) dt,$$

where the first equality follows from (32) and the second one from (31). The case (ii) follows then from the above relation. Finally, when $\lim_{h \rightarrow 0^+} \frac{\tau_h \epsilon_h^2}{h} = +\infty$, we deduce from (32) that

$$\int_{\mathbb{R}} \theta(t) M(t)(\mathcal{I}_g(b)\mathcal{I}_g(V)) dt = 0,$$

from which (iii) follows according to (31). □

6. RELATION TO SPECTRAL PROBLEMS: PROOF OF PROPOSITION 1.14

In this last section, we explain how time dependent semiclassical measures can be used to prove some results on the spectrum of the semiclassical Schrödinger operator

$$P_\epsilon(\hbar) := -\frac{\hbar^2 \Delta_g}{2} + \epsilon_\hbar^2 V.$$

Precisely, we prove Proposition 1.14 from the introduction which establishes the existence of distinct eigenvalues which are asymptotically close to each other. In order to do that, we revisit an argument that was given in [37] – proof of Theorem 2 from that reference – to show the following result.

Proposition 6.1. *Let (M, g) be a P_l -manifold. Assume that there exist $E_0 > 0$ and $0 < 2\delta_0 \leq E_0$ such that the function $s_{E_0, \delta_0}(\hbar)$ defined in the statement of Proposition 1.14 satisfies:*

$$\frac{s_{E_0, \delta_0}(\hbar)}{\hbar \epsilon_\hbar^2} \longrightarrow \infty, \quad \text{as } \hbar \longrightarrow 0^+.$$

Then one has:

$$\{\delta_\gamma : \gamma \text{ geodesic of } (M, g)\} \subset \mathcal{N}(\infty).$$

Proposition 1.14 is then a consequence of Proposition 6.1 and Corollary 1.10 which, recall, states that, under the assumptions of Proposition 1.14, any weak- \star accumulation point ν of a sequence $(|\psi_\hbar|^2 \text{vol}_g)_{0 < \hbar \leq 1}$ verifying

$$P_\epsilon(\hbar)\psi_\hbar = E(\hbar)\psi_\hbar, \quad \text{and } E(\hbar) \rightarrow E_0 \neq 0,$$

must satisfy

$$\nu(\gamma_0) = 0,$$

for every geodesic γ_0 that is not included in $\mathcal{C}(V)$. This shows in particular that, as soon as $\mathcal{C}(V) \neq M$, one has

$$\{\delta_\gamma : \gamma \text{ geodesic of } (M, g)\} \not\subset \mathcal{N}(\infty),$$

which contradicts Proposition 6.1.

We now give the proof of Proposition 6.1 which, for the reader's convenience, has been divided in several steps.

Let γ_0 be a geodesic of (M, g) ; let Γ_0 be a lift of γ_0 to \mathring{T}^*M such that, for $(x_0, \xi_0) \in \Gamma_0$ one has $\frac{\|\xi_0\|_{x_0}^2}{2} = E_0$. Our goal is to show that $\delta_{\gamma_0} \in \mathcal{N}(\infty)$ by constructing a sequence of eigenfunctions of $P_\epsilon(\hbar)$ with eigenvalues approaching E_0 such that its semiclassical measure is δ_{Γ_0} .

Suppose that $s_{E_0, \delta_0}(\hbar)$ satisfies the hypothesis of Proposition 6.1. Then it is possible to find a sequence $\tau_{\hbar_n} \rightarrow +\infty$ satisfying

$$(33) \quad \tau_{\hbar_n} \frac{s_{E_0, \delta_0}(\hbar_n)}{\hbar_n} \rightarrow +\infty,$$

and

$$(34) \quad \tau_{\hbar_n} \epsilon_{\hbar_n}^2 \rightarrow 0.$$

In the following, we omit the subscript n in order to alleviate the notations.

6.1. A sequence of “good” initial data. We fix a sequence of initial data $(u_{x_0, \xi_0}^{\hbar})_{0 < \hbar \leq 1}$ whose semiclassical measure is unique and given by the Dirac measure δ_{x_0, ξ_0} . This can be obtained from a sequence of normalized coherent states – see for instance Ch. 5 in [56]. Then, we pick a cutoff function $0 \leq \chi \leq 1$ in $C_c^\infty(\mathbb{R})$ which is equal to 1 in a small neighborhood of 0, say $[-1/2, 1/2]$, and 0 outside a slightly larger neighborhood, say $[-1, 1]$. We introduce the following truncation of our coherent states:

$$\tilde{u}_{x_0, \xi_0}^{\hbar} := \chi \left(\frac{P_0(\hbar) - E_0}{\delta_0} \right) u_{x_0, \xi_0}^{\hbar}.$$

Using results on functional calculus for semiclassical pseudodifferential operators – see for instance Ch. 14 in [56], one can verify that the operator $\chi \left(\frac{P_0(\hbar) - E_0}{\delta_0} \right)$ is an \hbar -pseudodifferential operator belonging to the set $\Psi^{-\infty, 0}(M)$ as defined in Appendix A. Recall that its principal symbol is

$$\chi_{\hbar}^0(x, \xi) := \chi \left(\frac{\|\xi\|^2/2 - E_0}{\delta_0} \right).$$

As the semiclassical measure of the sequence $(u_{x_0, \xi_0}^{\hbar})_{0 < \hbar \leq 1}$ is the Dirac measure in (x_0, ξ_0) , one can observe that

$$(35) \quad \|u_{x_0, \xi_0}^{\hbar} - \tilde{u}_{x_0, \xi_0}^{\hbar}\|_{L^2(M)} = o(1).$$

Thanks to the Calderón-Vaillancourt Theorem, one can also verify that

$$(36) \quad \left\| \chi \left(\frac{P_0(\hbar) - E_0}{\delta_0} \right) - \chi \left(\frac{P_{\epsilon}(\hbar) - E_0}{\delta_0} \right) \right\|_{L^2(M) \rightarrow L^2(M)} = o(1).$$

Denote by $\Pi_{E_j(\hbar)}$ the spectral projector corresponding to the eigenvalue $E_j(\hbar)$, one has

$$\tilde{u}_{x_0, \xi_0}^{\hbar, \epsilon} := \chi \left(\frac{P_{\epsilon}(\hbar) - E_0}{\delta_0} \right) u_{x_0, \xi_0}^{\hbar} = \sum_{j=1}^{N(\hbar)} \chi \left(\frac{E_j(\hbar) - E_0}{\delta_0} \right) \Pi_{E_j(\hbar)} u_{x_0, \xi_0}^{\hbar}.$$

Recall that the eigenvalues $E_j(\hbar)$ (and the spectral projectors) depend implicitly on ϵ_{\hbar} . When $\Pi_{E_j(\hbar)} u_{x_0, \xi_0}^{\hbar} \neq 0$, we define

$$v_{\hbar, \epsilon}^j := \frac{\Pi_{E_j(\hbar)} u_{x_0, \xi_0}^{\hbar}}{\|\Pi_{E_j(\hbar)} u_{x_0, \xi_0}^{\hbar}\|_{L^2}},$$

otherwise, we set $v_{\hbar, \epsilon}^j = 0$. Then, we write

$$\tilde{u}_{x_0, \xi_0}^{\hbar, \epsilon} = \sum_{j=1}^{N(\hbar)} \chi \left(\frac{E_j(\hbar) - E_0}{\delta_0} \right) \|\Pi_{E_j(\hbar)} u_{x_0, \xi_0}^{\hbar}\|_{L^2(M)} v_{\hbar, \epsilon}^j.$$

Finally, we introduce $c_{\hbar,\epsilon}^j := \chi \left(\frac{E_j(\hbar) - E_0}{\delta_0} \right)^2 \|\Pi_{E_j(\hbar)} u_{x_0, \xi_0}^\hbar\|_{L^2(M)}^2$. Note that this quantity is equal to 0 when $v_{\hbar,\epsilon}^j = 0$. Using (35) and (36), one can verify that

$$(37) \quad \sum_{j=1}^{N(\hbar)} c_{\hbar,\epsilon}^j = 1 + o(1).$$

6.2. Semiclassical measures of the Schrödinger equation. Let θ be a smooth function on \mathbb{R} such that $\hat{\theta}$ is compactly supported and satisfies $\hat{\theta}(0) = 1$. We also fix b in $\mathcal{C}_c^\infty(\mathring{T}^*M)$. According to Propositions 2.2 and 2.3 (point i) and to the fact that $\tau_\hbar \epsilon_\hbar^2 \rightarrow 0$, one has

$$\int_{\mathbb{R}} \theta(t) \left\langle u_{x_0, \xi_0}^\hbar, e^{\frac{it\tau_\hbar P_\epsilon(\hbar)}{\hbar}} \text{Op}_\hbar(b) e^{\frac{it\tau_\hbar P_\epsilon(\hbar)}{\hbar}} u_{x_0, \xi_0}^\hbar \right\rangle dt = \mathcal{I}_g(b)(x_0, \xi_0) + o(1),$$

as \hbar goes to 0. Using the conventions introduced in the previous paragraph, one deduces then

$$\sum_{1 \leq j, k \leq N(\hbar)} \hat{\theta} \left(\frac{\tau_\hbar(E_j(\hbar) - E_k(\hbar))}{\hbar} \right) (c_{\hbar,\epsilon}^j c_{\hbar,\epsilon}^k)^{\frac{1}{2}} \langle v_{\hbar,\epsilon}^j, \text{Op}_\hbar(b) v_{\hbar,\epsilon}^k \rangle = \mathcal{I}_g(b)(x_0, \xi_0) + o(1).$$

For $j \neq k$, one has

$$\frac{\tau_\hbar}{\hbar} |E_j(\hbar) - E_k(\hbar)| \geq \tau_\hbar \frac{s_{E_0, \delta_0}(\hbar)}{\hbar},$$

which tends to $+\infty$ according to (33). Thus, we get, as \hbar goes to 0,

$$(38) \quad \sum_{j=1}^{N(\hbar)} c_{\hbar,\epsilon}^j \langle v_{\hbar,\epsilon}^j, \text{Op}_\hbar(b) v_{\hbar,\epsilon}^j \rangle = \mathcal{I}_g(b)(x_0, \xi_0) + o(1).$$

Recall that $v_{\hbar,\epsilon}^j$ is either 0 or a normalized state in $L^2(M)$ which satisfies $P_\epsilon(\hbar) v_{\hbar,\epsilon}^j = E_j(\hbar) v_{\hbar,\epsilon}^j$. We will deduce from this asymptotic formula the existence of an accumulation element $\mu \in \mathcal{M}(\infty)$ satisfying $\mu(\Gamma_0) = 1$, which by invariance considerations must be equal to δ_{Γ_0} .

6.3. Estimating the variance. We now remark that the measure $b \mapsto \mathcal{I}_g(b)(x_0, \xi_0)$ is an ergodic measure which can be written as a convex sum of almost invariant and almost positive distribution. Thus, we can proceed as in the proof of the quantum ergodicity Theorem [56] to construct a subsequence converging to this measure. For that purpose, we start by estimating the variance:

$$V_\hbar(b) := \sum_{j=1}^{N(\hbar)} c_{\hbar,\epsilon}^j \left| \langle v_{\hbar,\epsilon}^j, \text{Op}_\hbar(b) v_{\hbar,\epsilon}^j \rangle - \mathcal{I}_g(b)(x_0, \xi_0) \right|^2.$$

One can introduce χ_1 a smooth compactly supported function on \mathring{T}^*M which is equal to 1 for $\frac{E_0}{2} \leq \|\xi\|^2 \leq 8E_0$ and which is invariant by the geodesic flow. By construction of the

states $v_{\hbar,\epsilon}^j$, we can rewrite $V_{\hbar}(b)$ as follows:

$$V_{\hbar}(b) = \sum_{j=1}^{N(\hbar)} c_{\hbar,\epsilon}^j \left| \left\langle v_{\hbar,\epsilon}^j, \text{Op}_{\hbar}(\tilde{b}\chi_1)v_{\hbar,\epsilon}^j \right\rangle \right|^2 + \mathcal{O}(\hbar),$$

where $\tilde{b} = b - \mathcal{I}_g(b)(x_0, \xi_0)$. Combining the Egorov Theorem to the fact that the $v_{\hbar,\epsilon}^j$ are eigenmodes, one finds, for every $T > 0$,

$$V_{\hbar}(b) = \sum_{j=1}^{N(\hbar)} c_{\hbar,\epsilon}^j \left| \left\langle v_{\hbar,\epsilon}^j, \text{Op}_{\hbar} \left(\frac{1}{T} \int_0^T (\tilde{b}\chi_1) \circ \varphi^s ds \right) v_{\hbar,\epsilon}^j \right\rangle \right|^2 + \mathcal{O}(\hbar),$$

where the constant in the remainder depends on T . Applying the Cauchy-Schwarz inequality and the composition rule for pseudodifferential operators, we obtain

$$V_{\hbar}(b) \leq \sum_{j=1}^{N(\hbar)} c_{\hbar,\epsilon}^j \left\langle v_{\hbar,\epsilon}^j, \text{Op}_{\hbar} \left(\left| \frac{1}{T} \int_0^T (\tilde{b}\chi_1) \circ \varphi^s ds \right|^2 \right) v_{\hbar,\epsilon}^j \right\rangle + \mathcal{O}(\hbar),$$

We now use the limit formula (38), and we derive that, for every $T > 0$,

$$\limsup_{\hbar \rightarrow 0} V_{\hbar}(b) \leq \mathcal{I}_g \left(\left| \frac{1}{T} \int_0^T (\tilde{b}\chi_1) \circ \varphi^s ds \right|^2 \right) (x_0, \xi_0).$$

We take the limit $T \rightarrow +\infty$, and we have, as $\hbar \rightarrow 0^+$,

$$(39) \quad \sum_{j=1}^{N(\hbar)} c_{\hbar,\epsilon}^j \left| \left\langle v_{\hbar,\epsilon}^j, \text{Op}_{\hbar}(b)v_{\hbar,\epsilon}^j \right\rangle - \mathcal{I}_g(b)(x_0, \xi_0) \right|^2 = o(1).$$

6.4. Bienaymé-Tchebychev inequality. As the eigenmodes are microlocalized on the energy layers $E_0 - \delta_0 \leq \frac{\|\xi\|_x^2}{2} \leq E_0 + \delta_0$, we fix $(b_k)_{k \in \mathbb{N}}$ a family of smooth functions in $\mathcal{C}_c^\infty(\dot{T}^*M)$ which defines by restriction a dense subset (in the \mathcal{C}^0 topology) of the continuous functions on $\{(x, \xi) \in T^*M : [E_0 - 2\delta_0, E_0 + 2\delta_0]\}$. We reestablish the dependence in the parameter n and we define, for every k in \mathbb{N} and every n in \mathbb{N} ,

$$A_{\hbar_n}^k := \left\{ 1 \leq j \leq N(\hbar_n) : \left| \left\langle v_{\hbar_n,\epsilon_n}^j, \text{Op}_{\hbar_n}(b_k)v_{\hbar_n,\epsilon_n}^j \right\rangle - \mathcal{I}_g(b_k)(x_0, \xi_0) \right|^2 \geq V_{\hbar_n}(b_k)^{1/2} \right\}.$$

From the Bienaymé-Tchebychev inequality, we get, as $n \rightarrow +\infty$,

$$\sum_{j \in A_{\hbar_n}^k} c_{\hbar_n,\epsilon_n}^j = o(1).$$

For every $k \geq 0$, we set $\mathbf{A}_{\hbar_n}^k := \cup_{l=0}^k A_{\hbar_n}^l$ which still satisfies

$$\sum_{j \in \mathbf{A}_{\hbar_n}^k} c_{\hbar_n,\epsilon_n}^j = o(1).$$

For every $k \in \mathbb{N}$, we take n_k large enough to ensure that, for every $n \geq k$, one has

$$\left| \sum_{j \in \mathbf{A}_{\hbar_n}^k} c_{\hbar_n, \epsilon_n}^j \right| \leq \frac{1}{k},$$

and we also impose the subsequence n_k to be increasing. Then, for every $n_k \leq n < n_{k+1}$, we let $\mathbf{A}_{\hbar_n} := \mathbf{A}_{\hbar_n}^k$. Then, thanks to (37), one has, as $n \rightarrow +\infty$,

$$\sum_{j \in \mathbf{A}_{\hbar_n}^c} c_{\hbar_n, \epsilon_n}^j = 1 + o(1).$$

By construction, this implies that there exists a sequence $1 \leq j_n \leq N(\hbar_n)$ such that, for every k in \mathbb{N} ,

$$\lim_{n \rightarrow +\infty} \langle v_{\hbar_n, \epsilon_n}^{j_n}, \text{Op}_{\hbar_n}(b_k) v_{\hbar_n, \epsilon_n}^{j_n} \rangle = \mathcal{I}_g(b_k)(x_0, \xi_0).$$

By density of the family b_k in the \mathcal{C}^0 -topology, we find that the limit measure is the measure carried by the closed geodesic issued from (x_0, ξ_0) . In other words, we have constructed a sequence of eigenmodes whose semiclassical measure is carried by the closed geodesic issued from (x_0, ξ_0) . Since the semiclassical measure is invariant by the geodesic flow, it must coincide with the uniform orbit measure δ_{Γ_0} . This concludes the proof of Proposition 6.1.

APPENDIX A. SEMICLASSICAL ANALYSIS ON MANIFOLDS

In this appendix, we review some basic facts on semiclassical analysis that can be found for instance in [56]. Recall that we define on \mathbb{R}^{2d} the following class of symbols:

$$S^{m,k}(\mathbb{R}^{2d}) := \left\{ (b_{\hbar}(x, \xi))_{\hbar \in (0,1]} \in \mathcal{C}^\infty(\mathbb{R}^{2d}) : |\partial_x^\alpha \partial_\xi^\beta b_{\hbar}| \leq C_{\alpha,\beta} \hbar^{-k} \langle \xi \rangle^{m-|\beta|} \right\}.$$

Let M be a smooth Riemannian d -manifold without boundary. Consider a smooth atlas (f_l, V_l) of M , where each f_l is a smooth diffeomorphism from $V_l \subset M$ to a bounded open set $W_l \subset \mathbb{R}^d$. To each f_l correspond a pull back $f_l^* : \mathcal{C}^\infty(W_l) \rightarrow \mathcal{C}^\infty(V_l)$ and a canonical map \tilde{f}_l from T^*V_l to T^*W_l :

$$\tilde{f}_l : (x, \xi) \mapsto (f_l(x), (Df_l(x)^{-1})^T \xi).$$

Consider now a smooth locally finite partition of identity (ϕ_l) adapted to the previous atlas (f_l, V_l) . That means $\sum_l \phi_l = 1$ and $\phi_l \in \mathcal{C}^\infty(V_l)$. Then, any observable b in $\mathcal{C}^\infty(T^*M)$ can be decomposed as $b = \sum_l b_l$, where $b_l = b\phi_l$. Each b_l belongs to $\mathcal{C}^\infty(T^*V_l)$ and can be pushed to a function $\tilde{b}_l = (\tilde{f}_l^{-1})^* b_l \in \mathcal{C}^\infty(T^*W_l)$. As in [56], define the class of symbols of order m and index k

$$(40) \quad S^{m,k}(T^*M) := \left\{ (b_{\hbar}(x, \xi))_{\hbar \in (0,1]} \in \mathcal{C}^\infty(T^*M) : |\partial_x^\alpha \partial_\xi^\beta b_{\hbar}| \leq C_{\alpha,\beta} \hbar^{-k} \langle \xi \rangle^{m-|\beta|} \right\}.$$

Then, for $b \in S^{m,k}(T^*M)$ and for each l , one can associate to the symbol $\tilde{b}_l \in S^{m,k}(\mathbb{R}^{2d})$ the standard Weyl quantization

$$\text{Op}_\hbar^w(\tilde{b}_l)u(x) := \frac{1}{(2\pi\hbar)^d} \int_{\mathbb{R}^{2d}} e^{\frac{i}{\hbar}\langle x-y, \xi \rangle} \tilde{b}_l\left(\frac{x+y}{2}, \xi; \hbar\right) u(y) dy d\xi,$$

where $u \in \mathcal{S}(\mathbb{R}^d)$, the Schwartz class. Consider now a smooth cutoff $\psi_l \in \mathcal{C}_c^\infty(V_l)$ such that $\psi_l = 1$ close to the support of ϕ_l . A quantization of $b \in S^{m,k}(T^*M)$ is then defined in the following way [56]:

$$(41) \quad \text{Op}_\hbar(b)(u) := \sum_l \psi_l \times \left(f_l^* \text{Op}_\hbar^w(\tilde{b}_l)(f_l^{-1})^* \right) (\psi_l \times u),$$

where $u \in \mathcal{C}^\infty(M)$. This quantization procedure Op_\hbar sends (modulo $\mathcal{O}(\hbar^\infty)$) $S^{m,k}(T^*M)$ onto the space of pseudodifferential operators of order m and of index k , denoted $\Psi^{m,k}(M)$ [56]. It can be shown that the dependence in the cutoffs ϕ_l and ψ_l only appears at order 1 in \hbar (Theorem 9.10 in [56]) and the principal symbol map $\sigma_0 : \Psi^{m,k}(M) \rightarrow S^{m,k}/S^{m-1,k-1}(T^*M)$ is then intrinsically defined. Most of the rules (for example the composition of operators, the Egorov and Calderón-Vaillancourt Theorems) that hold on \mathbb{R}^{2d} still hold in the case of $\Psi^{m,k}(M)$. Finally, we denote by $\Psi^{-\infty,k}(M)$ the set $\bigcap_{m \in \mathbb{R}} \Psi^{m,k}(M)$.

APPENDIX B. TIME-DEPENDENT SEMICLASSICAL MEASURES

The aim of this short appendix is to recall a few facts on the definition of time-dependent semiclassical measures – we refer to [38] for more details.

Let $(u_\hbar)_{0 < \hbar \leq 1}$ be a normalized sequence in $L^2(M)$ verifying the oscillation assumptions (6) and (7). For a given scale of times $\tau := (\tau_\hbar)_{\hbar \rightarrow 0^+}$ satisfying

$$\lim_{\hbar \rightarrow 0^+} \tau_\hbar = +\infty,$$

we denoted the Wigner distribution by

$$(42) \quad \forall a \in \mathcal{C}_c^\infty(T^*M), \quad w_\hbar(t\tau_\hbar)(a) := \langle v_\hbar(t\tau_\hbar), \text{Op}_\hbar^w(a)v_\hbar(t\tau_\hbar) \rangle,$$

where $v_\hbar(\tau')$ is the solution at time τ' of (5) with initial condition u_\hbar . Using the Calderón-Vaillancourt Theorem, we deduce the existence of a constant $C > 0$ and a positive integer D depending only on the manifold (M, g) such that

$$\left| \int_{\mathbb{R} \times T^*M} a(t, x, \xi) w_\hbar(t\tau_\hbar, dx, d\xi) dt \right| \leq C_K \sum_{|\alpha| \leq D} \hbar^{|\alpha|/2} \int_{\mathbb{R}} \|\partial_{x,\xi}^\alpha a(t, \cdot)\|_{\mathcal{C}^0(T^*M)} dt,$$

for every $a \in \mathcal{C}_c^\infty(\mathbb{R} \times T^*M)$. According to [46] (Ch. 3), the sequence $(\mu_\hbar : (t, x, \xi) \mapsto w_\hbar(t\tau_\hbar, x, \xi))_{\hbar > 0}$ is relatively compact in $\mathcal{D}'(\mathbb{R} \times T^*M)$. Thus, we can extract converging subsequences (for the weak- \star topology). In particular, for any accumulation point μ of this sequence and every $a \in \mathcal{C}_c^\infty(\mathbb{R} \times T^*M)$, one has

$$\left| \int_{\mathbb{R} \times T^*M} a(t, x, \xi) \mu(dt, dx, d\xi) \right| \leq C \int_{\mathbb{R}} \|a(t, \cdot)\|_{\mathcal{C}^0(T^*M)} dt.$$

Thus, μ can be extended to a continuous linear form on $L^1(\mathbb{R}, \mathcal{C}_0^0(T^*M))$, where $\mathcal{C}_0^0(T^*M)$ denotes the set of continuous functions vanishing at infinity. Consequently, the limit distribution $t \mapsto \mu(t, \cdot)$ belongs to $L^\infty(\mathbb{R}, \mathcal{M}(T^*M))$ (see for instance [11]), where $\mathcal{M}(T^*M)$ is the set of finite complex measures on T^*M . For any converging subsequence in $\mathcal{D}'(\mathbb{R} \times T^*M)$ (which we do not relabel), we note that the following also holds, for every $\theta \in L^1(\mathbb{R})$, and for every $a \in \mathcal{C}_c^\infty(T^*M)$

$$\lim_{h \rightarrow 0^+} \int_{\mathbb{R} \times T^*M} \theta(t) a(x, \xi) w_h(t\tau_h, dx, d\xi) dt = \int_{\mathbb{R} \times T^*M} \theta(t) a(x, \xi) \mu(t, dx, d\xi) dt.$$

Finally², according to the Gårding inequality, the limit distribution is in fact a positive measure for a.e. t in \mathbb{R} . Moreover, the frequency assumptions (6) and (7) and the fact that $\epsilon_h \rightarrow 0^+$ imply that, for every almost every t , $\mu(t)(\mathring{T}^*M) = 1$. Using Egorov Theorem, one can also verify that, for a.e. t in \mathbb{R} , $\mu(t, \cdot)$ is invariant by the geodesic flow φ^s .

APPENDIX C. GEOMETRY OF T^*M

In this appendix, we collect some classical results on Riemannian and symplectic geometry that appear at different stages of this work. Along the way, we recall classical notations that are used all along this article. We refer for instance the reader to [4, 42, 45, 50] for more details.

C.1. Musical isomorphisms. Recall that the Riemannian metric g on M induces two natural isomorphisms

$$\flat : T_x M \rightarrow T_x^* M, \quad v \mapsto g_x(v, \cdot),$$

and its inverse $\sharp : T_x^* M \rightarrow T_x M$. This natural isomorphism induces a positive definite form on $T_x^* M$ for which these isomorphisms are in fact isometries. We denote by g^* the corresponding metric.

C.2. Horizontal and vertical subbundles. Let $\rho = (x, \xi)$ be an element in T^*M . Denote by $\pi : T^*M \rightarrow M$ the canonical projection $(x, \xi) \mapsto x$. We introduce the so-called *vertical* subspace:

$$\mathcal{V}_\rho := \text{Ker}(d_\rho \pi) \subset T_\rho T^*M.$$

The fiber T_x^*M is a submanifold of T^*M that contains the point (x, ξ) . The tangent space to this submanifold at point (x, ξ) is the vertical subspace \mathcal{V}_ρ and it can be canonically identified with T_x^*M . We will now define the connection map. For that purpose, we fix Z in $T_\rho T^*M$ and $\rho(t) = (x(t), \xi(t))$ a smooth curve in T^*M such that $\rho(0) = \rho$ and $\rho'(0) = Z$. The connection map $\mathcal{K}_\rho : T_\rho T^*M \mapsto T_x^*M$ is the following application:

$$\mathcal{K}_\rho(Z) := \frac{D}{dt} \rho(0) = \nabla_{x'(0)} \xi(0),$$

where $\frac{D}{dt} \rho(t)$ is the covariant derivative of $\rho(t)$ along the curve $x(t)$. One can verify that this quantity depends only on the initial velocity Z of the curve (and not on the curve

²One can for instance follow the arguments given in Ch. 5 of [56].

itself) and the map is linear. The *horizontal* space is given by the kernel of this linear application, i.e.

$$\mathcal{H}_\rho := \text{Ker}(\mathcal{K}_\rho) \subset T_\rho T^*M.$$

There exists a natural vector bundle isomorphism between the pullback bundle $\pi^*(TM \oplus T^*M) \rightarrow T^*M$ and the canonical bundle $TT^*M \rightarrow T^*M$. The restriction of this isomorphism on the fibers above $\rho \in T^*M$ is given by

$$\theta(\rho) : T_\rho T^*M \rightarrow T_{\pi(\rho)}M \oplus T_{\pi(\rho)}^*M, \quad Z \mapsto (y, \eta) := (d_\rho \pi(Z), \mathcal{K}_\rho(Z)).$$

These coordinates (y, η) will allow us to express easily the different structures on T^*M . For instance, the Hamiltonian vector field X associated to p_0 (i.e. the generator of the geodesic flow) satisfies $\theta(\rho)X(\rho) = (\pi(\rho)^\sharp, 0)$.

C.3. Symplectic structure on T^*M . Recall that the canonical contact form on T^*M is given by the following expression:

$$\forall \rho = (x, \xi) \in T^*M, \quad \forall Z \in T_\rho T^*M, \quad \alpha_{x, \xi}(Z) = \xi(d_\rho \pi(Z)).$$

The canonical symplectic form on T^*M can then be defined as $\Omega = d\alpha$. Using our natural isomorphism, this symplectic form can be written as

$$\forall Z_1 \cong (y_1, \eta_1) \in T_\rho T^*M, \quad \forall Z_2 \cong (y_2, \eta_2) \in T_\rho T^*M, \quad \Omega_\rho(Z_1, Z_2) = \eta_1(y_2) - \eta_2(y_1).$$

C.4. Almost complex structure on T^*M . One can define the following map from $T_x M \oplus T_x^*M$ to itself:

$$\tilde{J}_x(y, \eta) = (\eta^\sharp, -y^\flat).$$

This map induces an almost complex structure on $T_\rho T^*M$ through the isomorphism $\theta(\rho)$. We denote this almost complex structure by J_ρ .

C.5. Riemannian metric on T^*M . The Sasaki metric g^S on T^*M is then defined as

$$g_\rho^S(Z_1, Z_2) := g_x^*(\mathcal{K}_\rho(Z_1), \mathcal{K}_\rho(Z_2)) + g_x(d_\rho \pi(Z_1), d_\rho \pi(Z_2)).$$

This is a positive definite bilinear form on $T_\rho T^*M$. The important point is that *this metric is compatible with the symplectic structure on T^*M through the almost complex structure*. Precisely, one has, for every $(Z_1, Z_2) \in T_\rho T^*M \times T_\rho T^*M$,

$$g_\rho^S(Z_1, Z_2) = \Omega_\rho(Z_1, J_\rho Z_2).$$

In fact, using the natural isomorphism, one has

$$\Omega_\rho(Z_1, J_\rho Z_2) = \eta_1(\eta_2^\sharp) + y_2^\flat(y_1) = g_x^*(\eta_1, \eta_2) + g_x(y_1, y_2).$$

C.6. A natural frame on T^*M . In the case M is an oriented surface, one can introduce a “natural basis” on $T_\rho T^*M$ as follows. Thanks to the fact that the manifold M is oriented with a Riemannian structure, one can define a notion of rotation by $\pi/2$ in every cotangent space T_x^*M (which is of dimension 2). Thus, given any $\xi \in T_x^*M \setminus \{0\}$ there exists a unique ξ^\perp such that $\{\xi, \xi^\perp\}$ is a direct orthogonal basis with $\|\xi\|_x = \|\xi^\perp\|_x$. We use this to define an orthogonal basis of \mathcal{V}_ρ for $\rho \in \mathring{T}^*M$:

$$Y_0(\rho) := (\theta(\rho))^{-1}(0, \xi), \text{ and } U(\rho) = (\theta(\rho))^{-1}(0, \xi^\perp).$$

Then, we can define an orthogonal basis of \mathcal{H}_ρ as follows

$$X_0(\rho) = J_\rho Y_0(\rho), \text{ and } W(\rho) = J_\rho U(\rho).$$

Note that X_0 is the geodesic vector field and that the family $\{X_0(\rho), W(\rho), Y_0(\rho), U(\rho)\}$ forms a direct orthogonal basis of $T_\rho \mathring{T}^*M$.

REFERENCES

- [1] N. Anantharaman, C. Fermanian-Kammerer, F. Macià *Long-time dynamics of completely integrable Schrödinger flows on the torus*, American J. Math. **137** (2015).
- [2] N. Anantharaman, F. Macià *Semiclassical measures for the Schrödinger equation on the torus*, JEMS, **16** (2014), 1253–1288.
- [3] D. Azagra, F. Macià *Concentration of symmetric eigenfunctions*, Nonlinear Anal. **73** (2010), 683–688.
- [4] A. Besse *Manifolds All of Whose Geodesics Are Closed*, Ergeb. Math. **93**, Springer-Verlag, New York, (1978)
- [5] M.L. Bialy, L. V. Polterovich *Lagrangian singularities of invariant tori of Hamiltonian systems with two degrees of freedom*, Invent. Math. **97** (1989), 291–303.
- [6] J. Bolte, T. Schwaibold *Stability of wave packet dynamics under perturbations*, Phys. Rev. E73 (2006) 026223.
- [7] S. Brooks, E. Le Masson, E. Lindenstrauss *Quantum Ergodicity and Averaging Operators on the Sphere*, arXiv:1505.03887 (2015)
- [8] J. Chazarain *Spectre d’un hamiltonien quantique et mécanique classique*, CPDE **6** (1980), 595–644
- [9] Y. Colin de Verdière *Sur le spectre des opérateurs elliptiques à bicaractéristiques toutes périodiques*, Comment. Math. Helv. **54** (1979), 508–522
- [10] M. Combes, D. Robert *A phase-space study of the quantum Loschmidt Echo in the semiclassical limit*, Ann. H. Poincaré **8** (2007), 91–108
- [11] J. Diestel and J.J. Uhl *Vector measures*. Mathematical Surveys, No. 15. American Mathematical Society, Providence, R.I., 1977.
- [12] M. Dimassi, J. Sjöstrand *Spectral asymptotics in the semi-classical limit*, London Math. Soc. Lecture notes series **268**, Cambridge University Press (1999)
- [13] J.J. Duistermaat *On global action-angle coordinates*, Comm. Pure Appl. Math. **33** (1980), 687–706.
- [14] J.J. Duistermaat and V. Guillemin *The spectrum of elliptic operators and periodic bicharacteristics*, Inv. Math. **29** (1975), 39–79
- [15] M. Einsiedler, T. Ward *Ergodic theory with a view towards number theory*, Graduate Texts in Mathematics **259**, Springer-Verlag London (2011)
- [16] S. Eswarathan, G. Rivière *Perturbation of the semiclassical Schrödinger equation on negatively curved surfaces*, J. Inst. Math. Jussieu, on line (2015)
- [17] P. Gérard *Mesures semi-classiques et ondes de Bloch*, Sem. EDP (Polytechnique) 1990–1991, Exp. 16 (1991)

- [18] P. Gérard *Description du défaut de compacité de l'injection de Sobolev*, ESAIM Control Optim. Calc. Var., **3** (1998) 213–233.
- [19] T. Gorin, T. Prosen, T.H. Seligman, M. Znidaric *Dynamics of Loschmidt echoes and fidelity decay*, Physics Reports **435** (2006) 33–156.
- [20] A. Goussev, R.A. Jalabert, H.M. Pastawski, D. Wisniacki *Loschmidt Echo*, Scholarpedia 7(8), 11687, arXiv:1206.6348 (2012)
- [21] D. Gromoll, K. Grove, *On metrics on S^2 all of whose geodesics are closed*, Invent. Math. **65** (1981/82), 175–177.
- [22] V. Guillemin *The Radon transform on Zoll surfaces*, Adv. Math. **22** (1976), 85–119.
- [23] V. Guillemin *Some spectral results for the Laplace operator with potential on the n -sphere*, Adv. in Math. **27** (1978), 273–286.
- [24] V. Guillemin *Some spectral results on rank one symmetric spaces*, Adv. in Math. **28** (1978), 129–137.
- [25] V. Guillemin *Band asymptotics in 2 dimension*, Adv. Math. **42** (1981), 248–282.
- [26] M.A. Hall, M. Hitrik, J. Sjöstrand *Spectra for semiclassical operators with periodic bicharacteristics in dimension two*, arXiv:1401.3371 (2014).
- [27] A. Hatcher *Algebraic topology*, Cambridge University Press (2002).
- [28] B. Helffer, D. Robert *Puits de potentiel généralisés et asymptotique semi-classique*, Ann. Inst. H. Poincaré Phys. Théor. **41** (1984), 291–331.
- [29] S. Helgason *Integral geometry and Radon transforms*. Springer, New York, 2011.
- [30] M. Hitrik *Eigenfrequencies for Damped Wave Equations on Zoll Manifolds*, Asymptot. Analysis **31** (2002), 265–277
- [31] M. Hitrik and J. Sjöstrand, *Non-selfadjoint perturbations of selfadjoint operators in 2 dimensions I*, Ann. Henri Poincaré **5** (2004), 1–73.
- [32] L. Hörmander, *The analysis of linear partial differential operators III*, Springer-Verlag Berlin Heidelberg New York Tokyo (1985)
- [33] L. Hörmander, *The analysis of linear partial differential operators IV*, Springer-Verlag Berlin Heidelberg New York Tokyo (1985)
- [34] P. Jacquod, C. Petitjean *Decoherence, Entanglement and Irreversibility in Quantum Dynamical Systems with Few Degrees of Freedom*, Adv. Phys. **58**, 67–196 (2009)
- [35] D. Jakobson and S. Zelditch *Classical limits of eigenfunctions for some completely integrable systems*, Emerging applications of number theory (Minneapolis, MN, 1996), 329–354, IMA Vol. Math. Appl., **109**, Springer, New York (1999)
- [36] B. Küster, P. Ramacher *Quantum ergodicity and symmetry reduction*, arXiv:1410.1096 (2014)
- [37] F. Macià *Some remarks on quantum limits on Zoll manifolds*, CPDE **33** (2008), 1137–1146
- [38] F. Macià *Semiclassical measures and the Schrödinger flow on Riemannian manifolds*, Nonlinearity **22** (2009), 1003–1020
- [39] F. Macià, G. Rivière *Semiclassical measures for the perturbed Schrödinger equation on \mathbb{T}^d* , in preparation
- [40] J. Moser, E. Zehnder, *Notes on dynamical systems*, Courant Lecture Notes in Mathematics, **12**. American Mathematical Society, Providence, RI, (2005).
- [41] D. Ojeda-Valencia, C. Villegas-Blas *Limiting eigenvalue distributions Theorems in semiclassical analysis*, Spectral Analysis of Quantum Hamiltonians: Spectral Days 2010, Springer
- [42] G.P. Paternáin *Geodesic Flows*, Progress in Mathematics, **180**. Birkhäuser Boston, Inc., Boston, MA, (1999).
- [43] A. Peres *Stability of quantum motion in chaotic and regular systems*, Phys. Rev. A **30** (1984), 1610–1615
- [44] G. Rivière *Long time dynamics of the perturbed Schrödinger equation on negatively curved surfaces*, arXiv:1412.4400 (2014)
- [45] R.O. Ruggiero *Dynamics and global geometry of manifolds without conjugate points*, Ensaios Mat. Vol. **12**, Soc. Bras. Mat. (2007)

- [46] L. Schwartz *Théorie des distributions*, Hermann, Paris (1966)
- [47] A. Uribe *Some spectral results on rank one symmetric spaces*, Adv. in Math. **58**, 285–299 (1985)
- [48] A. Uribe and S. Zelditch *Spectral statistics on Zoll surfaces*, Comm. Math. Phys. **154** (1993), 313–346
- [49] J.M. VanderKam *L^∞ norms and quantum ergodicity on the sphere* IMRN **7** (1997), 329–347
- [50] J. Weber, *J-holomorphic curves in cotangent bundles and the heat flow*, Ph.D. thesis / Dissertation, TU Berlin (1999)
- [51] A. Weinstein *Asymptotics of eigenvalue clusters for the Laplacian plus a potential*, Duke Math. Jour. **44** (1977), 883–892
- [52] S. Zelditch *Quantum ergodicity on the sphere*, Comm. Math. Phys. **146** (1994), 61–71
- [53] S. Zelditch *Maximally degenerate Laplacians*, Ann. Inst. Fourier **46** (1996), 547–587
- [54] S. Zelditch *Fine structure of Zoll spectra*, J. Funct. Anal. **143** (1997), 415–460
- [55] S. Zelditch *Gaussian beams on Zoll manifolds and maximally degenerate Laplacians*, arXiv:1404.2906 (2014)
- [56] M. Zworski *Semiclassical analysis*, Graduate Studies in Mathematics **138**, AMS (2012)

UNIVERSIDAD POLITÉCNICA DE MADRID. DCAIN, ETSI NAVALES. AVDA. ARCO DE LA VICTORIA
S/N. 28040 MADRID, SPAIN

E-mail address: Fabricio.Macia@upm.es

LABORATOIRE PAUL PAINLEVÉ (U.M.R. CNRS 8524), U.F.R. DE MATHÉMATIQUES, UNIVERSITÉ
LILLE 1, 59655 VILLENEUVE D'ASCQ CEDEX, FRANCE

E-mail address: gabriel.riviere@math.univ-lille1.fr