DEFORMATION THEORY OF BIALGEBRAS, HIGHER HOCHSCHILD COHOMOLOGY AND FORMALITY

GRÉGORY GINOT, SINAN YALIN

Abstract. A first goal of this paper is to precisely relate the homotopy theories of bialgebras and E_2 -algebras. For this, we construct a conservative and fully faithful ∞-functor from pointed conilpotent homotopy bialgebras to augmented E_2 -algebras which consists in an appropriate "cobar" construction. Then we prove that the (derived) formal moduli problem of homotopy bialgebras structures on a bialgebra is equivalent to the (derived) formal moduli problem of E_2 -algebra structures on this "cobar" construction. We show consequently that the E_3 -algebra structure on the higher Hochschild complex of this cobar construction, given by the solution to the higher Deligne conjecture, controls the deformation theory of this bialgebra. This implies the existence of an E_3 -structure on the deformation complex of a dg bialgebra, solving a long-standing conjecture of Gerstenhaber-Schack. On this basis we solve a long-standing conjecture of Kontsevich, by proving the E_3 -formality of the deformation complex of the symmetric bialgebra. This provides as a corollary a new proof of Etingof-Kazdhan deformation quantization of Lie bialgebras which extends to homotopy dg Lie bialgebras and is independent from the choice of an associator. Along the way, we establish new general results of independent interest about the deformation theory of algebraic structures, which shed a new light on various deformation complexes and cohomology theories studied in the literature.

Contents

| Int | roduction | 1 |
|-----------|---|----|
| Notations | | 9 |
| 1. | Recollections | 10 |
| 2. | Moduli problems of algebraic structures and deformation complexes | 16 |
| 3. | The "plus" construction and the corresponding deformation theory | 30 |
| 4. | Bialgebras versus iterated coalgebras | 35 |
| 5. | Bialgebras versus E_2 -algebras | 48 |
| 6. | Identification of Deformation complexes with higher Hochschild, | |
| | Gerstenhaber-Schack and Tamarkin complexes of $Pois_n$ -algebras | 54 |
| 7. | The E_3 -formality Theorem | 67 |
| 8. | Etingof-Kazdhan deformation quantization | 75 |
| Re | References | |

Introduction

A deep interplay between bialgebras and quantum groups on one side and higher algebras on the other side (for instance E_2 and E_3 -algebras) has been noticed for

many years and more have been expected for a long time. This led people to investigate the relationship between these two kinds of structures with the hope to establish some equivalence between their deformation theories, which could help to understand various related problems on both sides. One of the first goals of this paper is to give a precise relation between their respective homotopy theories and to use it to study the deformation theory of dg-bialgebras.

Algebras governed by E_n -operads and their deformation theory play a prominent role in a variety of topics such as the study of iterated loop spaces, Goodwillie-Weiss calculus for embedding spaces, deformation quantization of Poisson manifolds and Lie bialgebras, and factorization homology [47, 48, 52, 54, 8, 19, 20, 26, 30, 36, 39, 40, 44, 49, 55, 62, 69, 72, 77]. These algebras form a hierarchy of "more and more" commutative and homotopy associative structures, interpolating between homotopy associative or A_{∞} -algebras (the E_1 case) and E_{∞} -algebras (the colimit of the E_n 's, which is homotopy equivalent to differential graded commutative algebras in characteristic zero). Their natural cohomology theory, the higher Hochschild cohomology, is a suitable generalization of the Hochschild cohomology of associative algebras.

On the other hand, bialgebras are central in various topics of algebraic topology, representation theory and mathematical physics (often incarnated as Hopf algebras) [9, 10, 5, 16, 17, 29, 58, 59]. They consist of two structures, an associative algebra structure and a coassociative coalgebra structure, related by a compatibility condition such that bialgebras can be alternately defined as algebras in coalgebras or vice-versa. Their natural cohomology theory, the Gerstenhaber-Schack cohomology, is a suitable mixing between Hochschild cohomology of algebras and co-Hochschild cohomology of coalgebras.

Two important and long-standing conjectures expected from the relationship between these two kind of structures are the following. The first one, enunciated by Gerstenhaber and Schack (in a wrong way) at the beginning of the 90's [29], characterizes the structure of the complex controlling the deformation theory of bialgebras, which remained quite mysterious for a while. It is a "differential graded bialgebra version" of the famous Deligne conjecture for associative differential graded algebras (see for instance [72] and [47]). The second one, enunciated by Kontsevich in his celebrated work on deformation quantization of Poisson manifolds [48], is a formality statement for the deformation complex of the symmetric bialgebra, which should imply as a corollary Drinfeld's and Etingof-Kazdhan's deformation quantization theory (see [9], [16] and [17]).

This situation is somehow a generalization to bialgebras of the situation encountered in deformation quantization of Poisson manifolds: the formality of the E_2 -operad implies the Deligne conjecture, which states the existence of an E_2 -algebra structure on the Hochschild complex of an associative algebra, and this allows to prove that the Hochschild complex of the ring of functions on a Poisson manifold is formal. This formality implies, in turn, the deformation quantization of such a manifold. Here, the formality of the E_3 -operad should allow to prove a bialgebra version of this Deligne conjecture, implying the formality of the Gerstenhaber-Schack complex of the symmetric bialgebra, inducing in turn Drinfeld and Etingof-Kazdhan deformation quantization of Lie bialgebras.

In this paper, we solve this bialgebra version of the Deligne conjecture proposed by Gerstenhaber-Schack as well as Kontsevich's E_3 -formality conjecture, both at a

greater level of generality than the original statements, by implementing methods coming from derived algebraic geometry (see in particular [52], [54] and [76]) as well as from previous results of the authors (see in particular [36] and [82]) and by solving partially a conjecture of Francis-Gaistgory.

We deduce from it a generalization of Etingof-Kadhan's celebrated deformation quantization.

Moreover, we enlighten along the way the role of various versions of deformation complexes of algebraic structures considered in the literature, by characterizing their associated derived formal moduli problems in full generality. Our new methods have thus an independent interest going even further than the solutions to the two aforementioned conjectures. We now describe our main results and conclude with some perspectives for future works.

0.1. From bialgebras to E_2 -algebras. The cobar construction defining a Quillen equivalence between conilpotent dg coalgebras and augmented dg algebras can be lifted to a functor from the category of associative coassociative bialgebras to the category of E_2 -algebras [43]. A natural question [19] was thus to know whether this implies an equivalence of homotopy theories (in the sense of $(\infty, 1)$ -categories) between bialgebras and E_2 -algebras. The appropriate answer to this problem is more subtle, and we solve it by using an appropriate notion of cobar construction for bialgebras, which intertwines a bar construction on the algebra part of the structure with a cobar construction on the coalgebra part:

Theorem 0.1. (1) There exists a bar-cobar adjunction

$$\mathcal{B}_{E_1}^{enh}: E_1 - Alg^{0-con}(dgCog^{conil}) \rightleftarrows E_1 - Cog^{conil}(dgCog^{conil}): \Omega_{E_1}^{enh}$$

inducing an equivalence of $(\infty, 1)$ -categories between 0-connected homotopy associative algebras in conilpotent dg coalgebras (0-connected conilpotent homotopy associative bialgebras) and conilpotent homotopy coassociative coalgebras in conilpotent dg coalgebras.

(2) The equivalence above induces an equivalence of $(\infty, 1)$ -categories

$$(\mathcal{B}^{enh,pt}_{E_1}(-)_-)_+:E_1-Alg^{aug,con}(dgCog^{conil})\rightleftarrows E_1-Cog^{conil,pt}(dgCog^{conil}):(\Omega^{enh,pt}_{E_1}(-)_-)_+$$

between connected augmented conilpotent homotopy associative bialgebras and pointed conilpotent homotopy coassociative coalgebras in conilpotent dg coalgebras.

This solves a conjecture of Francis-Gaitsgory [19] in the case where the base category is the category of conilpotent dg coalgebras, respectively the category of pointed conilpotent dg coalgebras. Let us note that we work here with non-negatively graded chain complexes.

Corollary 0.2. The left adjoint of Theorem 0.1(2) induces a fully faithful ∞ -functor

$$\tilde{\Omega}: E_1 - Alg^{aug,con}(dgCog^{conil}) \hookrightarrow E_2 - Alg^{aug}$$

embedding connected augmented conilpotent homotopy associative bialgebras into augmented E_2 -algebras.

We use the above results to study deformation of bialgebras in terms of those of E_2 -algebras, by the means of the theory of moduli spaces of algebraic structures.

0.2. Moduli spaces associated to deformation complexes. One of the main tool for studying deformation is given by deformation complexes of algebraic structures. In a part of this paper which is of independent interest, we prove several new results about deformation theory of algebraic structures. In particular, we give a conceptual explanation of the differences between various deformation complexes appearing in the literature by explaining which kind of deformations each of these complexes controls. We later use these results to identify the deformation complexes of dg-bialgebras with those of its (appropriate) cobar construction.

Deformation theory is classically encoded by moduli problem and their associated dg-Lie-algebras (for instance see [38, 48, 52, 63]). Formal moduli problems are simplicial presheaves over augmented artinian cdgas satisfying some extra properties with respect to homotopy pullbacks, which control the infinitesimal deformation theory of points on a given moduli space (variety, scheme, derived stack). According to Lurie's equivalence theorem [52], (derived) formal moduli problems and dg Lie algebras are equivalent as ∞-categories, so to a given formal moduli problem controlling the infinitesimal neighbourhood of a point on a moduli space corresponds a dg Lie algebra called the deformation complex of this point. Here we are interested in moduli spaces of algebraic structures and formal moduli problems controlling the deformations of such structures. A convenient formalism to deal with algebraic structures at a high level of generality, in order to encompass not only algebras but also bialgebras, is the notion of properad [79], a suitable generalization of operads. In the differential graded setting, algebraic structures are deformed as algebraic structures up to homotopy, for instance dg associative algebras are deformed as A_{∞} -algebras. Such a notion can be properly defined by considering cofibrant resolutions of properads. Given a cofibrant dg properad P_{∞} and a P_{∞} -algebra structure $\psi: P_{\infty} \to End_X$ on a complex X, there is a formal moduli problem $P_{\infty}\{X\}^{\psi}$ controlling the deformation theory of ψ . The associated deformation complex is an explicit Lie algebra noted $g_{P,X}^{\psi}$. In this paper, we prove the following general criterion to compare formal moduli problems between two kind of algebras:

Theorem 0.3. Let $F: P_{\infty} - Alg \to Q_{\infty} - Alg$ be a fully faithful and conservative ∞ -functor inducing functorially in A, for every augmented artinian edga A, a fully faithful and conservative ∞ -functor $F: P_{\infty} - Alg(Mod_A) \to Q_{\infty} - Alg(Mod_A)$. Then F induces an equivalence of formal moduli problems

$$\underline{P_{\infty}\{X\}}^{\psi} \sim \underline{Q_{\infty}\{F(X)\}}^{F(\psi)},$$

where $F(\psi)$ is the Q_{∞} -algebra structure on the image $F(X,\psi)$ of (X,ψ) under F, hence an equivalence of tangent L_{∞} -algebras

$$g_{P,X}^{\psi} \sim g_{Q,F(X)}^{F(\psi)}$$
.

Moreover, we get the following fiber sequence comparing the deformation complex of an algebra (X,ψ) with the homotopy Lie algebra $Lie(\underline{haut}_{P_{\infty}}(X,\psi))$ tangent to the derived algebraic group $\underline{haut}_{P_{\infty}}(X,\psi)$ (in the sense of [20]) of homotopy automorphisms of this algebra:

Proposition 0.4. There is a homotopy fiber sequence of L_{∞} -algebras

$$g_{P,X}^{\psi} \to Lie(\underline{haut}_{P_{\infty}}(X,\psi)) \to Lie(\underline{haut}(X)).$$

However, one should note that the deformation complex $g_{P,X}^{\psi}$ does not give exactly the usual cohomology theories. For instance, let us consider the case of the Hochschild complex Hom $(A^{\otimes >0},A)$ of a dg associative algebra A. This Hochschild complex is bigraded, with a cohomological grading induced by the grading of A and a weight grading given by the tensor powers $A^{\otimes \bullet}$. It turns out that the part Hom(A,A) of weight 1 in the Hochschild complex is missing in $g_{Ass,A}^{\psi}$.

To correct this, and in particular to get the correct deformation complexes, we use a "plus" construction $g_{P^+,X}^{\psi^+}$ which gives the right cohomology theory and can be obtained by a slight modification of the properad P, see § 3. Moreover, we relate it to the homotopy automorphisms of the algebra:

Theorem 0.5. There is a quasi-isomorphism of L_{∞} -algebras

$$g_{P^+,X}^{\psi^+} \simeq Lie(\underline{haut}_{P_{\infty}}(X,\psi)).$$

The conceptual explanation behind this phenomenon is that $g_{P,X}^{\psi}$ controls the deformations of the P_{∞} -algebra structure over a fixed complex X, whereas $g_{P^+,X}^{\psi^+}$ controls deformations of this P_{∞} -algebra structure plus compatible deformations of the differential of X, that is, deformations of the P_{∞} -algebra structure up to self quasi-isomorphisms of X (and in particular up to automorphisms). This is the role of the part Hom(X,X) appearing for instance in Hochschild cohomology.

For instance, in the case of a an associative dg algebra A, the complex $g_{Ass^+,A}^{\psi^+} \cong Hom(A^{\otimes >0},A)[1]$ computes the reduced Hochschild cohomology of A, where the right hand side is a sub-complex of the standard Hochschild cochain complex shifted down by 1 equipped with its standard Lie algebra structure (due to Gerstenhaber). The complex $g_{Ass,A}^{\psi} \cong Hom(A^{\otimes >1},A)[1]$ is the one controlling the formal moduli problem of deformations of A with fixed differential, where the right hand side is the subcomplex of the previous shifted Hochschild cochain complex where we have removed the Hom(A,A) component. There is also a third complex, the full shifted Hochschild complex $Hom(A^{\otimes \geq 0},A)[1)$, which controls not the deformations of A itself but the linear deformations of its dg category of modules Mod_A [46, 62].

In the case of n-Poisson algebras (Poisson algebras with a Poisson bracket of degree 1-n), we prove in Section 6 that the Tamarkin deformation complex [74] (which we denote $CH^{(\bullet>0)}_{Pois_n}(A)[n]$ since it is the part of positive weight in the full Poisson complex [7]) controls deformations of A into dg-Pois $_n$ -algebras, that is, it is the tangent Lie algebra $g^{\psi^+}_{Pois_n^+,A}$ of $\underline{haut}_{Pois_n}(A)$. We also prove that the deformation complex $g^{\psi}_{Pois_n,R}$ of the formal moduli problem $\underline{Pois}_{n\infty}\{A\}^{\psi}$ of homotopy n-Poisson algebra structures deforming ψ is given by the L_{∞} -algebra $CH^{(\bullet>1)}_{Pois_n}(A)[n]$, which is a further truncation of $CH_{Pois_n}(A)[n]$. Concerning the full shifted Poisson complex, we conjecture the following:

Conjecture. Let $n \geq 2$ and A be an n-Poisson algebra. The L_{∞} -algebra structure of the full shifted Poisson complex $CH_{Pois_n}^*(A)[n]$ controls the deformations of Mod_A into E_{n-1} -monoidal dg categories.

In the same way, we relate deformation complexes of bialgebras with truncations of the full Gerstenhaber-Schack complex (see § 6.2), and we conjecture the following:

Conjecture. Let B be a conilpotent dg bialgebra. The full shifted Gerstenhaber-Schack complex $C_{GS}^{full}(B,B)[2]$ controls the deformations of Mod_B as an E_2 -monoidal dg category, that is, a braided monoidal dg category.

0.3. Gerstenhaber-Schack conjecture. Another open problem is to prove that the higher Hochschild complex of the cobar construction of a bialgebra is a deformation complex of this bialgebra. It is important since it allows to reduce questions of deformations of the bialgebras to those of E_2 -structures for which more tools are available.

We deduce it as a consequence of a more general result stating the equivalence of formal moduli problems in the sense of Lurie [54] between the moduli problem $\underline{Bialg_{\infty}\{B\}}^{\varphi}$ of homotopy bialgebra structures on a bialgebra B deforming the bialgebra structure $\varphi:Bialg\to End_B$ (where Bialg is the prop of bialgebras) and the moduli problem $\underline{E_2\{\tilde{\Omega}B\}}^{\tilde{\Omega}\varphi}$ of E_2 -algebra structures deforming the E_2 -algebra structure $\tilde{\Omega}\varphi:E_2\to End_{\tilde{\Omega}B}$ on its cobar construction $\tilde{\Omega}B$. For this, we use the characterization of such moduli problems obtained in [82]. We deduce that the Gerstenhaber-Schack complex of B is quasi-isomorphic to the higher Hochschild complex of ΩB as L_{∞} -algebras, for the L_{∞} structure on this Hochschild complex induced by its E_3 structure. The existence of this E_3 structure on the higher Hochschild complex is a higher version of Deligne conjecture proved in [20] and [36]. Precisely, we prove the following results:

Theorem 0.6. Let B be a pointed conilpotent homotopy associative dg bialgebra. Let $\varphi: Bialg_{\infty} \to End_B$ be this homotopy bialgebra structure on B, and let $\tilde{\Omega}\varphi: E_2 \to End_{\tilde{\Omega}B}$ be the corresponding E_2 -algebra structure on its cobar construction $\tilde{\Omega}B$.

(1) There is a homotopy equivalence of formal moduli problems

$$Bialg_{\infty}\{B\}^{\varphi} \simeq E_2\{\tilde{\Omega}B\}^{\tilde{\Omega}\varphi}.$$

This homotopy equivalence induces a quasi-isomorphism of L_{∞} -algebras

$$g^{\varphi}_{Bialg,B} \stackrel{\sim}{\to} g^{\tilde{\Omega}\varphi}_{E_2,\tilde{\Omega}B}$$

between the deformation complex of B and the deformation complex of $\tilde{\Omega}B$.

(2) There is a homotopy equivalence of formal moduli problems

$$\underline{Bialg_{\infty}^{+}}\{B\}^{\varphi^{+}} \simeq \underline{E_{2}^{+}}\{\tilde{\Omega}B\}^{\tilde{\Omega}\varphi^{+}}.$$

This homotopy equivalence induces a quasi-isomorphism of L_{∞} -algebras

$$C^*_{GS}(B,B) \stackrel{\sim}{\to} T_{\tilde{\Omega}(B)}$$

between the Gerstenhaber-Schack complex of B and the (truncated) E_2 -Hochschild complex¹ of $\tilde{\Omega}B$.

Then we extend these equivalences to E_3 -algebra equivalences, and even extend this to the full Gerstenhaber-Schack complex $C_{GS}^{full}(B,B)$. We get eventually the following solution to a longstanding problem:

¹that is the E_2 -cotangent complex $T_{\tilde{\Omega}(B)}$ which is equivalent to $\mathbb{R}Der_{E_2}(\tilde{\Omega}B, \tilde{\Omega}B)$

Corollary 0.7 (Gerstenhaber-Schack conjecture). (1) There is an E_3 -algebra structures on $C_{GS}^*(B,B)$ and a unital E_3 -algebra structure on $C_{GS}^{full}(B,B)$) such that the following diagram

$$\tilde{\Omega}B[-1] \xrightarrow{} T_{\tilde{\Omega}(B)} \xrightarrow{} CH_{E_2}^*(\tilde{\Omega}B, \tilde{\Omega}B)$$

$$\stackrel{\simeq}{\parallel} \qquad \stackrel{\simeq}{\sim} \qquad \stackrel{\simeq}{\parallel} \simeq$$

$$\tilde{\Omega}B[-1] \xrightarrow{} C_{GS}^*(B, B) \xrightarrow{} C_{GS}^{full}(B, B)$$

is a commutative diagram of non-unital E_3 -algebras with vertical arrows being equivalences.

(2) The E_3 -algebra structure on $C_{GS}^*(B,B)$ is a refinement of its L_{∞} -algebra structure controlling the deformation theory of the bialgebra B.

This corollary provides a stronger version of a longstanding conjecture stated by Gerstenhaber-Schack in [29, Section 8] (the existence of a homotopy graded Lie algebra structure on the Gerstenhaber-Schack complex) for the three different versions of the Gerstenhaber-Schack and E_2 -Hochschild complexes. This also solves a stronger version of a conjecture proposed by Kontsevich in [48] in 2003 (existence of an E_3 structure in the case where B = Sym(V) is the symmetric bialgebra on a vector space V).

0.4. Kontsevich formality conjecture and deformation quantization of Lie bialgebras. In the introduction of his work on deformation quantization of Poisson manifolds [48], Kontsevich conjectured that Etingof-Kazhdan quantization should be the "degree zero part" of a much more general formality theorem of the deformation complex Def(Sym(V)) of the symmetric bialgebra Sym(V) on a vector space V. This deformation complex should possess an E_3 -algebra structure whose underlying L_{∞} -structure controls the deformations of this bialgebra, and should be formal as an E_3 -algebra.

The existence of such an E_3 -algebra structure on Def(Sym(V)) controling the deformations of Sym(V) is a particular case of Corollary 0.7. To prove the E_3 formality of Def(Sym(V)), one has to compare two E_3 -algebra structures on the Gerstenhaber-Schack cohomology of Sym(V), the one coming from the action of the operad of 3-Poisson algebras $P_3 \cong H^*E_3$ via the formality $E_3 \xrightarrow{\sim} H^*E_3 \cong P_3$ of E_3 operads (see [50] over $\mathbb R$ and more generally [28] over $\mathbb Q$), and the one transferred from Def(Sym(V)). The following formality result was conjectured by Kontsevich (in the case where V is a vector space) in the introduction of [48]:

Theorem 0.8. The deformation complex of the symmetric bialgebra Sym(V) on a \mathbb{Z} -graded cochain complex V whose cohomology is of finite dimension in each degree is formal over \mathbb{Q} as an E_3 -algebra.

We obtain it by using the relationship with E_2 -Hochschild cohomology (Corollary 0.7) and the higher HKR-theorem for the latter [7].

We then (see § 8) obtain a new proof of Etingof-Kazhdan quantization theorem from the underlying L_{∞} formality given by Theorem 0.8, the properties of Maurer-Cartan spaces for filtered L_{∞} -algebras [81] and by using the results of [59]. In fact, after the appropriate completion, Def(Sym(V)) controls the (homotopy) $\mathbb{K}[[\hbar]]$ -bialgebra structures quantizing the (homotopy) Lie bialgebra structures on

V, which are controlled by $H_{GS}^*(Sym(V), Sym(V))$ which by formality also controls deformation of dg-bialgebras. In particular, our work shows that quantization of bialgebras follows from a higher analogue of the Kontsevich approach to quantization of Poisson manifolds in [48, 72] as was conjectured from the early 2000's.

Let us add two important remarks. First, what we get is actually a generalization of Etingof-Kazhdan quantization to homotopy dg Lie bialgebras, which encompasses the case of usual Lie bialgebras. Second, since the E_3 formality does not rely on the choice of a Drinfeld associator (contrary to the E_2 formality), we thus solve another open problem by showing that such a quantization does not rely on the choice of an associator. Precisely, we get the following results:

Corollary 0.9. (1) The L_{∞} -formality underlying Theorem 0.8 induces a generalization of Etingof-Kazdhan deformation quantization theorem to homotopy dg Lie bialgebras whose cohomology is of finite dimension in each degree. In the case where V is a vector space, this gives a new proof of Etingof-Kazdhan's theorem.

(2) Deformation quantization of homotopy dg Lie bialgebras whose cohomology is of finite dimension in each degree does not rely on the choice of a Drinfeld associator.

In the case of a Lie algebra equipped with a Casimir element, the invariance under the choice of an associator was already known by Drinfeld in [9]. A similar result was announced in [12] for finite dimensional Lie bialgebras but without complete proof. Our corollary provides an alternative complete proof of this statement, as well as an extension of this invariance result to homotopy dg Lie bialgebras with finite dimensional cohomology which is completely new.

0.5. **Perspectives.** The new methods developed in this paper to approach deformation theory and quantization problems have several possible continuations. A first perspective is to get a better understanding of the moduli problem associated to the full E_n -Hochschild complex, which should control the deformation theory of E_n -monoidal dg categories and is deeply related to the deformation quantization of shifted Poisson structures in derived algebraic geometry (see for instance [77, Theorem 5.2]). In particular, in the case of the full E_2 -Hochschild complex, the equivalence with the full Gerstenhaber-Schack complex $C_{GS}^{full}(B,B)$ proved in this paper (Corollary 0.7) will imply that $C_{GS}^{full}(B,B)$ controls the deformations of Mod_B as an E_2 -monoidal dg category, that is, a braided monoidal dg category. This is still a conjecture at present. The E_3 -formality of $C_{GS}^{full}(Sym(g), Sym(g))$, where g is a Lie bialgebra, then corresponds to the deformation quantization of Mod_g with the equivalence relation on quantizations given by Drinfeld's gauge equivalence [9] (equivalence of braided monoidal categories). In the case where g = Lie(G) is the tangent Lie algebra of a reductive algebraic group G, this result has to be compared with the quantization of the derived classifying stack BG in the sense of [8].

A second perspective is to recover the major results in deformation quantization obtained in [14] and [15] as consequences of suitable variants of our Formality Theorem 0.8. In particular, we expect the quantization of coboundary Lie bialgebras of [15] to be a consequence of this formality suitably extended to algebras over framed little disks operads (see for instance [66] for a definition of framed little disks, and [32],[67] for formality results). The coboundary on the bialgebra side should correspond to the framing on the little disks side. Let us note that this quantization would be then automatically independent from the choice of an associator.

Both perspectives will be investigated by the authors in future works.

NOTATIONS

The reader will find below a list of the main notations used at several places in this article.

- ullet We work over a field of characteristic zero noted \mathbb{K} .
- $Ch_{\mathbb{K}}$ is the category of \mathbb{Z} -graded chain complexes over \mathbb{K} .
- Let (C, W_C) be a relative category, also called a category with weak equivalences. Here C is a category and W_C its subcategory of weak equivalences. The hammock localization (see [11]) of such a category with respect to its weak equivalences is noted $L^H(C, W_C)$, and the mapping spaces of this simplicial localization are noted $L^H(C, W_C)(X, Y)$.
- Several categories of algebras and coalgebras will have a dedicated notation: cdga for the category of commutative differential graded algebras, dgArt for the category of artinian cdgas, dgCog for the category of dg coassociative coalgebras and dgLie for the category of dg Lie algebras.
- Given a cdga A, the category of A-modules is noted Mod_A . More generally, if \mathcal{C} is a symmetric monoidal category tensored over $Ch_{\mathbb{K}}$, the category of A-modules in \mathcal{C} is noted $Mod_A(\mathcal{C})$.
- Given a dg Lie algebra g, its Chevalley-Eilenberg algebra is noted $C_{CE}^*(g)$ and its Chevalley-Eilenberg coalgebra is noted $C_*^{CE}(g)$.
- More general categories of algebras and coalgebras over operads or properads will have the following generic notations: given a properad P, we will note P-Alg the category of dg P-algebras and given an operad P we will note P-Cog the category of dg P-coalgebras.
- Given a properad P, a cofibrant resolution of P is noted P_{∞} .
- When the base category is a symmetric monoidal category \mathcal{C} other than $Ch_{\mathbb{K}}$, we note $P-Alg(\mathcal{C})$ the category of P-algebras in \mathcal{C} and $P-Cog(\mathcal{C})$ the category of P-coalgebras in \mathcal{C} .
- These various categories of algebras and coalgebras will be restricted to several subcategories for which we will add the following super scripts: aug for augmented and coaug for coaugmented (for example, $P Alg^{aug}$ is the category of augmented P-algebras), con for connected, 0 con for 0-connected, conil for conilpotent, and pt for pointed.
- Algebras over properads form a relative category for the weak equivalences defined by chain quasi-isomorphisms. The subcategory of weak equivalences of P-Alg is noted wP-Alg.
- Given a properad P and a complex X, we will consider an associated convolution Lie algebra noted $g_{P,X}$ which will give rise to two deformation complexes: the deformation complex $g_{P,X}^{\varphi}$ controling the formal moduli problem of deformations of a P-algebra structure φ on X, and a variant $g_{P+X}^{\varphi^+}$ whose role will be explained in Section 3.
- We will consider various moduli functors in this paper, defined as simplicial presheaves over artinian augmented cdgas: the simplicial presheaf of P_{∞} -algebra structures on X noted $\underline{P_{\infty}\{X\}}$, the formal moduli problem of deformations of a given P_{∞} -algebra structure φ on X noted $\underline{P_{\infty}\{X\}}^{\varphi}$, and the derived algebraic group of homotopy automorphisms of (X, φ) noted

 $\underline{haut}_{P_{\infty}}(X,\varphi)$. The derived algebraic group of automorphisms of X as a chain complex will be noted $\underline{haut}(X)$.

- The little n-disks operads have several variants we will distinguish as follows: E_n for the operad of non-unitary E_n -algebras, uE_n for the operad of E_n -algebras with a unit up to homotopy, and suE_n for the operad of strictly unitary E_n -algebras. The same notations hold for the operad $Pois_n$ of n-Poisson algebras.
- We will use two tensor products of operads. The tensor product arity by arity, also called the Hadamar tensor product, will be noted $P \otimes_H Q$. The Boardman-Vogt tensor product will be noted \otimes (see [18] for a definition).
- The standard (full) complex computing the (André-Quillen) cohomology theory of n-Poisson algebra A is denoted $CH_{Pois_n}^*(A,A)$. Tamarkin standard deformation complex is the subcomplex in weight greater than 0, that we denote $CH_{Pois_n}^{(\bullet>0)}(A,A)$. There is also another truncation $CH_{Pois_n}^{(\bullet>1)}(A,A)$ complex has two gradings, the cohomological grading from the one of A and a weight grading for the subcomplex in weight greater than 1. The same notations will be used for some standard complexes computing E_n -Hochschild cohomology of E_n -algebras.
- The tangent complex of an E_n -algebra A in the sense of [20] is noted T_A .
- The centralizer of an E_n -algebra morphism f in the sense of [54] is noted $\mathfrak{z}_{E_n}(f)$, the version for n-Poisson algebras is noted $\mathfrak{z}_{Pois_n}(f)$.
- Several kinds of bar and cobar constructions will play an important role in this paper. The functors Bar and Ω are the classical bar and cobar constructions respectively for augmented associative algebras and coaugmented coassociative coalgebras. The operadic Koszul duality functors $Bar^{(n)}$ and $Cobar^{(n)}$ are adjoint functors defining respectively the bar construction for augmented E_n -algebras and the cobar construction for conilpotent E_n -coalgebras, which can be modeled respectively by an n-iterated bar construction and an n-iterated cobar construction (hence the notations). The versions for n-Poisson algebras are noted $Bar_{Pois_n}^{(n)}$ and $Cobar_{Pois_n}^{(n)}$. Another Koszul duality functor is the one used in [52], which is an endofunctor of the category of augmented E_n -algebras noted \mathcal{D}^n .
- More generally, given a Koszul operad P in a symmetric monoidal ∞ category \mathcal{C} , there is a bar construction $\mathcal{B}_P: P-Alg(\mathcal{C}) \to \mathcal{C}$, which can
 be enhanced by the Barr-Beck-Lurie theorem [54] to a bar functor $\mathcal{B}_P^{enh}:$ $P-Alg(\mathcal{C}) \to P^! Cog^{conil}(\mathcal{C})$ where $P^!$ is the Koszul dual operad of P.
 Similarly, there is a cobar construction Ω_P and an enhanced version Ω_P^{enh} .

1. Recollections

The goal of this section is to briefly review the key notions and results at the heart of the present paper.

1.1. Symmetric monoidal categories over a base category.

Definition 1.1. Let \mathcal{C} be a symmetric monoidal category. A symmetric monoidal category over \mathcal{C} is a symmetric monoidal category $(\mathcal{E}, \otimes_{\mathcal{E}}, 1_{\mathcal{E}})$ endowed with a symmetric monoidal functor $\eta: \mathcal{C} \to \mathcal{E}$, that is, an object under \mathcal{C} in the 2-category of symmetric monoidal categories.

This defines on \mathcal{E} an external tensor product $\otimes : \mathcal{C} \times \mathcal{E} \to \mathcal{E}$ by $C \otimes X = \eta(C) \otimes_{\mathcal{E}} X$ for every $C \in \mathcal{C}$ and $X \in \mathcal{E}$. This external tensor product is equipped with the following natural unit, associativity and symmetry isomorphisms:

- $(1) \ \forall X \in \mathcal{E}, 1_{\mathcal{C}} \otimes X \cong X,$
- $(2) \ \forall X \in \mathcal{E}, \forall C, D \in \mathcal{C}, (C \otimes D) \otimes X \cong C \otimes (D \otimes X),$
- $(3) \ \forall C \in \mathcal{C}, \forall X, Y \in \mathcal{E}, C \otimes (X \otimes Y) \cong (C \otimes X) \otimes Y \cong X \otimes (C \otimes Y).$

The coherence constraints of these natural isomorphisms (associativity pentagons, symmetry hexagons and unit triangles which mix both internal and external tensor products) come from the symmetric monoidal structure of the functor η .

We will implicitly assume throughout the paper that all small limits and small colimits exist in \mathcal{C} and \mathcal{E} , and that each of these categories admit an internal hom bifunctor. We suppose moreover the existence of an external hom bifunctor $Hom_{\mathcal{E}}(-,-): \mathcal{E}^{op} \times \mathcal{E} \to \mathcal{C}$ satisfying an adjunction relation

$$\forall C \in \mathcal{C}, \forall X, Y \in \mathcal{E}, Mor_{\mathcal{E}}(C \otimes X, Y) \cong Mor_{\mathcal{C}}(C, Hom_{\mathcal{E}}(X, Y))$$

(so \mathcal{E} is naturally an enriched category over \mathcal{C}).

Example 1.2. Let I be a small category; the I-diagrams in a symmetric monoidal category \mathcal{C} form a symmetric monoidal category over \mathcal{C} . The internal tensor product is defined pointwise, and the external tensor product is defined by the functor η which associates to $X \in \mathcal{C}$ the constant I-diagram C_X on X. The external hom $Hom_{\mathcal{C}^I}(-,-): \mathcal{C}^I \times \mathcal{C}^I \to \mathcal{C}$ is given by

$$Hom_{\mathcal{C}^I}(X,Y) = \int_{i \in I} Hom_{\mathcal{C}}(X(i),Y(i)).$$

Throughout this paper we will deal with symmetric monoidal categories equipped with a model structure. We assume that the reader is familiar with the basics of model categories. We refer to to Hirschhorn [42] and Hovey [41] for a comprehensive treatment of homotopical algebra. We just recall the axioms of symmetric monoidal model categories formalizing the interplay between the tensor and the model structures.

Definition 1.3. (1) A symmetric monoidal model category is a symmetric monoidal category C equipped with a model category structure such that the following axioms holds:

MM0. For any cofibrant object X of \mathcal{C} , the map $Q1_{\mathcal{C}} \otimes X \to 1_{\mathcal{C}} \otimes X \cong X$ induced by a cofibrant resolution $Q1_{\mathcal{C}} \to 1_{\mathcal{C}}$ of the unit $1_{\mathcal{C}}$ is a weak equivalence.

MM1. The pushout-product $(i_*, j_*): A \otimes D \oplus_{A \otimes C} B \otimes C \to B \otimes D$ of cofibrations $i: A \mapsto B$ and $j: C \mapsto D$ is a cofibration which is also acyclic as soon as i or j is so.

(2) Suppose that \mathcal{C} is a symmetric monoidal model category. A symmetric monoidal category \mathcal{E} over \mathcal{C} is a symmetric monoidal model category over \mathcal{C} if the axiom MM1 holds for both the internal and external tensor products of \mathcal{E} .

Example 1.4. The usual projective model category $Ch_{\mathbb{K}}$ of unbounded chain complexes over a field \mathbb{K} forms a symmetric monoidal model category.

A property of the pushout-product axiom MM1 which will be useful later is that it is equivalent to the following dual version:

Lemma 1.5. (cf. [41, Lemma 4.2.2]) In a symmetric monoidal model category C, the axiom MM1 is equivalent to the following one:

MM1'. The morphism

$$(i^*, p_*): Hom_{\mathcal{C}}(B, X) \to Hom_{\mathcal{C}}(A, X) \times_{Hom_{\mathcal{C}}(A, Y)} Hom_{\mathcal{C}}(B, Y)$$

induced by a cofibration $i: A \rightarrow B$ and a fibration $p: X \rightarrow Y$ is a fibration in C which is also acyclic as soon as i or p is so.

1.2. Operads, Props and their algebras.

1.2.1. Props and their algebras. Let \mathcal{C} be a symmetric monoidal category. A Σ -biobject is a double sequence $\{M(m,n)\in\mathcal{C}\}_{(m,n)\in\mathbb{N}^2}$ where each M(m,n) is equipped with a right action of Σ_m and a left action of Σ_n commuting with each other.

Definition 1.6. A prop is a Σ -biobject endowed with associative horizontal composition products

$$\circ_h: P(m_1, n_1) \otimes P(m_2, n_2) \to P(m_1 + m_2, n_1 + n_2),$$

associative vertical composition products

$$\circ_v: P(k,n) \otimes P(m,k) \to P(m,n)$$

and units $1 \to P(n, n)$ which are neutral for \circ_v . These products satisfy the exchange law

$$(f_1 \circ_h f_2) \circ_v (g_1 \circ_h g_2) = (f_1 \circ_v g_1) \circ_h (f_2 \circ_v g_2)$$

and are compatible with the actions of symmetric groups.

Morphisms of props are equivariant morphisms of collections compatible with the composition products.

Definition 1.7. (1) To any object X of \mathcal{C} we can associate an endomorphism prop End_X defined by

$$End_X(m,n) = Hom_{\mathcal{C}}(X^{\otimes m}, X^{\otimes n}).$$

(2) A P-algebra is an object $X \in \mathcal{C}$ equipped with a prop morphism $P \to End_X$.

We can also define a P-algebra in a symmetric monoidal category over C:

Definition 1.8. Let \mathcal{E} be a symmetric monoidal category over \mathcal{C} .

- (1) The endomorphism prop of $X \in \mathcal{E}$ is given by $End_X(m,n) = Hom_{\mathcal{E}}(X^{\otimes m}, X^{\otimes n})$ where $Hom_{\mathcal{E}}(-,-)$ is the external hom bifunctor of \mathcal{E} .
- (2) Let P be a prop in \mathcal{C} . A P-algebra in \mathcal{E} is an object $X \in \mathcal{E}$ equipped with a prop morphism $P \to End_X$.

There is a functorial free prop construction F leading to an adjunction

$$F: \mathcal{C}^{\mathbb{S}} \rightleftarrows Prop: U$$

with the forgetful functor U. The category of Σ -biobjects $\mathcal{C}^{\mathbb{S}}$ is a diagram category over \mathcal{C} , so it inherits the usual projective model structure of diagrams, which is a cofibrantly generated model category structure. In the case of unbounded chain complexes over a field of characteristic zero, this model structure can be transferred along the free-forgetful adjunction:

Theorem 1.9. (cf. [23, Theorem 5.5]) The category of dg props Prop equipped with the classes of componentwise weak equivalences and componentwise fibrations forms a cofibrantly generated model category.

1.2.2. Properads. Composing operations of two Σ -biobjects M and N amounts to consider 2-levelled directed graphs (with no loops) with the first level indexed by operations of M and the second level by operations of N. Vertical composition by grafting and horizontal composition by concatenation allows one to define props as before. The idea of properads is to mimick operads (for operations with several outputs), which are defined as monoids in Σ -objects, by restricting the vertical composition product to connected graphs. The unit for this connected composition product \boxtimes_c is the Σ -biobject I given by $I(1,1) = \mathbb{K}$ and I(m,n) = 0 otherwise. The category of Σ -biobjects then forms a symmetric monoidal category $(Ch_{\mathbb{K}}^{\mathbb{S}}, \boxtimes_c, I)$.

Definition 1.10. A dg properad (P, μ, η) is a monoid in $(Ch_{\mathbb{K}}^{\mathbb{S}}, \boxtimes_{c}, I)$, where μ denotes the product and η the unit. It is augmented if there exists a morphism of properads $\epsilon: P \to I$. In this case, there is a canonical isomorphism $P \cong I \oplus \overline{P}$ where $\overline{P} = ker(\epsilon)$ is called the augmentation ideal of P.

Morphisms of properads are morphisms of monoids in $(Ch_{\mathbb{K}}^{\mathbb{S}}, \boxtimes_{c}, I)$.

Properads have also their dual notion, namely coproperads:

Definition 1.11. A dg coproperad (C, Δ, ϵ) is a comonoid in $(Ch_{\mathbb{K}}^{\mathbb{S}}, \boxtimes_{c}, I)$.

As in the prop case, there exists a free properad functor $\mathcal F$ forming an adjunction

$$\mathcal{F}: Ch^{\mathbb{S}}_{\mathbb{K}} \rightleftarrows Properad: U$$

with the forgetful functor U. There is an explicit construction of the free properad in terms of direct sums of labelled graphs for which we refer the reader to [79]. Dually, there exists a cofree coproperad functor denoted $\mathcal{F}_c(-)$ having the same underlying Σ -biobject. Moreover, according to [57], this adjunction equips dg properads with a cofibrantly generated model category structure with componentwise fibrations and weak equivalences.

There is also a notion of algebra over a properad similar to an algebra over a prop, since the endomorphism prop restricts to an endomorphism properad. Properads are general enough to encode a wide range of bialgebra structures such as associative and coassociative bialgebras, Lie bialgebras, Poisson bialgebras, Frobenius bialgebras for instance.

1.2.3. Algebras over operads. Operads are used to parametrize various kind of algebraic structures consisting of operations with one single output. Fundamental examples of operads include the operad As encoding associative algebras, the operad Com of commutative algebras, the operad Lie of Lie algebras and the operad Pois of Poisson algebras. They can be defined as monoids in the category of Σ -objects (collections $\{P(n)\}_{n\in\mathbb{N}}$ with an action of Σ_n on each P(n)), for the appropriate composition product. Dg operads form a model category with bar-cobar resolutions and Koszul duality [51]. An algebra X over a dg operad P can be defined in any symmetric monoidal category \mathcal{E} over $Ch_{\mathbb{K}}$, alternatively as an algebra over the corresponding monad or as an operad morphism $P \to End_X$ where $End_X(n) = Hom_{\mathcal{E}}(X^{\otimes n}, X)$ and $Hom_{\mathcal{E}}$ is the external hom bifunctor. The category of P-algebras satisfies good homotopical properties, namely:

Theorem 1.12. (see [22, Theorem 12.3.A]) Let P be a Σ -cofibrant dg operad and \mathcal{E} be a cofibrantly generated symmetric monoidal model category over $Ch_{\mathbb{K}}$. Then the category \mathcal{E}^P of P-algebras in \mathcal{E} inherits a cofibrantly generated semi-model category structure such that the forgetful functor $U: \mathcal{E}^P \to \mathcal{E}$ creates weak equivalences and

fibrations, and that the generating (acyclic) cofibrations are the images under the free P-algebra functor of the generating (acyclic) cofibrations of \mathcal{E} .

Remark 1.13. In general, algebras over properads and props do not inherit such a model category structure, since there is no free algebra functor in this case.

Definition 1.14. (1) Let P be an operad. A P-coalgebra is a complex C equiped with linear applications $\rho_n: P(n) \otimes C \to C^{\otimes n}$ for every $n \geq 0$. These maps are Σ_n -equivariant and associative with respect to the operadic compositions.

(2) Each $p \in P(n)$ gives rise to a cooperation $p^*: C \to C^{\otimes n}$. The coalgebra C is usually said to be conilpotent if for each $c \in C$, there exists $N \in \mathbb{N}$ so that $p^*(c) = 0$ when we have $p \in P(n)$ with n > N.

If \mathbb{K} is a field of characteristic zero and the P(n) are finite dimensional, then it is equivalent to define a P-coalgebra via a family of applications $\overline{\rho}_n: C \to P(n)^* \otimes_{\Sigma_n} C^{\otimes n}$.

Theorem 1.15. The category $dgCog^{conil}$ of non-negatively graded conilpotent coassociative dg coalgebras forms a cofibrantly generated model category with quasi-isomorphisms as weak equivalences and degreewise injections as cofibrations.

This follows verbatim from the argument line of [80, Theorem 0.1] adapted to the non-negatively graded setting, or the argument line of [31] adapted to the conilpotent case. In both cases, the model structure is transferred via the adjunction between the forgetful functor and the reduced tensor coalgebra functor (which is both the cofree coalgebra functor in the positively graded setting of [80] and the cofree conilpotent coalgebra functor in the non-negatively graded setting of [31]). The fact that this model category is cofibrantly generated is crucial in the proof of Theorem 0.1(2).

We will also need a more general model category structure on conilpotent coalgebras over dg operads, however without needing it to be cofibrantly generated. The existence of such a model category results from the following arguments. There are very general assumptions under which a category of coalgebras over a comonad in a model category inherits a model category structure [37], which can be applied to the case of coalgebras over differential non-negatively graded operads. Given a differential non-negatively graded operad P, to check that the category $Ch_{\mathbb{K}}$ has the needed properties to get a left-induced model structure on the category $P - Cog^{conil}$ of dg P-coalgebras from [37, Theorem 5.8], one just specializes the results of [37, Section 6] (enunciated for right dg modules over a dga A) to the case $A = \mathbb{K}$:

Theorem 1.16. Let P be a differential non-negatively graded operad. The category $P - Cog^{conil}$ inherits a model category structure such that the forgetful functor $U: P - Cog^{conil} \to Ch_{\mathbb{K}}$ creates the cofibrations and the weak equivalences.

1.3. E_n -operads, higher Hochschild cohomology and the Deligne conjecture. Configuration spaces of n-disks into a bigger n-disk gather into a topological operad D_n called the little n-disks operad. An E_n -operad is a dg operad quasiisomorphic to the singular chains C_*D_n of the little n-disks operad. The formality of the little 2-disks operad (i.e. the fact that it is quasi-isomorphic as an operad to its cohomology) is the key point to prove Deligne conjecture about the homotopy Gerstenhaber structure (that is, the E_2 -algebra structure) of the Hochschild complex, which in turn provided an alternative method for deformation quantization of Poisson manifolds [47], [48], [72], [73]. Its formality holds over $\mathbb Q$ and relies on the choice of a Drinfeld associator (which exists over \mathbb{Q} , as proved in [9]), a process closely related to deformation quantization of Lie bialgebras as well. The formality of E_n -operads for $n \geq 3$ was sketched over \mathbb{R} in [47] and fully proved in [50]. This result was recently superseded in [28], where this formality was proved to hold over Q and to be intrinsic (under a mild technical assumption satisfied in particular by the little n-disks operads). An important observation to note here is that contrary to the case n=2, the formality of E_n -operads for $n\geq 3$ does not rely on the choice of an associator.

Given an ordinary associative (or E_1) algebra A, its endomorphisms $Hom_{biMod_A}(A, A)$ in the category $biMod_A$ of A-bimodules form nothing but the center Z(A) of A. Deriving this hom object gives the Hochschild complex $CH^*(A, A)$ of A, and the Hochschild cohomology $HH^*(A, A)$ of A satisfies $HH^0(A, A) = Z(A)$. One says that the Hoschchild complex is the derived center of A, and the Deligne conjecture can then be reformulated as "the derived center of an E_1 -algebra forms an E_2 -algebra". This sentence enlarges to a similar statement for E_n -algebras.

Definition 1.17. The (full) Hochschild complex of an E_n -algebra A, computing its higher Hochschild cohomology, is the derived hom $CH_{E_n}^*(A,A) = \mathbb{R}Hom_A^{E_n}(A,A)$ in the category of (operadic) A-modules over E_n .

The Deligne conjecture endows the Hochschild cochain complex with an E_{n+1} -algebra structure [36, Theorem 6.28] or [20, 54]. Associated to an E_n -algebra A, one also has its cotangent complex L_A , which classifies square-zero extensions of A [20, 54].

Definition 1.18 ([20]). The tangent complex T_A of an E_n -algebra A is the dual $T_A := Hom_A^{E_n}(L_A, A) \cong \mathbb{R}Der(A, A)$.

Francis [20, 54] has proved that $T_A[-n]$ has a canonical structure of E_{n+1} -algebra and further that we have a fiber sequence $T_A[-n] \to CH_{E_n}^*(A,A) \to A$ where the first map is a map of E_{n+1} -algebras.

1.4. **Deformation quantization of Lie bialgebras.** Lie bialgebras originally arose from Poisson-Lie groups in mathematical physics. Poisson-Lie groups are Lie groups with a compatible Poisson structure, and appear as gauge groups of various classical mechanical systems such as integrable systems. The tangent space T_eG of a Poisson-Lie group G has more structure than a Lie algebra, because of the Poisson bracket, which induces a compatible Lie coalgebra structure on T_eG so that T_eG forms Lie bialgebra. The compatibility relation between the bracket and

the cobracket is called the Drinfeld's compatibility relation or the cocycle relation. Quantizing tangent spaces of Poisson-Lie groups gives solutions to the quantum Yang-Baxter equation, which allow to build "exactly solvable" quantum systems from classical mechanical systems. Deformation quantization of Lie bialgebras produces quantum groups, which are relevant not only for mathematical physics but also for low-dimensional topology (quantum invariants of knots and 3-manifolds), and are deeply related to algebraic geometry and number theory (moduli spaces of curves, Grothendieck-Teichmüller groups, multizeta values) via the Drinfeld associators [9].

The problem of a universal quantization of Lie bialgebras raised by Drinfeld was solved by Etingof and Kazhdan [16], [17]. A deformation quantization of a Lie bialgebra g is a topologically free Hopf algebra H over the ring of formal power series $\mathbb{K}[[\hbar]]$ such that $H/\hbar H$ is isomorphic to U(q) (the enveloping algebra of g) as a co-Poisson bialgebra. Such a Hopf algebra is called a quantum universal enveloping algebra (QUE for short). Conversely, the quasi-classical limit of a QUE algebra H is the Lie bialgebra of primitive elements $g = Prim(H/\hbar H)$ whose cobracket is induced by the coproduct of H. For technical reasons (the need of a "quasi-triangular structure" on the Lie bialgebra, that is, of a classical r-matrix), the initial construction of Etingof-Kazdhan takes as input a quasi-triangular Lie bialgebra D(g) called the double of g. Then they equip the category $Mod_{D(g)}[[\hbar]]$ (same objects than $Mod_{D(q)}$ but the vector spaces of morphisms are tensored by the ring of formal series), called the Drinfeld category, with a braided monoidal structure induced by the r-matrix of D(q) and the choice of a Drinfeld associator. The forgetful functor from the Drinfeld category to the category of modules is a braided monoidal functor, so by the tannakian formalism the Drinfeld category is equivalent to the category of modules over a topologically free Hopf algebra quantizing D(g). The QUE algebra of g is then a certain Hopf subalgebra of this Hopf algebra.

As explained in the sequel [17] with the formalism of cyclic categories, and later in [13] in the formalism of props, deformation quantization of Lie bialgebras can be reformulated as the existence of a prop morphism

$$QUE \to UE_{cP}[[\hbar]]$$

between the prop of QUE algebras and the prop $UE_{cP}[[\hbar]]$ of topologically free co-Poisson $\mathbb{K}[[\hbar]]$ -bialgebras. This prop morphism induces a quantization functor from the category of $UE_{cP}[[\hbar]]$ -algebras to the category of QUE-algebras. The category of $UE_{cP}[[\hbar]]$ -algebras is equivalent to the category of topologically free Lie bialgebras over $\mathbb{K}[[\hbar]]$ (the enveloping algebra of a Lie bialgebra is a co-Poisson bialgebra). The existence of such a prop morphism relies on the use of a Drinfeld associator. An alternative propic approach to deformation quantization, which will be useful for us in this paper, is presented in [59]. We will go back to this with more details in Section 8.

2. Moduli problems of algebraic structures and deformation complexes

2.1. Formal moduli problems and (homotopy) Lie algebras. Formal moduli problems arise when one wants to study the deformation theory of a point of a given moduli space (variety, scheme, stack, derived stack). The general principle

of moduli problems is that the deformation theory of a given point in a formal neighbourhood of this point (that is, the formal completion of the moduli space at this point) is controlled by a certain tangent dg Lie algebra. This idea of a correspondence between formal moduli problems and dg Lie algebras arose from unpublished work of Deligne, Drinfed and Feigin, and was developed further by Goldman-Millson, Hinich, Kontsevich, Manetti among others. However, there was no systematic recipe to build a dg Lie algebra for a given moduli problem, and even worse, different dg Lie algebras could represent the same moduli problem. To overcome these difficulties, one has to consider moduli problems in a derived setting. The rigorous statement of an equivalence between derived formal moduli problems and dg Lie algebras was proved independently by Lurie in [52] and by Pridham in [63].In this paper, what we will call moduli problems are actually derived moduli problems.

Briefly, formal moduli problems are functors $F: dgArt^{aug}_{\mathbb{K}} \to sSet$ from augmented artinian commutative differential graded algebras to simplicial sets, such that $F(\mathbb{K}) \simeq pt$ and F preserves certain pullbacks (we refer the reader to [52] for more details). The value $F(\mathbb{K})$ corresponds to the point of which we study the formal neighbourhood, the evaluation $F(\mathbb{K}[t]/(t^2))$ on the algebra of dual numbers encodes infinitesimal deformations of this point, and the $F(\mathbb{K}[t]/(t^n))$ are polynomial deformations of a higher order, for instance. Formal moduli problems form a full sub- ∞ -category noted $FMP_{\mathbb{K}}$ of the ∞ -category of simplicial presheaves over augmented artinian cdgas. By [52, Theorem 2.0.2], this ∞ -category is equivalent to the ∞ -category $dgLie_{\mathbb{K}}$ of dg Lie algebras. Moreover, one side of the equivalence is made explicit, and is equivalent to the nerve construction of dg Lie algebras studied thoroughly by Hinich in [38]. The homotopy invariance of the nerve relies on nilpotence conditions on the dg Lie algebra. In the case of formal moduli problems, this nilpotence condition is always satisfied because one tensors the Lie algebra with the maximal ideal of an augmented artinian cdga.

It turns out that this nerve construction can be extended to homotopy Lie algebras, that is, L_{∞} -algebras. There are two equivalent definitions of an L_{∞} -algebra:

Definition 2.1. (1) An L_{∞} -algebra is a graded vector space $g = \{g_n\}_{n \in \mathbb{Z}}$ equipped with maps $l_k : g^{\otimes k} \to g$ of degree 2-k, for $k \geq 1$, satisfying the following properties:

- $l_k(..., x_i, x_{i+1}, ...) = -(-1)^{|x_i||x_{i+1}|} l_k(..., x_{i+1}, x_i, ...)$
- for every $k \ge 1$, the generalized Jacobi identities

$$\sum_{i=1}^{k} \sum_{\sigma \in Sh(i,k-i)} (-1)^{\epsilon(i)} l_k(l_i(x_{\sigma(1)},...,x_{\sigma(i)}), x_{\sigma(i+1)},...,x_{\sigma(k)}) = 0$$

where σ ranges over the (i, k-i)-shuffles and

$$\epsilon(i) = i + \sum_{j_1 < j_2, \sigma(j_1) > \sigma(j_2)} (|x_{j_1}||x_{j_2}| + 1).$$

(2) An L_{∞} -algebra structure on a graded vector space $g = \{g_n\}_{n \in \mathbb{Z}}$ is a coderivation $Q: Sym^{\bullet \geq 1}(g[1]) \to Sym^{\bullet \geq 1}(g[1])$ of degree 1 of the cofree cocommutative coalgebra $Sym^{\bullet \geq 1}(g[1])$ such that $Q^2 = 0$.

The bracket l_1 is actually the differential of g as a cochain complex. When the brackets l_k vanish for $k \geq 3$, then one gets a dg Lie algebra. The dg algebra $C^*(g)$

obtained by dualizing the dg coalgebra of (2) is called the Chevalley-Eilenberg algebra of g.

A L_{∞} algebra g is filtered if it admits a decreasing filtration

$$g = F_1 g \supseteq F_2 g \supseteq \dots \supseteq F_r g \supseteq \dots$$

compatible with the brackets: for every $k \geq 1$,

$$l_k(F_rg, g, ..., g) \in F_rg.$$

We suppose moreover that for every r, there exists an integer N(r) such that $l_k(g,...,g) \subseteq F_r g$ for every k > N(r). A filtered L_{∞} algebra g is complete if the canonical map $g \to \lim_r g/F_r g$ is an isomorphism.

The completeness of a L_{∞} algebra allows to define properly the notion of Maurer-Cartan element:

Definition 2.2. (1) Let g be a dg L_{∞} -algebra and $\tau \in g^1$, we say that τ is a Maurer-Cartan element of g if

$$\sum_{k \ge 1} \frac{1}{k!} l_k(\tau, ..., \tau) = 0.$$

The set of Maurer-Cartan elements of g is noted MC(g).

(2) The simplicial Maurer-Cartan set is then defined by

$$MC_{\bullet}(g) = MC(g \hat{\otimes} \Omega_{\bullet}),$$

, where Ω_{\bullet} is the Sullivan cdga of de Rham polynomial forms on the standard simplex Δ^{\bullet} (see [70]) and $\hat{\otimes}$ is the completed tensor product with respect to the filtration induced by g.

The simplicial Maurer-Cartan set is a Kan complex, functorial in g and preserves quasi-isomorphisms of complete L_{∞} -algebras. The Maurer-Cartan moduli set of g is $\mathcal{MC}(g) = \pi_0 MC_{\bullet}(g)$: it is the quotient of the set of Maurer-Cartan elements of g by the homotopy relation defined by the 1-simplices. When g is a complete dg Lie algebra, it turns out that this homotopy relation is equivalent to the action of the gauge group $\exp(g^0)$ (a prounipotent algebraic group acting on Maurer-Cartan elements), so in this case this moduli set coincides with the one usually known for Lie algebras. We refer the reader to [82] for more details about all these results. We also recall briefly the notion of twisting by a Maurer-Cartan element. The twisting of a complete L_{∞} algebra g by a Maurer-Cartan element τ is the complete L_{∞} algebra g^{τ} with the same underlying graded vector space and new brackets l_k^{τ} defined by


$$l_k^{\tau}(x_1, ..., x_k) = \sum_{i \ge 0} \frac{1}{i!} l_{k+i}(\underbrace{\tau, ..., \tau}_{i}, x_1, ..., x_k)$$

where the l_k are the brackets of g.

To conclude, we explain why Lurie's equivalence [52, Theorem 2.0.2] lifts from the ∞ -category of dg Lie algebras dgLie to the ∞ -category of L_∞ -algebras $L_\infty - Alg$. Let $p:L_\infty \stackrel{\sim}{\to} Lie$ be the cofibrant resolution of the operad Lie encoding L_∞ -algebras. This morphism induces a functor $p^*:dgLie \to L_\infty - Alg$ which associates to any dg Lie algebra the L_∞ -algebra with the same differential, the same bracket of arity 2 and trivial higher brackets in arities greater than 2. It turns out that this functor is a right Quillen functor belonging to a Quillen equivalence

$$p_1: L_{\infty} - Alg \leftrightarrows dgLie: p^*$$

since p is a quasi-isomorphism of Σ -cofibrant operads (see [22, Theorem 16.A] for the general result). Quillen equivalences induce equivalences of the ∞ -categories associated to these model categories (which can be realized, for instance, by taking the coherent nerve of the simplicial localization of these model categories). We have a commutative triangle of ∞ -categories



where ψ and $\tilde{\psi}$ send a Lie algebra, respectively an L_{∞} -algebra, to its nerve functor or Maurer-Cartan space. The maps p^* and ψ are weak equivalences of ∞ -categories in the chosen model category of ∞ -categories (which can be for instance the one of quasicategories [53], but actually any model works). By the two-out-of-three property of weak equivalences, this implies that $\tilde{\psi}: L_{\infty} - Alg \to FMP_{\mathbb{K}}$ is a weak equivalence of ∞ -categories.

2.2. Fully faithful ∞ -functors. In the next section, we will need the following useful criteria to obtain fully faithful ∞ -functors between categories of algebras, and see under which conditions they induce equivalences between the associated formal moduli problems:

Lemma 2.3. Let $F:(\mathcal{C},W_{\mathcal{C}}) \rightleftarrows (\mathcal{D},W_{\mathcal{D}}): G$ be an adjunction of relative categories (that is, the functors F and G preserves weak equivalences) such that the unit and counit of this adjunction are pointwise weak equivalences. Then F induces an equivalence of ∞ -categories with inverse G.

Proof. Let us denote by RelCat the category of relative categories. The objects are the relative categories and the morphisms are the relative functors, that is, the functors restricting to functors between the categories of weak equivalences. By [4, Theorem 6.1], there is an adjunction between the category of bisimplicial sets and the category of relative categories

$$K_{\xi}: sSets^{\Delta^{op}} \leftrightarrows RelCat: N_{\xi}$$

(where K_{ξ} is the left adjoint and N_{ξ} the right adjoint) which lifts any Bousfield localization of the Reedy model structure of bisimplicial sets into a model structure on RelCat. In the particular case of the Bousfield localization defining the model category CSS of complete Segal spaces [65, Theorem 7.2], one obtains a Quillen equivalent homotopy theory of ∞ -categories in RelCat [4].

A way to build the ∞ -category associated to a relative category $(\mathcal{C}, W_{\mathcal{C}})$ is to take a functorial fibrant resolution $N_{\xi}(\mathcal{C}, W_{\mathcal{C}})^f$ of the bisimplicial set $N_{\xi}(\mathcal{C}, W_{\mathcal{C}})$ in CSS to get a complete Segal space. So we want to prove that $N_{\xi}F^f$ is a weak equivalence of CSS. For this, let us note first that the assumption on the adjunction between F and G implies that F is a strict homotopy equivalence in RelCat in the sense of [4]. By [4, Proposition 7.5 (iii)], the functor N_{ξ} preserves homotopy equivalences, so $N_{\xi}F$ is a homotopy equivalence of bisimplicial sets, hence a Reedy weak equivalence. Since CSS is a Bousfield localization of the Reedy model structure on bisimplicial sets, Reedy weak equivalences are weak equivalences in CSS, then by applying the fibrant resolution functor $(-)^f$ we conclude that $N_{\xi}F^f$ is a weak equivalence of complete Segal spaces.

From this we deduce:

Proposition 2.4. Let $F:(\mathcal{C},W_{\mathcal{C}})\to(\mathcal{D},W_{\mathcal{D}})$ be a functor of relative categories (that is, a functor preserving weak equivalences). If F is fully faithful, then it induces a fully faithful ∞ -functor at the level of the associated ∞ -categories.

Proof. Since F is fully faithful, it induces an equivalence of relative categories between \mathcal{C} and $F(\mathcal{C})$, hence an equivalence of the associated ∞ -categories by Lemma 2.3, which exactly means that F induces a fully faithful ∞ -functor between \mathcal{C} and \mathcal{D} . \square

Corollary 2.5. A surjection of props $\varphi: P \twoheadrightarrow Q$ induces a fully faithful ∞ -functor

$$\varphi^*: Q - Alg \hookrightarrow P - Alg,$$

where weak equivalences on both sides are defined by quasi-isomorphisms.

In the formalism of Dwyer-Kan's hammock localization, a fully faithful functor $F: \mathcal{C} \to \mathcal{D}$ is a functor satisfying the following property: for every two objects X and Y of \mathcal{C} , it induces a weak equivalence of simplicial mapping spaces

$$L^{H}(\mathcal{C}, W_{\mathcal{C}})(X, Y) \stackrel{\sim}{\to} L^{H}(\mathcal{D}, W_{\mathcal{D}})(F(X), F(Y)).$$

In particular, the associated functor Ho(F) at the level of homotopy categories is fully faithful (but not an equivalence). We would like this weak equivalence to restrict at the level of homotopy automorphisms. For this, let us recall that a functor $F: \mathcal{C} \to \mathcal{D}$ is conservative if the following property holds: let f be a morphism of \mathcal{C} , if F(f) is a weak equivalence of \mathcal{D} then f is a weak equivalence of \mathcal{C} .

Lemma 2.6. Let $F:(\mathcal{C},W_{\mathcal{C}})\to(\mathcal{D},W_{\mathcal{D}})$ be a functor of relative categories. Let us suppose that F is conservative and fully faithful. Then the restriction of F to the subcategories of weak equivalences

$$wF:W_{\mathcal{C}}\to W_{\mathcal{D}}$$

is fully faithful and induces, for every $X \in \mathcal{C}$, a weak equivalence of homotopy automorphisms

$$L^H W_{\mathcal{C}}(X, X) \stackrel{\sim}{\to} L^H W_{\mathcal{D}}(F(X), F(X)),$$

where L^HW_C is Dwyer-Kan's hammock localization of W_C with respect to itself.

Proof. Since F preserves weak equivalences, it induces a functor $wF:W_{\mathcal{C}}\to W_{\mathcal{D}}$. Moreover, the functor wF is clearly faithful as well. Now let $f:F(X)\to F(Y)$ be a weak equivalence of \mathcal{D} . Since F is full, there exists a morphism g of \mathcal{C} such that f=F(g). Since F is conservative, the fact that f is a weak equivalence of \mathcal{D} implies that g is a weak equivalence of \mathcal{C} , hence wF is full. According to Proposition 2.4, the functor wF gives a fully faithful ∞ -functor, so at the level of hammock localizations it induces a weak equivalence

$$L^H W_{\mathcal{C}}(X,Y) \stackrel{\sim}{\to} L^H W_{\mathcal{D}}(F(X),F(Y))$$

for every two objects X and Y of \mathcal{C} , in particular

$$L^H W_{\mathcal{C}}(X, X) \stackrel{\sim}{\to} L^H W_{\mathcal{D}}(F(X), F(X)).$$

Corollary 2.7. A surjection of props $\varphi: P \twoheadrightarrow Q$ induces weak equivalences of homotopy automorphisms

$$L^H wQ - Alg(X,X) \stackrel{\sim}{\to} L^H wP - Alg(\varphi^*X,\varphi^*X)$$

for every Q-algebra X.

2.3. Moduli problems of algebraic structures. Moduli spaces of algebraic structures were originally defined as simplicial sets, in the setting of simplicial operads [64]. This notion can be extended to algebras over differential graded props as follows (see [82]):

Definition 2.8. Let P_{∞} be a cofibrant prop and X be a complex. The moduli space of P_{∞} -algebra structures on X is the simplicial set $P_{\infty}\{X\}$ defined by

$$P_{\infty} = Mor_{Prop}(P_{\infty}, End_X \otimes \Omega_{\bullet}),$$

where the prop $End_X \otimes \Omega_{\bullet}$ is defined by $End_X \otimes \Omega_{\bullet})(m,n) = End_X(m,n) \otimes \Omega_{\bullet}$ and Ω_{\bullet} is the Sullivan cdga of the standard simplex Δ^{\bullet} .

This simplicial set enjoys the following properties:

Proposition 2.9. (1) The simplicial set $P_{\infty}\{X\}$ is a Kan complex and

$$\pi_0 P_{\infty} \{X\} = [P_{\infty}, End_X]_{Ho(Prop)}$$

is the set of homotopy classes of P_{∞} -algebra structures on X.

(2) Any weak equivalence of cofibrant props $P_{\infty} \xrightarrow{\sim} Q_{\infty}$ induces a weak equivalence of Kan complexes $Q_{\infty}\{X\} \xrightarrow{\sim} P_{\infty}\{X\}$.

For the remaining part of the paper, we will restrict ourselves to dg properads. Cofibrant resolutions of a properad P can always be obtained as a cobar construction $\Omega(C)$ on some coproperad C (which is usually the bar construction or the Koszul dual if P is Koszul). Given a cofibrant resolution $\Omega(C) \stackrel{\sim}{\to} P$ of P and another properad Q, one considers the convolution dg Lie algebra $Hom_{\Sigma}(\overline{C},Q)$ consisting in morphisms of Σ -biobjects from the augmentation ideal of C to Q. The Lie bracket is the antisymmetrization of the convolution product. This convolution product is defined similarly to the convolution product of morphisms from a coalgebra to an algebra, using the infinitesimal coproduct of C and the infinitesimal product of Q. The total complex $Hom_{\Sigma}(\overline{C},Q)$ is a complete dg Lie algebra. More generally, if P is a properad with minimal model $(\mathcal{F}(s^{-1}C), \partial) \stackrel{\sim}{\to} P$ for a certain homotopy coproperad C (see [56, Section 4] for the definition of homotopy coproperads), and Q is any properad, then the complex $Hom_{\Sigma}(\overline{C},Q)$ is a complete dg L_{∞} algebra. The relationship between the simplicial mapping space of morphisms $P_{\infty} \to Q$ and the Lie theory of the convolution L_{∞} -algebra $Hom_{\Sigma}(\overline{C},Q)$ is provided by the following theorem:

Theorem 2.10. (cf. [82, Theorem 2.10,Corollary 4.21]) Let P be a dg properad equipped with a minimal model $P_{\infty} := (\mathcal{F}(s^{-1}C), \partial) \stackrel{\sim}{\to} P$ and Q be a dg properad. The simplicial presheaf

$$Map(P_{\infty}, Q) : A \in dgArt^{aug}_{\mathbb{K}} \mapsto Map_{Prop}(P_{\infty}, Q \otimes A)$$

is equivalent to the simplicial presheaf

$$\underline{MC_{\bullet}}(Hom_{\Sigma}(\overline{C},Q)):A\in dgArt^{aug}_{\mathbb{K}}\mapsto MC_{\bullet}(Hom_{\Sigma}(\overline{C},Q)\otimes A)$$

associated to the complete L_{∞} -algebra $Hom_{\Sigma}(\overline{C},Q)$.

By [81, Corollary 2.4], the tensor product $MC_{\bullet}(Hom_{\Sigma}(\overline{C},Q)\otimes A)$ does not need to be completed because A is artinian. In the following, we will also consider the simplicial presheaf

$$\underline{MC^{fmp}_{\bullet}(Hom_{\Sigma}(\overline{C},Q))}: A \in dgArt^{aug}_{\mathbb{K}} \mapsto MC_{\bullet}(Hom_{\Sigma}(\overline{C},Q) \otimes m_{A}),$$

where m_A is the maximal ideal of A. This presheaf is the formal moduli problem associated to $Hom_{\Sigma}(\overline{C}, Q)$ under Lurie's equivalence theorem.

We will use such results in the case where $Q = End_X$. In this case, the simplicial presheaf $\underline{Map}(P_{\infty},Q)$ will be noted $\underline{P_{\infty}\{X\}}$. Let us note that $\underline{P_{\infty}\{X\}}$ can be alternately defined by

$$A \mapsto P_{\infty} \otimes A\{X \otimes A\}_{Mod_A},$$

where $P_{\infty} \otimes A\{X \otimes A\}_{Mod_A}$ is the mapping space of dg props in A-modules $Map(P_{\infty} \otimes A, End_{X \otimes A}^{Mod_A})$ where $End_{X \otimes A}^{Mod_A}$ is the endormorphism prop of $X \otimes A$ taken in the category of A-modules. That is, the simplicial moduli space of P_{∞} -algebra structures on $X \otimes A$ in the category of A-modules. Indeed, since Mod_A is tensored over $Ch_{\mathbb{K}}$, on can make P_{∞} act on A-modules either by morphisms of dg props in A-modules from $P_{\infty} \otimes A$ to the endomorphism prop defined by the internal hom of Mod_A , or by morphisms of dg props from P_{∞} to the endomorphism prop defined by the external hom of Mod_A .

This theorem applies in particular to the case of a Koszul properad, which includes for instance Frobenius algebras, Lie bialgebras and their variants such as involutive Lie bialgebras in string topology. It applies also to more general situations such as the properad *Bialg* encoding associative and coassociative bialgebras, which is homotopy Koszul [56, Proposition 41].

By Proposition 2.9, the simplicial set $\underline{P_{\infty}\{X\}}(A)$ classifies $P_{\infty} \otimes A$ -algebra structures on $X \otimes A$. However, the simplicial presheaf $\underline{P_{\infty}\{X\}}$ is not a formal moduli problem, since $\underline{P_{\infty}\{X\}}(\mathbb{K})$ is in general not contractible. The formal moduli problem $\underline{P_{\infty}\{X\}}^{\psi}$ controlling the formal deformations of a P_{∞} -algebra structure $\psi: P_{\infty} \xrightarrow{} End_X$ on X is defined, on any augmented artinian cdga A, by the homotopy fiber

$$\underline{P_{\infty}\{X\}}^{\psi}(A) = hofib(\underline{P_{\infty}\{X\}}(A) \to \underline{P_{\infty}\{X\}}(\mathbb{K}))$$

taken over the base point ψ , the map being induced by the augmentation $A \to \mathbb{K}$. The twisting of $Hom_{\Sigma}(\overline{C}, End_X)$ by a properad morphism $\psi: P_{\infty} \to End_X$ is the deformation complex of ψ , and we have an isomorphism

$$g_{P,X}^{\psi} = Hom_{\Sigma}(\overline{C}, End_X)^{\psi} \cong Der_{\psi}(\Omega(C), End_X)$$

where the right-hand term is the complex of derivations with respect to ψ [57, Theorem 12].

Proposition 2.11. The tangent L_{∞} -algebra of the formal moduli problem $\underline{P_{\infty}\{X\}}^{\psi}$ is given by

$$g_{P,X}^{\psi} = Hom_{\Sigma}(\overline{C}, End_X)^{\psi}.$$

Proof. Let A be an augmented artinian cdga. By Theorem 2.10, we have the homotopy equivalences

$$\frac{P_{\infty}\{X\}^{\psi}(A)}{=} \sim hofib(\underline{MC_{\bullet}}(g_{P,X})(A) \to \underline{MC_{\bullet}}(g_{P,X})(\mathbb{K}))
= hofib(\underline{MC_{\bullet}}(g_{P,X} \otimes A) \to \underline{MC_{\bullet}}(g_{P,X}))
\simeq \underline{MC_{\bullet}}(hofib_{L_{\infty}}(g_{P,X} \otimes A \to g_{P,X}))$$

where $hofib_{L_{\infty}}(g_{P,X} \otimes A \to g_{P,X})$ is the homotopy fiber, over the Maurer-Cartan element ψ , of the L_{∞} -algebra morphism $g_{P,X} \otimes A \to g_{P,X}$ given by the tensor product of the augmentation $A \to \mathbb{K}$ with $g_{P,X}$. This homotopy fiber is nothing but $g_{P,X}^{\varphi} \otimes m_A$, where m_A is the maximal ideal of A, so there is an equivalence of formal moduli problems

$$\underline{P_{\infty}\{X\}}^{\psi} \simeq \underline{MC_{\bullet}^{fmp}}(g_{P,X}^{\psi}).$$

By Lurie's equivalence theorem, this means that $g_{P,X}^{\psi}$ is the Lie algebra of the formal moduli problem $P_{\infty}\{X\}^{\psi}$.

In this paper, we will also be interested in derived algebraic groups in the sense of [20] and deformation complexes governing homotopy automorphisms of algebras over properads. Let us first note that for any cdga A, the category Mod_A is a (cofibrantly generated) symmetric monoidal model category tensored over chain complexes, so that one can define the category $P_{\infty} - Alg(Mod_A)$ of P_{∞} -algebras in Mod_A .

Definition 2.12. For a chain complex X, we denote $\underline{haut}(X)$ the derived algebraic group of homotopy automorphisms of the underlying complex X taken in the model category of chain complexes. It is defined by

$$A \mapsto haut_{Mod_A}(X \otimes A),$$

where $haut_{Mod_A}$ is the simplicial monoid of homotopy automorphisms in the category of A-modules. Since for every A we have $haut(X) \sim haut_{Mod_A}(X \otimes A)$, it can be defined alternately by the constant functor

$$A \mapsto haut(X)$$
.

Further, let O be an operad, let O_{∞} be a cofibrant resolution of O, and let $(X,\psi:O_{\infty}\to End_X)$ be an O-algebra structure on X. We define $\underline{haut}_{O_{\infty}}(X,\psi)$ to be the derived algebraic group associated to the simplicial monoid of homotopy automorphisms of (X,ψ) in the model category of O_{∞} -algebras, that is the simplicial sub-monoid of self weak equivalences in the usual homotopy mapping space $Map_{O_{\infty}-Alg}(X,X)$. Such simplicial monoids of self weak-equivalences are defined in any model category, see for instance [42, Chapter 17]. The (weak) simplicial presheaf is defined by

$$A \mapsto haut_{O_{\infty}}(X \otimes A, \psi \otimes A)_{Mod_A}$$

where $haut_{O_{\infty}}(X \otimes A, \psi \otimes A)_{Mod_A}$ is the simplicial monoid of homotopy automorphisms of $(X \otimes A, \psi \otimes A) \in O_{\infty} - Alg(Mod_A)$.

By [11] we have a homotopy equivalence $\underline{haut}_{O_{\infty}}(X,\psi) \simeq \underline{L}^H w O_{\infty} - \underline{Alg}(X,\psi)$, where $\underline{L}^H w O_{\infty} - \underline{Alg}(X,\psi)$ is the derived algebraic group of homotopy automorphisms in the Dwyer-Kan hammock localization of O_{∞} -algebras with respect to quasi-isomorphisms [11]. When P is a prop but not an operad, we do not have a

model structure on P_{∞} -algebras to define $\underline{haut}_P(X,\psi)$ but we can still define the hammock localization $\underline{L}^H w P_{\infty} - \underline{Alg}(X,\psi)$. To such a derived algebraic group, one can associate a formal moduli problem $B\underline{L}^H w P_{\infty} - \underline{Alg}(X,\psi)$ given by the strictification (see [1, Section I.2.3.1]) of the weak simplicial presheaf

$$A \in dgArt^{aug}_{\mathbb{K}} \mapsto L^H wP_{\infty} - Alg(Mod_A)(X \otimes A, \psi \otimes A),$$

where B is the classifying complex functor for simplicial monoids (see [20] for the definition of such a formal moduli problem in the case of algebras over operads). We introduce two related and useful classification space constructions. The assignment

$$A \mapsto wP_{\infty} - Alg(Mod_A),$$

where the w(-) stands for the subcategory of weak equivalences, defines a weak presheaf of categories in the sense of [1, Definition I.56], sending a morphism $A \to B$ to the functor $-\otimes_A B$, which is symmetric monoidal, hence lifted at the level of P_{∞} -algebras. This weak presheaf can be strictified into a presheaf of categories. Applying the nerve functor then defines a simplicial presheaf of Dwyer-Kan classification spaces that we note $\underline{\mathcal{N}wP_{\infty}-Alg}$. The simplicial presheaf associated to $A\mapsto Mod_A$ is the simplicial presheaf of quasi-coherent modules of [75, Definition 1.3.7.1] that we note $\underline{\mathcal{N}wCh_{\mathbb{K}}}$. The loop construction on these simplicial presheaves is defined as follows. The loop space on $\underline{\mathcal{N}wP_{\infty}-Alg}$ based at a P_{∞} -algebra (X,ψ) is the strictification of the weak simplicial presheaf

$$\Omega_{(X,\psi)} \mathcal{N} w P_{\infty} - Alg : A \mapsto \Omega_{(X \otimes A, \psi \otimes A)} \mathcal{N} w P_{\infty} - Alg(Mod_A).$$

The loop space on $P_{\infty}\{X\}$ based at a morphism $\psi: P_{\infty} \to End_X$ is the strictification of the weak simplicial presheaf

$$\Omega_{\psi}P_{\infty}\{X\}: A \mapsto \Omega_{\psi \otimes A}P_{\infty} \otimes A\{X \otimes A\}_{Mod_A}.$$

One defines similarly the loop space $\Omega_{\psi} \underline{P_{\infty}\{X\}}^{\psi}$ on the formal moduli problem $P_{\infty}\{X\}^{\psi}$.

Rezk's homotopy pullback theorem [64], generalized to algebras over cofibrant dg props in [83, Theorem 0.1], states that the simplicial moduli space of P_{∞} -algebras structures on X is the homotopy fiber of a forgetful map between the Dwyer-Kan classification space of the ∞ -category of P_{∞} -algebras and the classification space of the ∞ -category of chain complexes, over the base point given by a P_{∞} -algebra structure on X. This result points towards two crucial ideas about the deformation theory of algebras over a prop, which will be crucial in our paper but are of course of independent interest as well.

First, in derived algebraic geometry, a Zariski open immersion of derived Artin stacks $F \hookrightarrow G$ induces a weak equivalence between the tangent complex over a given point of F and the tangent complex over its image in G [75]. We would like a somehow similar statement about the tangent Lie algebras of our formal moduli problems of algebraic structures, when an ∞ -category of algebras "embeds" into another ∞ -category of algebras. The idea is that [83, Theorem 0.1] tells us that the formal moduli problem $P_{\infty}\{X\}^{\psi}$ is the "tangent space" over (X,ψ) to the Dwyer-Kan classification space of the ∞ -category of P_{∞} -algebras. An "immersion" $F: P_{\infty} - Alg \hookrightarrow Q_{\infty} - Alg$ should induce an equivalence between the formal moduli problem of formal deformations of (X,ψ) in P_{∞} -algebras and the formal moduli problem of formal deformations of $F(X,\psi)$ in P_{∞} -algebras, hence

an equivalence between their tangent L_{∞} -algebras. Here the word "immersion" has to be understood as "fully faithful conservative ∞ -functor".

Second, the result [83, Theorem 0.1] implies the existence of a long exact sequence relating the homotopy type of $P_{\infty}\{X\}$ to the homotopy type of the homotopy automorphisms $haut_{P_{\infty}}(X,\psi)$ of $\overline{(X,\psi)}$ inside the ∞ -category of P_{∞} -algebras. These homotopy automorphisms form a derived algebraic group [20] with an associated L_{∞} -algebra $Lie(\underline{haut}_{P_{\infty}}(X,\psi))$, thus there should be a homotopy fiber sequence of L_{∞} -algebras relating $Lie(\underline{haut}_{P_{\infty}}(X,\psi))$ to the tangent L_{∞} -algebra of $P_{\infty}\{X\}^{\psi}$.

To formalize properly the two ideas above, we need the following generalization of Rezk's homotopy pullback theorem:

Proposition 2.13. Let P_{∞} be a cofibrant prop and X be a chain complex.

(1) The forgetful functor $P_{\infty} - Alg \rightarrow Ch_{\mathbb{K}}$ induces a homotopy fiber sequence

$$\underline{P_{\infty}\{X\}} \to \underline{\mathcal{N}wP_{\infty} - Alg} \to \underline{\mathcal{N}wCh_{\mathbb{K}}}$$

of simplicial presheaves over augmented artinian cdgas, taken over the base point X.

(2) This homotopy fiber sequence induces an equivalence of derived algebraic groups

$$\Omega_{\psi} \underline{P_{\infty}\{X\}} \simeq \underline{L^H w P_{\infty} - Alg}(X, \psi).$$

Proof. (1) We explain briefly how [83, Theorem 0.1] can be transposed in the context of simplicial presheaves of cdgas. The identification of the homotopy fiber of the forgetful map

$$\mathcal{N}wP_{\infty} - Alg \to \underline{\mathcal{N}wCh_{\mathbb{K}}}$$

with the simplicial presheaf $P_{\infty}\{X\}$ follows from the two following facts. First, we can identify it pointwise with $Map(P_{\infty}\otimes A, End_{X\otimes A}^{Mod_A})$, where $End_{X\otimes A}^{Mod_A}$ is the endomorphism prop of $X\otimes A$ in the category of A-modules. This comes from the extension of [83, Theorem 0.1] to P_{∞} -algebras in A-modules, which holds true trivially by replacing chain complexes by A-modules as target category in the universal functorial constructions of [83, Section 2.2] (A-modules are equipped with exactly the same operations than chain complexes which are needed in this construction: directs sums, suspensions, twisting cochains). Second, there is an isomorphism of simplicial sets

$$Map(P_{\infty} \otimes A, End_{X \otimes A}^{Mod_A}) \cong P_{\infty}\{X\}(A)$$

(see for instance [82, Section 3]).

(2) The connected components of $\mathcal{N}wP_{\infty} - Alg$ are the classifying complexes of homotopy automorphisms of P_{∞} -algebras, hence a decomposition

$$\mathcal{N}wP_{\infty} - Alg \cong \coprod_{[Y,\phi] \in \pi_0 \mathcal{N}wP_{\infty} - Alg} BL^H wP_{\infty} - Alg(Y,\phi)$$

where $[Y,\phi]$ ranges over quasi-isomorphism classes of P_{∞} -algebras. The moduli space $P_{\infty}\{X\}$ thus decomposes into

$$P_{\infty}\{X\} \cong \coprod_{[X,\psi] \in \pi_0 \mathcal{N} w P_{\infty} - Alg} BL^H w P_{\infty} - Alg(X,\psi)$$

where $[X, \psi]$ ranges over quasi-isomorphism classes of P_{∞} -algebras having X as underlying complex. This induces a homotopy equivalence

$$\Omega_{\psi} P_{\infty} \{X\} \simeq L^H w P_{\infty} - Alg(X, \psi)$$

which sends the constant loop at ψ to $id_{(X,\psi)}$ and any loop to a homotopy automorphism of (X,ψ) . The naturality of this homotopy equivalence follows from the commutativity of the following square for every morphism $f:A\to B$ of augmented artinian cdgas:

$$\Omega_{\psi} \underbrace{P_{\infty}\{X\}}_{(A)}(A) \xrightarrow{\sim} L^{H}wP_{\infty} - Alg(Mod_{A})(X \otimes A, \psi \otimes A) \\
-\otimes_{A}B \downarrow \qquad \qquad -\otimes_{A}B \downarrow \\
\Omega_{\psi} \underbrace{P_{\infty}\{X\}}_{(B)}(B) \xrightarrow{\sim} L^{H}wP_{\infty} - Alg(Mod_{B})(X \otimes B, \psi \otimes B)$$

Hence we get the desired equivalence

$$\Omega_{\psi} \underline{P_{\infty}\{X\}} \simeq \underline{L^H w P_{\infty} - Alg}(X, \psi).$$

The relation between $\Omega_{\psi} \underline{P_{\infty}\{X\}}^{\psi}$ and the derived algebraic group of homotopy automorphisms of (X, ψ) is then given by the following homotopy fiber sequence:

Proposition 2.14. There is a homotopy fiber sequence of derived algebraic groups

$$\Omega_{\psi} P_{\infty} \{X\}^{\psi} \to \underline{L}^H w P_{\infty} - Alg(X, \psi) \to \underline{haut}(X),$$

hence a homotopy fiber sequence of the associated L_{∞} -algebras

$$g^{\psi}_{P,X} \to Lie(\underline{L^H w P_{\infty} - Alg}(X,\psi)) \to Lie(\underline{haut}(X)).$$

Proof. Let A be an augmented artinian cdga. On the one hand, the simplicial set $\underline{P_\infty\{X\}}^\psi(A)$ is identified to the Kan subcomplex of $\underline{P_\infty\{X\}}(A)$ generated by the set of points $\phi: P_\infty \otimes A \to End_{X\otimes A}^{Mod_A}$ such that their reduction modulo A satisfies $\phi \otimes_A \mathbb{K} \sim \psi$, where \mathbb{K} is an A-module via the augmentation $A \to \mathbb{K}$. That is, the simplicial set $P_\infty\{X\}^\psi(A)$ is the moduli space of A-linear deformations of ψ .

On the other hand, the homotopy fiber of the map

$$-\otimes_A \mathbb{K}: L^H w P_{\infty} - Alg(X, \psi)(A) \to haut(X),$$

which is the reduction modulo A induced by the augmentation $A \to \mathbb{K}$, is identified with the Kan subcomplex of $\underline{L^H w P_\infty - Alg}(X, \psi)(A) = L^H w P_\infty - Alg(Mod_A)(X \otimes A, \psi \otimes A)$ generated by the set of points $f: (X \otimes A, \psi \otimes A) \overset{\sim}{\leftarrow} \bullet \overset{\sim}{\to} (X \otimes A, \psi \otimes A)$ such that $f \otimes_A \mathbb{K} \sim id_X$ (recall that vertices of the hammock localization are given by finite zigzags of weak equivalences).

Now recall from Proposition 2.13 that there is a simplicial homotopy equivalence $\Omega_{\psi}\underline{P_{\infty}}\{X\}(A)\cong\Omega_{\psi\otimes A}P_{\infty}\otimes A\{X\otimes A\}_{Mod_A}\simeq L^HwP_{\infty}-Alg(Mod_A)(X\otimes A,\psi\otimes A)$ sending the constant loop at $\psi\otimes A$ to $id_{(X\otimes A,\psi\otimes A)}$ and generally any loop to a homotopy automorphism of $(X\otimes A,\psi\otimes A)$. Such a loop is given by a pointed simplicial map $\partial\Delta^1\to P_{\infty}\otimes A\{X\otimes A\}_{Mod_A}$ sending the point 0 to $\psi\otimes A$. At the level of geometric realizations of these Kan complexes, this is equivalent to define a pointed topological map $l:S^1\to |P_{\infty}\otimes A\{X\otimes A\}_{Mod_A}|$ such that $l(0)=\psi\otimes A$, where 0 is the base point of S^1 . Here we use the fact that the geometric realization is a Quillen equivalence of model categories between simplicial sets with the Kan-Quillen model structure and topological spaces with the model structure induced by weak homotopy equivalences and Serre fibrations. The space $\Omega_{\psi\otimes A}|\underline{P_{\infty}\{X\}}^{\psi}(A)|$ is then the subspace of loops $l:S^1\to |P_{\infty}\otimes A\{X\otimes A\}_{Mod_A}|$ such that for every

point p of S^1 , we have $l(p) \otimes_A \mathbb{K} \sim \psi$. By the geometric realization of the homotopy equivalence above, such loops l are sent to homotopy automorphisms $f_l : (X \otimes A, \psi \otimes A) \stackrel{\sim}{\leftarrow} \bullet \stackrel{\sim}{\rightarrow} (X \otimes A, \psi \otimes A)$ of $(X \otimes A, \psi \otimes A)$ such that $f_l \otimes_A \mathbb{K} \sim id_X$. That is, we get a homotopy equivalence

$$\Omega_{\psi \otimes A} | P_{\infty} \{X\}^{\psi}(A) | \sim |hofib(-\otimes_A \mathbb{K})|$$

hence the simplicial homotopy equivalence above restricts to

$$\Omega_{\psi \otimes A} P_{\infty} \{X\}^{\psi}(A) \sim hofib(-\otimes_A \mathbb{K}).$$

The fiber sequence of L_{∞} -algebras then follows by taking the Lie algebras of the derived algebraic groups as in [20]. Concerning the identification of the fiber with $g_{P,X}^{\psi}$, recall that the Lie algebra of loops over a formal moduli problem gives the tangent Lie algebra of this formal moduli problem.

Remark 2.15. Let us note that $g_{P,X}^{\psi}$ is thus different from the Lie algebra of the loop space $\Omega_{\psi}P_{\infty}\{X\} \simeq \underline{L}^HwP_{\infty} - Alg(X,\psi)$. The later will be described in Section 3.

Relying on these results, we prove the following comparison result:

Theorem 2.16. Let $F: P_{\infty} - Alg \to Q_{\infty} - Alg$ be a fully faithful and conservative ∞ -functor inducing functorially in A, for every augmented artinian edga A, a fully faithful and conservative ∞ -functor $F: P_{\infty} - Alg(Mod_A) \to Q_{\infty} - Alg(Mod_A)$. Then F induces an equivalence of formal moduli problems

$$P_{\infty}\{X\}^{\psi} \sim Q_{\infty}\{F(X)\}^{F(\psi)},$$

where $F(\psi)$ is the Q_{∞} -algebra structure on the image $F(X,\psi)$ of X,ψ under F, hence an equivalence of the associated L_{∞} -algebras

$$g_{P,X}^{\psi} \sim g_{Q,F(X)}^{F(\psi)}$$
.

Proof. Let $F: P_{\infty} - Alg \to Q_{\infty} - Alg$ be a fully faithful ∞ -functor inducing a fully faithful and conservative ∞ -functor $F: P_{\infty} - Alg(Mod_A) \to Q_{\infty} - Alg(Mod_A)$ for every augmented artinian cdga A. At the level of homotopy automorphisms, which can be seen equivalently as loops in Dwyer-Kan's classification spaces, this means that the commutative triangle

$$\underbrace{NwP_{\infty} - Alg} \xrightarrow{NwF} \underbrace{NwQ_{\infty} - Alg}_{NwU},$$


$$\underbrace{NwV}_{NwU} \xrightarrow{NwU}_{NwU}$$

where $\mathcal{N}w(-)$ is the simplicial nerve of the subcategory of weak equivalences (here quasi-isomorphisms) and U is the forgetful functor, induces a commutative triangle

$$\Omega_{(X,\psi)} \underbrace{\mathcal{N}wP_{\infty} - Alg}^{\sim} \xrightarrow{\sim} \Omega_{F(X,\psi)} \underbrace{\mathcal{N}wQ_{\infty} - Alg}_{\Omega_{X} \mathcal{N}wCh_{\mathbb{K}}}$$

where the horizontal arrow induced by $\mathcal{N}wF$ is a weak equivalence by Lemma 2.6.

Consequently, we get the morphism of homotopy fiber sequences of derived algebraic groups



where the homotopy fibers are given by Proposition 2.14. The upper horizontal arrow is a weak equivalence of derived algebraic groups

$$\Omega_{\psi} \underline{P_{\infty}\{X\}}^{\psi} \stackrel{\sim}{\to} \Omega_{F(\psi)} \underline{Q_{\infty}\{F(X)\}}^{F(\psi)},$$

hence an equivalence of the associated L_{∞} -algebras, which are the tangent L_{∞} -algebras of the associated formal moduli problems. By Lurie's equivalence theorem, this means that we have a weak equivalence of formal moduli problems

$$\underline{P_{\infty}\{X\}}^{\psi} \stackrel{\sim}{\to} \underline{Q_{\infty}\{F(X)\}}^{F(\psi)}$$

as well. \Box

In the case of algebras and coalgebras over operads, we can get a similar result for a more general base category:

Proposition 2.17. (1) Let P and Q be two dg operads. Let C be a cofibrantly generated symmetric monoidal model category over $Ch_{\mathbb{K}}$. We assume that the categories of P-algebras and Q-algebras are equipped with their standard model structure. Let

$$F: P - Ala(\mathcal{C}) \rightarrow Q - Ala(\mathcal{C})$$

be a fully faithful and conservative ∞ -functor inducing functorially in A, for every augmented artinian cdga A, a fully faithful and conservative ∞ -functor $F: P_{\infty} - Alg(Mod_A(\mathcal{C})) \to Q_{\infty} - Alg(Mod_A(\mathcal{C}))$, where $Mod_A(\mathcal{C})$ is the category of A-modules in \mathcal{C} . Let $X \in \mathcal{C}$ be equipped with a P-algebra structure $\varphi: \to End_X$ and let $\varphi: Q \to End_{F(X)}$ be its image under the functor F. Then there exists a homotopy equivalence of formal moduli problems

$$\underline{P\{X\}_{\mathcal{C}}}^{\varphi} \simeq \underline{Q\{F(X)\}_{\mathcal{C}}}^{\phi}.$$

(2) Let us now assume that Q satisfies the necessary assumptions so that $Q - Cog(\mathcal{C})$ forms a model category. Let

$$F: P - Alg(\mathcal{C}) \rightarrow Q - Cog(\mathcal{C})$$

be a fully faithful ∞ -functor. Let $X \in \mathcal{C}$ be equipped with a P-algebra structure $\varphi : \to End_X$ and let $\varphi : Q \to coEnd_{F(X)}$ be its image under the functor F. Then there exists a homotopy equivalence of formal moduli problems

$$\underline{P\{X\}_{\mathcal{C}}}^{\varphi} \simeq \underline{Q^{\vee}\{F(X)\}_{\mathcal{C}}}^{\phi},$$

where $Q^{\vee}\{F(X)\}_{\mathcal{C}}$ denotes the simplicial presheaf of Q-coalgebra structures on X.

Proof. Proof of (1). The functor F induces a commutative square of categories

$$P - Alg(\mathcal{C}) \xrightarrow{U \circ F} \mathcal{C}$$

$$\downarrow F \qquad \qquad \downarrow =$$

$$Q - Alg(\mathcal{C}) \xrightarrow{U} \mathcal{C}$$

where $U:Q-Alg(\mathcal{C})\to\mathcal{C}$ is the forgetful functor. In [61], Muro builds a stack of algebras over a nonsymmetric operad $\underline{Alg_{\mathcal{C}}}(P)$, whose evaluation at an algebra A is the classification space of cofibrant P-algebras in A-modules (that is, the nerve of the subcategory of weak equivalences between cofibrant objects). For every algebra A, there is a forgetful functor from P-algebras in A-modules to A-modules, inducing a morphism of simplicial sets $\mathcal{N}wP-Alg(Mod_A)\to\mathcal{N}wMod_A$, hence a morphism of stacks $\underline{Alg_{\mathcal{C}}}(P)\to\underline{QCoh}$ (where \underline{QCoh} is the stack of quasi-coherent modules). For operads in chain complexes over a field of characteristic zero, this construction extends to symmetric operads. Thus the commutative square above also induces a commutative square of stacks

$$\frac{\mathcal{N}wP - Alg(\mathcal{C})}{\downarrow} \longrightarrow \frac{\mathcal{N}w\mathcal{C}}{\downarrow} = \\
\mathcal{N}wQ - Alg(\mathcal{C}) \longrightarrow \frac{\mathcal{N}w\mathcal{C}}{\downarrow}$$

via this construction, hence a morphism of the induced homotopy fiber sequences of stacks.

In order to relate these homotopy fibers to moduli spaces of algebraic structures, we need a symmetric version of [61, Theorem 4.6], which consists in a generalization of Rezk's homotopy pullback theorem [64] to non-symmetric operads in monoidal model categories. A key point in the proof of [61, Theorem 4.6] is that extension and restriction of structures along a weak equivalence of admissible non-symmetric operads form a Quillen pair. However, if one restricts to the setting of algebras in a cofibrantly generated symmetric monoidal model category tensored over another cofibrantly generated symmetric monoidal model category, then a weak equivalence of symmetric Σ -cofibrant operads similarly induces a Quillen pair between the corresponding categories of algebras by [22, Theorem 16.A]. In this framework, [61, Theorem 4.6] can be extended to the symmetric case by following exactly the same argument line. When the base category is $Ch_{\mathbb{K}}$, then the endomorphism operad of any object of \mathcal{C} , defined by the external dg hom of \mathcal{C} , is fibrant because fibrations of operads are defined aritywise and every chain complex over a field of characteristic zero is fibrant. In our situation we can thus drop the fibrancy-cofibrancy condition on X stated in [61, Theorem 4.6]. The stack of P-algebra structures on an A-module M is the homotopy pullback of the morphism $Alg_{\mathcal{C}}(P) \to QCoh$ along the morphism $\mathbb{R}Spec(A) \to QCoh$ representing M [61, Section 5]. This gives a stack version of Rezk's homotopy pullback theorem [64]. Under our assumptions, this theorem holds also in the symmetric case.

The proof concludes as follows. The commutative square of stacks

$$\frac{\mathcal{N}wP - Alg(\mathcal{C})}{\downarrow} \longrightarrow \frac{\mathcal{N}w\mathcal{C}}{\downarrow} = \\
\mathcal{N}wQ - Alg(\mathcal{C}) \longrightarrow \frac{\mathcal{N}w\mathcal{C}}{\downarrow}$$

induces, for every choice of a basepoint $X \in \mathcal{C}$, a commutative square of derived algebraic groups

$$\Omega_{(X,\varphi)} \underbrace{\mathcal{N}wP - Alg(\mathcal{C})}^{\Omega \mathcal{N}w(U \circ F)} \Omega_X \underbrace{\mathcal{N}w\mathcal{C}}_{X} \\
= \Omega_{(F(X),\phi)} \underbrace{\mathcal{N}wQ - Alg(\mathcal{C})}_{\Omega \mathcal{N}wU} \underbrace{\Omega_X \mathcal{N}w\mathcal{C}}_{\Omega \mathcal{N}wU}$$

The homotopy fibers of $\Omega \mathcal{N}w(U \circ F)$ and $\Omega \mathcal{N}wU$ are identified respectively with the formal moduli problems $P\{X\}_{\mathcal{C}}^{\varphi}$ and $Q\{F(X)\}_{\mathcal{C}}^{\varphi}$. The map $\Omega \mathcal{N}wF$ is a weak equivalence, so the homotopy fibers are weakly equivalent as well.

Proof of (2). The arguments are the same than in the proof of (1), using that Q-coalgebra structures on a given object X can be defined by operad morphisms $Q \to coEnd_X$, where $coEnd_X$ is the coendomorphism operad defined by $coEnd_X(n) = Hom_{\mathcal{C}}(X, X^{\otimes n})$.

3. The "plus" construction and the corresponding deformation theory

The plus construction, originally due to Merkulov, will be crucial later in the paper, to consider deformation complexes of algebraic structures which also encode compatible deformations of the differential. In this section, we explain the homotopical counterpart of this construction, that is, how the corresponding deformation complex controls the derived algebraic group of homotopy automorphisms of an algebra over a properad.

Recall from [59] that there is an endofunctor $(-)^+: Prop \to Prop$ on the category of properads which associates to any dg properad P with presentation $\mathcal{F}(E)/(R)$ and differential δ a properad P^+ with presentation $\mathcal{F}(E^+)/(R)$ and differential δ^+ . The Σ -biobject E^+ is defined by $E^+(1,1) = E(1,1) \oplus \mathbb{K}[1]$ and $E^+(m,n) = E(m,n)$ otherwise. This means that we added a generating operation u of degree 1, with one input and one output. We would like this generator to twist the differential of a complex X when we consider a P^+ -algebra structure on X. For this, the differential δ is modified accordingly to have in particular $\delta^+(u) = u \otimes u \in E(1,1) \otimes E(1,1)$. Consequently, properad morphisms $\varphi^+: P^+ \to End_{(X,d)}$ for a given complex X with differential d corresponds to properad morphisms $P \to End_{(X,d-\varphi^+(u))}$ for X equipped with the twisted differential $d - \varphi^+(u)$. In particular, if X is a graded vector space then P^+ -algebra structures on X equip X simultaneously with a P-algebra structure and a compatible differential. Let us reinterpret this construction by defining the following operad:

Definition 3.1. The operad of differentials Di is the quasi-free operad $Di = (\mathcal{F}(E), \partial)$, where $E(1) = \mathbb{K}\delta$ with δ a generator of degree 1, E(n) = 0 for $n \neq 1$ and $\partial(\delta) = \delta \circ \delta$ is the operadic composition $\circ : Di(1) \otimes Di(1) \to Di(1)$.

We will do an abuse of notation and still note Di the properad freely generated by this operad.

Lemma 3.2. Let (V, d_V) be a complex.

- (1) A Di-algebra structure $\phi: Di \to End_V$ on V is a twisted complex $(V, d_V \delta_V)$ where δ_V is the image of the operatic generator δ under ϕ .
- (2) A morphism of Di-algebras $f:(V,d_V-\delta_V)\to (W,d_W-\delta_W)$ is a chain morphism $f:(V,d_V)\to (W,d_W)$ which satisfies moreover $f\circ (d_V-\delta_V)=(d_W-\delta_W)\circ f$ (it is a morphism of twisted complexes).

Proof. (1) The morphism ϕ is entirely determined by the image of the generator δ . Since

$$Di(1) \rightarrow Hom(V, V)$$

is a morphism of complexes, its compatibility with the differentials reads

$$\phi(\partial(\delta)) = d_V \circ \delta_V + \delta_V \circ d_V$$

which gives the equation of twisting cochains

$$\delta_V^2 = d_V \circ \delta_V + \delta_V \circ d_V,$$

hence

$$(d_V - \delta_V)^2 = d_V^2 + \delta_V^2 - d_V \circ \delta_V - \delta_V \circ d_V = 0.$$

(2) A Di-algebra structure on V is given by a morphism $Di(V) \to V$, and a Di-algebra morphism $f: V \to W$ is a chain morphism fitting in the commutative square

$$Di(V) \xrightarrow{Di(f)} Di(W) .$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$V \xrightarrow{f} W$$

Since a Di-algebra structure is determined by the image of the generator δ via $Di(1) \otimes V \to V$, this amounts to the commutativity of the square

$$Di(1) \otimes V \xrightarrow{Di(1) \otimes f} Di(1) \otimes W ,$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$V \xrightarrow{f} W$$

which is exactly saying that f is a morphism of twisted complexes.

Let us note that Di is a non-negatively graded quasi-free operad, hence a cofibrant operad, so we do not need to take a resolution of it to consider the associated simplicial presheaf of Di-algebra structures.

Lemma 3.3. The based loop space at a given complex of the simplicial presheaf of Di-algebra structures is equivalent to the derived algebraic group of homotopy automorphisms of this complex, that is

$$\Omega_{(V,d_V)}\underline{Di\{(V,d_V)\}} \sim \underline{haut(V,d_V)}$$

where the loops are based at the trivial Di-algebra structure $Di \xrightarrow{0} End_{(V,d_V)}$.

Proof. First, recall from Proposition 2.13(2) the equivalence

$$\Omega_{(V,d_V)}Di\{(V,d_V)\} \sim haut_{Di}(V,d_V).$$

There is a fully faithful functor $i:Ch_{\mathbb{K}}\hookrightarrow Di-Alg$ which sends any chain complex X to itself seen as a twisted complex with twisting $\delta_X=0$, that is, the Di-algebra (X,0) for the trivial Di-algebra structure $0:Di\to End_X$. By definition, it sends quasi-isomorphisms of chain complexes to quasi-isomorphisms of complexes with trivial twisting, that is, it preserves weak equivalences. It is clearly conservative as well, and it satisfies $i(X\otimes A)=i(X)\otimes A$ for every augmented artinian cdga A, so it induces a weak equivalence

$$\underline{haut_{Di}(V, d_V)} \sim \underline{haut(V, d_V)}.$$

Let us note that the plus construction obviously preserves quasi-isomorphisms of props. Indeed, the only effect of the plus construction on the cohomology of P is to add a new generator of arity (1,1) to H^*P whose square is zero. Let $\varphi: P \xrightarrow{\sim} Q$ be a quasi-isomorphism of dg props whose collections of generators are respectively E_P and E_Q , such that $E_P^+(1,1) = E(1,1) \oplus \mathbb{K}u_P$ and $E_Q^+(1,1) = E(1,1) \oplus \mathbb{K}u_Q$. Then $H^*(\varphi^+)$ sends $[u_P]$ to $[u_Q]$ (where [-] denotes the cohomology class) and coincides with $H^*(\varphi)$ on the other generators. The only relations satisfied by $[u_P]$ and $[u_Q]$ are that they are both of square zero so $H^*(\varphi^+)$ is still a prop isomorphism, hence φ^+ is a quasi-isomorphism.

The functor $(-)^+$ takes quasi-free properads satisfying the conditions of [57, Corollary 40] to quasi-free properads satisfying the same conditions. By [57, Corollary 40], such properads are cofibrant and by [57, Theorem 42], every properad admits a cofibrant resolution of this form. So to any dg properad P we can associate a cofibrant properad P_+^+ resolving P^+ .

Lemma 3.4. There is a homotopy cofiber sequence of properads

$$Di \to P_{\infty}^+ \to P_{\infty}$$
.

Proof. The initial morphism $i: I \to P_{\infty}$ induces a morphism $i^+: I^+ \to P_{\infty}^+$, and I^+ is nothing but Di so we get our first morphism

$$i^+: Di \to P_{\infty}^+,$$

which is a morphism of cofibrant properads. Now we would like to compare P_{∞}^+ and $P_{\infty} \vee Di$, where \vee stands for the coproduct of properads (see [57, Appendix A.3] for its definition). Since the free properad functor \mathcal{F} is a left adjoint, it preserves coproducts and thus comes with natural isomorphisms $\mathcal{F}(M \oplus N) \cong \mathcal{F}(M) \vee \mathcal{F}(N)$. If we consider the coproduct $P_{\infty} \vee Q_{\infty}$ of two quasi-free properads $P_{\infty} = (\mathcal{F}(M), \partial_P)$ and $Q_{\infty} = (\mathcal{F}(M), \partial_Q)$, then via the previous isomorphism we can define a differential on $\mathcal{F}(M \oplus N)$ by taking the derivation associated to

$$\partial_P|_M \oplus \partial_Q|_N : M \oplus N \to \mathcal{F}(M) \oplus \mathcal{F}(N) \hookrightarrow \mathcal{F}(M \oplus N)$$


by universal property of derivations and the fact that this morphism satisfies the twisting cochain equation. In the case where Q = Di, it turns out that the free

properad underlying P_{∞}^+ is $\mathcal{F}(M \oplus \mathbb{K}d)$ and the differential on P_{∞}^+ (see [59]) coincides with the one above, yielding a properad isomorphism

$$P_{\infty}^+ \cong P_{\infty} \vee Di$$
.

We deduce that the homotopy fiber of i^+ is the homotopy fiber of the canonical map $Di \to P_{\infty} \vee Di$. This map is a cofibration between cofibrant objects, so its homotopy cofiber is a strict cofiber, hence the cokernel of i^+ which is exactly P_{∞} as expected.

Now we fix a diagram of properad morphisms



which induces a homotopy fiber sequence of formal moduli problems

$$P_{\infty}\{X\}^{\psi} \to P_{\infty}^{+}\{X\}^{\psi^{+}} \to Di\{X\}^{0}$$

because the functor $Map(-,End_X)$ is a simplicial mapping space with fibrant target in a model category, so it sends homotopy colimits to homotopy limits. Equivalently this gives a homotopy fiber sequence of the associated L_{∞} -algebras

$$g_{P,X}^{\psi} \to g_{P+,X}^{\psi^+} \to g_{Di,X}^0$$
.

We obtain

Theorem 3.5. There is a quasi-isomorphism of L_{∞} -algebras

$$g_{P+X}^{\psi^+} \simeq Lie(L^H w P_{\infty} - Alg(X, \psi)),$$

in particular

$$g_{P+X}^{\psi^+} \simeq Lie(\underline{haut}_{P_{\infty}}(X,\psi))$$

if P_{∞} is an operad (where $\underline{haut}_{P_{\infty}}(X, \psi)$ is the derived algebraic group from Definition 2.12).

Proof. We have to build a comparison quasi-isomorphism between the homotopy fiber sequence

$$g_{P,X}^{\psi} \to g_{P^+,X}^{\psi^+} \to g_{Di,X}^0$$

and the homotopy fiber sequence


$$g^{\psi}_{P,X} \to Lie(\underline{L^H w P_{\infty} - Alg(X, \psi)}) \to Lie(\underline{haut}(X)).$$

of Proposition 2.14. For this, we consider the following commutative triangle

$$\underbrace{L^{H}wP_{\infty} - Alg(X, \psi)}_{H_{*} \circ U} \xrightarrow{U} \underbrace{haut(X)}_{H^{*}}$$

$$\underbrace{\underline{Aut}(H_{*}X)}$$

where U is the map induced by the forgetful functor, H_* is the map induced by the homology functor and $\underline{Aut}(H_*X)$ is the constant functor $A \mapsto Aut(H_*X)$. This triangle induces a morphism at the level of homotopy fiber sequences



Using the same arguments as in Proposition 2.14, one relates the homotopy automorphisms to loops over the appropriate moduli problems

$$hofib_1 \simeq \Omega_{\psi^+} \underline{P_{\infty}^+} \{X\}^{\psi^+}$$

and

$$hofib_2 \simeq \Omega_0 Di\{X\}^0.$$

Moreover, the base of these homotopy fiber sequences is a discrete space, which implies that the homotopy fiber and the total space have the same homotopy type: indeed, a homotopy fiber sequence $F \to E \to B$ induces a homotopy fiber sequence $\Omega B \to F \to E$. If B is discrete, then ΩB is contractible, which implies that the map $F \to E$ is a homotopy equivalence.

The resulting commutative square

$$\Omega_{\psi^{+}} \underbrace{P_{\infty}^{+} \{X\}}^{\psi^{+}} \xrightarrow{} \Omega_{0} \underbrace{Di\{X\}}^{0}$$

$$\downarrow \sim \qquad \qquad \downarrow \sim$$

$$\underline{L^{H}wP_{\infty} - Alg(X, \psi) \xrightarrow{U} \underbrace{haut}(X)}$$

induces a morphism of homotopy fiber sequences

$$hofib \longrightarrow \Omega_{\psi^{+}} \underbrace{P_{\infty}^{+}\{X\}}^{\psi^{+}} \longrightarrow \Omega_{0} \underline{Di\{X\}}^{0}$$

$$\downarrow \sim \qquad \qquad \downarrow \sim$$

$$\Omega_{\psi} \underline{P_{\infty}\{X\}}^{\psi} \longrightarrow \underline{L^{H}wP_{\infty} - Alg}(X, \psi) \xrightarrow{U} \underline{haut}(X)$$

where the bottom fiber sequence is the one of Proposition 2.14. The upper homotopy fiber hofib can be identified with those loops in $\Omega_{\psi^+} \underbrace{P_{\infty}^+ \{X\}}^{\psi^+}$ which preserves the trivial Di-algebra structure $0: Di \to End_X$ on X, that is, preserving the differential on X. So hofib is nothing but $\Omega_{\psi} \underbrace{P_{\infty} \{X\}}^{\psi}$. We thus get the desired quasi-isomorphism of homotopy fiber sequences.

A useful corollary for us will be the following identification of the tangent complex T_A of an E_n -algebra (Definition 1.18):

Corollary 3.6. The E_n -Hochschild tangent complex T_A of an E_n -algebra A is naturally weakly equivalent as an L_∞ -algebra to $g_{E^+}^{\psi^+}$.

$$T_A \simeq Lie(\underline{haut}_{E_n}(A, \psi)) \simeq g_{E_n^+, A}^{\psi^+},$$

where ψ^+ is the E_n^+ -algebra structure on A trivially induced by its E_n -algebra structure $\psi: E_n \to End_A$ as above, and $\underline{haut}_{E_n}(A)$ is the derived algebraic group of homotopy automorphisms of A as an E_n -algebra.

Proof. According to [20, Lemma 4.31], the homotopy Lie algebra of homotopy automorphisms $Lie(\underline{haut}_{E_n}(A, \psi))$ is equivalent to the tangent complex T_A of A. Hence Theorem 3.5 implies the corollary.

Remark 3.7. Theorem 3.5 shows that the + construction is crucial to study deformation of dg-algebras and not just deformations of algebraic structures on a fixed complex. Let us illustrate the difference between these two moduli problems in a standard example: dg associative algebras. We recall that the operad E_1 is weakly equivalent to the operad Ass of associative dg algebras. By Lemma 3.6, we have that $g_{Ass^+,A}^{\psi^+} \cong T_A$ and by Francis [20], the upper fiber sequence of Theorem 6.7 shows that $T_A \cong Hom(A^{\otimes >0}, A)$ [1] where the right hand side is a sub-complex of the standard Hochschild cochain complex [51] shifted down by 1, with its standard Lie algebra structure (due to Gerstenhaber). On the other hand, a computation similar to the one of Corollary 3.6 using the operad of associative algebras instead of E_n shows that

$$g_{Ass,A}^{\psi} \cong Hom(A^{\otimes > 1}, A)[1]$$

, where the right hand side is just the subcomplex of the previous shifted Hochschild cochain complex where we have removed the Hom(A,A) component.

In general, if \mathcal{O} is an operad, and X is a non-graded \mathcal{O} -algebra, $\underline{\mathcal{O}}_{\infty}\{X\}^{\psi}$ is the moduli space of all \mathcal{O} -algebra structure on X, while $\underline{\mathcal{O}}_{\infty}^+\{X\}^{\psi^+}$ is the moduli space of \mathcal{O} -algebra structures on X up to automorphisms (if X is a chain complex, then the same is true with dg-algebras structures and self-quasi-isomorphisms instead).

4. Bialgebras versus iterated coalgebras

The main purpose of this section is to prove Theorem 0.1(1). Along the way, we establish general results about triple coresolutions and totalizations for dg coalgebras which dualize the ones obtained in [27] and are of independent interest. Part (2) of Theorem 0.1 as well as Corollary 0.2 follows from an adaptation of Part (1) to the context of pointed algebras and will be proved in Section 5.

4.1. **Bar-cobar adjunction.** We recall the construction of bar-cobar adjunctions from [19] for algebras and coalgebras in a stable symmetric monoidal $(\infty, 1)$ -category \mathcal{C} . Let O be an operad with an augmentation $\epsilon: O \to I$, then the associated functor

$$\epsilon^* = triv_O : \mathcal{C} \to O - Alg(\mathcal{C})$$

has a left adjoint

$$\mathcal{B}_O: O-Alg(\mathcal{C}) \to \mathcal{C}.$$

It follows from Barr-Beck-Lurie's theorem [54, Theorem 4.7.4.5] that this bar construction can be enhanced in

$$O - Alg(\mathcal{C}) \xrightarrow{\mathcal{B}_{O}^{enh}} \mathcal{C}$$

where $oblv_O$ is the forgetful functor and $(\mathcal{B}_O \circ triv_O) - Cog(\mathcal{C})$ is the category of coalgebras over the comonad $\mathcal{B}_O \circ triv_O$. Now, by [19, Lemma 3.3.4] there is a morphism of comonads

$$\mathcal{B}_O \circ triv_O \stackrel{\sim}{\to} \overline{F_{BO}^c},$$

where BO is the operadic bar construction on O and $\overline{F_{BO}^c}$ is the comonad whose category of coalgebras is the category $BO - Cog^{conil}(\mathcal{C})$ of conilpotent BO-coalgebras, inducing a new commutative triangle (see [19, Corollary 3.3.5])

$$BO - Cog^{conil}(\mathcal{C})$$

$$O - Alg(\mathcal{C}) \xrightarrow{\mathcal{B}_O} \mathcal{C}$$

In the dual situation, given a cooperad P with a coaugmentation $\eta: P \to I$, there is an adjunction

$$\eta^* = triv_P : \mathcal{C} \leftrightarrows P - Cog^{conil}(\mathcal{C}) : \Omega_P$$

which can be enhanced by Barr-Beck-Lurie's theorem in a commutative triangle

$$P - Cog^{conil}(\mathcal{C}) \xrightarrow{\Omega_P^{enh}} \mathcal{C}$$

where $oblv_P$ is the forgetful functor and $(\Omega_P \circ triv_P) - Alg(\mathcal{C})$ is the category of algebras over the monad $\Omega_P \circ triv_P$. By [19, Lemma 3.3.9] there is a morphism of monads

$$\overline{F_{\Omega P}} \to \Omega_P \circ triv_P$$
,

where ΩP is the operadic cobar construction on P and $\overline{F_{\Omega P}}$ is the monad whose category of algebras is the category $\Omega P - Alg(\mathcal{C})$ of ΩP -algebras, inducing a new commutative triangle (see [19, Corollary 3.3.11])

$$\Omega P - Alg(\mathcal{C}) \qquad .$$

$$P - Cog^{conil}(\mathcal{C}) \xrightarrow{\Omega_P} \mathcal{C}$$

If the operad O is derived Koszul in the sense that the canonical map $\Omega \circ B(O) \xrightarrow{\sim} O$ is a weak equivalence (something always true in our setting), then the two constructions above return an adjunction of ∞ -categories

$$\mathcal{B}_{O}^{enh}: O-Alg(\mathcal{C}) \rightleftarrows BO-Cog^{conil}(\mathcal{C}): \Omega_{O}^{enh}.$$

Let us note that if the operad O is Koszul in the usual sense, for instance when O is an E_n -operad [24], then one can replace BO by the Koszul dual operad $O^!$ in the adjunction above. One then wonders whether such an adjunction is an equivalence of ∞ -categories. Such a result cannot be true in full generality, however, it is conjectured [19, Conjecture 3.4.5] that it holds true when one restricts O-algebras to the subcategory of nilpotent O-algebras. An O-algebra A is nilpotent if there exists an integer N such that the morphism

$$P(n) \to Hom_{\mathcal{C}}(A^{\otimes n}, A),$$

which is the arity n component of the operad morphism $P \to End_A$ defining the P-algebra structure of A, is homotopic to the zero map. However, in our framework we will prove a version of this conjecture that holds for the notion of 0-connected dg algebra, that is, an algebra whose underlying complex is concentrated in positive degrees.


4.2. Plan of the proof of Theorem 0.1(1). We are going to use the following dual version of [54, Theorem 4.7.4.5] in the comonadic setting:

Theorem 4.1. Let C be an ∞ -category.

(1) An adjunction of ∞ -categories

$$F: \mathcal{C} \rightleftarrows \mathcal{D}: G$$

induces a commutative triangle



where $(F \circ G) - Cog(\mathcal{D})$ is the category of coalgebras over the comonad $F \circ G$ in \mathcal{D} and obly is the forgetful functor.

(2) Let us suppose that C admits totalizations $Tot : C^{\Delta} \to C$. If F is conservative and preserves totalizations, then F^{enh} is an equivalence of ∞ -categories.

Remark 4.2. As we will explain with more details in the next subsection, cosimplicial objects in any model category admits a totalization functor, so the theorem above applies in particular to ∞ -categories associated to model categories.

Since we know that the comonads $\mathcal{B}_O \circ triv_O$ and $\overline{F_{BO}^c}$ are weakly equivalent, the associated ∞ -categories of coalgebras are equivalent as well, and the later is nothing but the ∞ -category of conilpotent BO-coalgebras. Thus it remains to prove that \mathcal{B}_O is conservative and commutes with totalizations up to weak equivalence. Our goal is to prove such a result in the case where $\mathcal{C} = dgCog^{conil}$ is the category of conilpotent coassociative coalgebras in non-negatively graded chain complexes.

In order to get explicit models for the bar construction and the totalization, we work in the framework of model categories. If \mathcal{E} is a cofibrantly generated symmetric monoidal model category over a base category \mathcal{C} and O is a Σ -cofibrant operad in \mathcal{C} , then the category of O-algebras form a cofibrantly generated semi-model category, a slight weakening of the notion of model category sufficient for our purposes (see [22, Section 12.1] for the definition and properties of semi-model categories, and [22, Theorem 12.3.A] for the proof of this result). Moreover, the functor η^* is a right Quillen functor and his left adjoint $\eta_! = \mathcal{B}_O$ is consequently a left Quillen functor

[22, Theorem 16.A]. In this context, one can show that \mathcal{B}_O is weakly equivalent to the derived functor of indecomposables (see for instance [51, Chapter 12]). Here we would like to consider dg operads acting on conilpotent dg coalgebras. For this, we need the following result:

Proposition 4.3. The category $dgCog^{conil}$ is a cofibrantly generated closed symmetric monoidal model category.

Proof. We know that $dgCog^{conil}$ forms a cofibrantly generated model category with colimits, weak equivalences and cofibrations created by the forgetful functor. We also know from [2, Lemma 1.3.36] that the chain tensor product of two conilpotent coalgebras is again a conilpotent coalgebra, so that the forgetful functor creates also the tensor product of $dgCog^{conil}$. Since $Ch_{\mathbb{K}}$ satisfies the pushout-product axiom, which is by definition made out of pushout, coproduct, cofibrations and weak equivalences, this implies that $dgCog^{conil}$ satisfies the pushout-product axiom as well. The existence of an internal hom bifunctor follows verbatim from the argument line of [2, Theorem 2.5.1], given that:

- [2, Proposition 2.1.10] restricts to the full subcategory $dgCog^{conil}$ of dgCog;
- [2, Proposition 2.1.12] restricts to the nilpotent case, either by following verbatim the proof of [80, Section 2.1], or by [30, Proposition 1.20] (since we work over a field of characteristic zero, every cooperad is exact);
- the forgetful functor creates small colimits;
- the forgetful functor creates the tensor product of conilpotent coalgebras.

This concludes the proof.

Taking $\mathcal{E}=dgCog^{conil},$ $\mathcal{C}=Ch_{\mathbb{K}}$ and O an E_1 -operad, we get a bar-cobar adjunction

$$\mathcal{B}_{E_1}^{enh}: E_1 - Alg(dgCog^{conil}) \rightleftarrows E_1 - Cog^{conil}(dgCog^{conil}): \Omega_{E_1}^{enh}$$

which forms a Quillen adjunction of semi-model categories. Our main goal is now to prove the following properties of the (not enhanced) bar construction functor:

Theorem 4.4. The functor \mathcal{B}_{E_1} is conservative when restricted to 0-connected E_1 -algebras, and commutes with totalizations up to weak equivalence when restricted to the essential image of a functorial fibrant resolution.

Since we wants to apply the comonadic Barr-Beck-Lurie theorem, we work in an ∞ -categorical setting where the ∞ -category of fibrant objects of a model category is equivalent to the ∞ -category associated to this model category, hence the assumptions of Theorem 4.4. More precisely, the inclusion of the full subcategory of fibrant objects into the model category induces a Dwyer-Kan equivalence of simplicial localizations [11, Proposition 5.2].

4.2.1. Conservativity. We note

$$\mathcal{B}_{E_1}^{\mathbb{K}}: E_1 - Alg(Ch_{\mathbb{K}}) \to Ch_{\mathbb{K}}$$

the left adjoint of the functor η^* induced by the augmentation $\eta: E_1 \to I$ at the level of dg algebras, to distinguish it from the functor

$$\mathcal{B}_{E_1}: E_1 - Alg(dgCog^{conil}) \to dgCog^{conil}$$

induced by the augmentation $\eta: E_1 \to I$ at the level of dg algebras in $dgCog^{conil}$. Recall that this left adjoint is by construction weakly equivalent to the derived functor of the indecomposables, which is the Quillen homology complex computing Quillen homology of algebras over the corresponding operad [51, Section 12.1].

First we want to prove the conservativity of $\mathcal{B}_{E_1}^{\mathbb{K}}$ on 0-connected E_1 -algebras. For this, we use the formalism of right modules over operads thoroughly developed in [22], characterizing in particular bar constructions as functors naturally associated to right modules. We prove that checking the conservativity of $\mathcal{B}_{E_1}^{\mathbb{K}}$ boils down to the conservativity of the classical bar construction of dg associative algebras.

Given the projection morphism $\pi: E_1 \xrightarrow{\sim} Ass$, according to [22, Theorem 7.2.2], there is a natural isomorphism

$$S_{E_1}(B_{E_1}, -) \circ \pi^* \cong S_{Ass}(\pi_! B_{E_1}, -)$$

where π^* is the restriction of structures fitting in the adjunction

$$\pi_!: E_1 - Alg \rightleftarrows Ass - Alg: \pi^*$$

and $\pi_!$ is the extension of structures fitting in the adjunction

$$\pi_1: E_1 - Mod \rightleftharpoons Ass - Mod: \pi^*$$

between right Ass-modules and right E_1 -modules. Since π is a weak equivalence, by [22, Theorem 16.B] the adjunction above between right modules is a Quillen equivalence, so the weak equivalence of right E_1 -modules $B_{E_1} \stackrel{\sim}{\to} \pi^* B_{Ass}$ corresponds by adjunction to a weak equivalence of right Ass-modules $\pi_! B_{E_1} \stackrel{\sim}{\to} B_{Ass}$. Moreover, since $\pi_!$ is a left Quillen functor, the right module $\pi_! B_{E_1}$ is still cofibrant, so this is a weak equivalence of cofibrant right Ass-modules. By [22, Theorem 15.1.A], it induces consequently a natural weak equivalence

$$S_{E_1}(B_{E_1}, -) \circ \pi^* \cong S_{Ass}(\pi_! B_{E_1}, -) \stackrel{\sim}{\to} S_{Ass}(B_{Ass}, -),$$

hence

$$\mathcal{B}_{E_1}^{\mathbb{K}}\circ\pi^*\sim Bar$$

where π^* is the restriction of structures and Bar is the operadic bar construction on dg associative algebras. The functor Bar is conservative on 0-connected associative algebras (alternately, the classical bar construction is conservative on connected associative algebras), see for instance [51, Chapter 11]. By the natural weak equivalence above, this implies that $\mathcal{B}_{E_1}^{\mathbb{K}} \circ \pi^*$ is conservative on 0-connected associative algebras as well. By the Quillen equivalence between dg associative algebras and E_1 -algebras, the conservativity of $\mathcal{B}_{E_1}^{\mathbb{K}} \circ \pi^*$ implies that the functor $\mathcal{B}_{E_1}^{\mathbb{K}}$ itself is conservative when restricted to 0-connected E_1 -algebras. Now recall from [22, Section 3.3.5] that \mathcal{B}_{E_1} , as an extension functor adjoint to the restriction functor η^* , is obtained by a certain reflexive coequalizer

$$E_1(A) \rightrightarrows A \to \mathcal{B}_{E_1}(A)$$

in $dgCog^{conil}$ for every $A \in ob(E_1 - Alg(dgCog^{conil}))$, where $E_1(A)$ is the free E_1 -algebra on A determined by the formula

$$E_1(A) = \bigoplus_{n \ge 1} (E_1(n) \otimes A^{\otimes n})_{\Sigma_n}.$$

Here $(-)_{\Sigma_n}$ denotes the coinvariants under the diagonal action of the symmetric group Σ_n . Since the forgetful functor $U: dgCog^{conil} \to Ch_{\mathbb{K}}$ creates the tensor product and the small colimits, we actually have

$$U \circ \mathcal{B}_{E_1} = \mathcal{B}_{E_1}^{\mathbb{K}}.$$

Weak equivalences in $E_1 - Alg(dgCog^{conil})$ and $dgCog^{conil}$ are created in chain complexes, so the conservativity of $\mathcal{B}_{E_1}^{\mathbb{K}}$ on $E_1 - Alg^{0-con}(Ch_{\mathbb{K}})$, where the upper script 0-con stands for 0-connected algebras, implies the conservativity of \mathcal{B}_{E_1} on $E_1 - Alg^{0-con}(dgCog^{conil})$.

4.2.2. Commuting with totalizations. We need the conormalization of a cosimplicial coalgebra structure to be compatible with an E_1 -algebra structure. The conormalization of cosimplicial coalgebras forms a lax monoidal functor (see [68, Section 2]) for a good review of its properties in the simplicial case) that consequently lifts to cosimplicial E_1 -algebras in dg coalgebras. Let us note

$$Tot_{E_1-Ass^{op}}: E_1-Alg(dgCog^{conil})^{\Delta} \to E_1-Alg(dgCog^{conil})$$

and

$$Tot_{Ass^{op}}: (dgCog^{conil})^{\Delta} \to dgCog^{conil}$$

the totalization functors, which exist since we work with model categories). Let us also note $Res^{\bullet}(C) \in E_1 - Alg(dgCog^{conil})^{\Delta}$, for $C \in E_1 - Alg(dgCog^{conil})$, a certain functorial cosimplicial resolution of C which will be defined in the next section and whose totalization gives a functorial fibrant resolution of C. Our aim is to prove the following properties of the bar construction with respect to these functors:

Proposition 4.5. (1) There is an isomorphism

$$Tot_{E_1-Ass^{op}}(Res^{\bullet}(C)) \stackrel{\cong}{\to} Tot_{Ass^{op}}(Res^{\bullet}(C))$$

 $in E_1 - Alg(dgCog^{conil}).$

(2) There is an isomorphism

$$\overline{N^*}Res^{\bullet}(C) \stackrel{\sim}{\cong} Tot_{Ass^{op}}(Res^{\bullet}(C))$$

in $E_1 - Alg(dgCog^{conil})$, where $\overline{N^*}$ is the reduced conormalization.

(3) There is a weak equivalence

$$\mathcal{B}_{E_1}(\overline{N^*}Res^{\bullet}(C)) \stackrel{\sim}{\to} \overline{N^*}\mathcal{B}_{E_1}(Res^{\bullet}(C))$$

in $dgCog^{conil}$.

This gives the commutation of the bar construction \mathcal{B}_{E_1} with the totalizations, up to weak equivalence, on the image of a functorial fibrant resolution on E_1 – $Alg(dgCog^{conil})$. The two next subsections are devoted to define explicit models for totalizations and cosimplicial resolutions that will allow us to prove this proposition.

4.3. Triple coresolution and totalization for dg coalgebras. The crux of our method here is to adapt to dg coalgebras a dual version of the operadic arguments of [26, Appendix B], [27] and [28, Lemma B.8].

The adjunction

$$U: dgCog^{conil} \rightleftarrows Ch_{\mathbb{K}}: F^{c}$$

between the forgetful functor U and the cofree coalgebra functor F^c gives rise to a comonad $\overline{F^c} = U \circ F^c$ over $Ch_{\mathbb{K}}$ equipped with a coproduct $\nu : \overline{F^c} \to \overline{F^c} \circ \overline{F^c}$,

a counit $\epsilon: \overline{F^c} \to Id$ and a coaction $\rho: F^c \to F^c \circ \overline{F^c}$ of F^c on $\overline{F^c}$. For every $C \in dgCog^{conil}$, the triple coresolution $Res^{\bullet}(C) \in (dgCog^{conil})^{\Delta}$ is defined by

$$Res^{n}(C) = F^{c} \circ \underbrace{F^{c} \circ \cdots \circ F^{c}}_{n}(U(C))$$

= $F^{c}(K^{n}(C))$

where

$$K^n(C) = \underbrace{\overline{F^c} \circ \cdots \circ \overline{F^c}}_{n}(U(C)).$$

The cofaces

$$d^i: Res^{n-1}(C) \to Res^n(C)$$

are induced by ρ for i=0, by ν on the i^{th} factor for $1\leq i\leq n-1$ and by the coalgebra structure $U(C)\to F^c(U(C))$ of C for i=n. The codegeneracies

$$s^j: Res^{n+1}(C) \to Res^n(C)$$

are induced by ϵ on the $(j+i)^{th}$ factor.

Now recall that for every model category \mathcal{C} , the category of cosimplicial objects \mathcal{C}^{Δ} can be equipped with the Reedy model structure. The existence of simplicial frames in any model category (see [42, Chapter 16]) ensures that for any cosimplicial object $C^{\bullet} \in \mathcal{C}^{\Delta}$, we can pick a simplicial frame $(C^{\bullet})^{\Delta^{\bullet}}$ which forms a simplicial object in \mathcal{C}^{Δ} . This allows us to define the totalization of C^{\bullet} by the end formula

$$Tot(C^{\bullet}) = \int_{n \in \Delta} (C^n)^{\Delta^n}.$$

When C^{\bullet} is Reedy fibrant, any simplicial frame on C^{\bullet} is a simplicial resolution of C^{\bullet} , and the properties of such resolutions implies that the totalization of C^{\bullet} is a fibrant object of C which is invariant, up to weak equivalence, under a change of simplicial frame.

Proposition 4.6. For any dg coalgebra C, the cosimplicial dg coalgebra $Res^{\bullet}(C)$ is Reedy fibrant.

Proof. The cofaces and codegeneracies of $Res^{\bullet}(C)$ restrict to $K^{\bullet}(C)$ except for d^0 , so we set $d^0=0$ to get a full cosimplicial structure on $K^{\bullet}(C)$ which induces the cosimplicial structure of $Res^{\bullet}(C)$ by applying the cofree coalgebra functor. Then the remaining part of the proof is essentially a dual version of the proof of [27, Proposition 2.2]. The tree-decomposition argument of [27, Proposition 2.2] dualizes to the coalgebra setting by using trees with the converse orientation (one single input and several outputs), so that the n-simplices $K^n(C)$ decompose into a direct sum

$$K^n(C) = M^n K^{\bullet}(C) \oplus N^n K^{\bullet}(C)$$

where $M^nK^{\bullet}(C)$ is the n^{th} matching object of $K^{\bullet}(C)$ and $N^nK^{\bullet}(C)$ is the degree n part of the conormalization of C (for the graduation induced by the cosimplicial structure of $K^{\bullet}(C)$). Moreover, we have $M^nRes^{\bullet}(C) = F^c(M^nK^{\bullet}(C))$ because F^c commutes with small limits and the codegeneracies of $Res^{\bullet}(C)$ are induced by those of $K^{\bullet}(C)$ under F^c , hence

$$Res^n(C) = M^n Res^{\bullet}(C) \wedge F^c(N^n K^{\bullet}(C))$$

where \wedge stands for the product in $dgCog^{conil}$. The matching morphisms

$$\mu_n: Res^n(C) \to M^n Res^{\bullet}(C)$$

are the projections on the first factor defined by the pullbacks

$$Res^{n}(C) \longrightarrow F^{c}(N^{n}K^{\bullet}(C))$$

$$\downarrow^{\mu_{n}} \qquad \qquad \downarrow$$

$$M^{n}Res^{\bullet}(C) \longrightarrow 0.$$

By definition of the model structure on $dgCog^{conil}$, the cofree coalgebra functor F^c is a right Quillen functor so $F^c(N^nK^{\bullet}(C))$ is fibrant (every chain complex over a field is fibrant in the projective model structure). Since fibrations are stable under pullbacks, the maps μ_n are fibrations for every integer n, which exactly means that $Res^{\bullet}(C)$ is Reedy fibrant.

Our goal is now to prove that the totalization of the triple coresolution gives a (functorial) fibrant resolution in $dgCog^{conil}$. For this, we give an explicit model of this totalization by chosing an appropriate simplicial frame. Recall that for any cosimplicial set K^{\bullet} , its conormalization $N^*(K^{\bullet})$ forms an augmented simplicial associative algebra, and dually its normalization $N_*(K_{\bullet})$ forms a coaugmented cosimplicial coassociative coalgebra (they have actually a richer structure, respectively that of an algebra and a coalgebra over the Barratt-Eccles operad [6]). The coaugmentation ideal of $N_*(K_{\bullet})$, that is, its reduced normalization $\overline{N_*}(K_{\bullet})$, hence forms an object of $dgCog^{conil}$.

Proposition 4.7. The collection $Res^{\bullet}(C)^{\Delta^{\bullet}} = \{F^c(K^n(C) \otimes \overline{N^*}(\Delta^m))\}_{n,m \in \mathbb{N}}$ forms a simplicial frame of $Res^{\bullet}(C)$ in $(dgCog^{conil})^{\Delta}$.

Proof. We first show that the functor $(-) \otimes \overline{N^*}(\Delta^{\bullet})$ defines a simplicial frame on the full subcategory $F^c(Ch_{\mathbb{K}})$ of $dgCog^{conil}$ formed by cofree coalgebras. Let $C \in dgCog^{conil}$ and $X \in Ch_{\mathbb{K}}$, we have a sequence of natural isomorphisms

$$\begin{array}{ccc} Mor_{dgCog^{conil}}(C, F^{c}(X \otimes \overline{N^{*}}(\Delta^{m}))) & \stackrel{\cong}{\to} & Mor_{Ch_{\mathbb{K}}}(C, X \otimes \overline{N^{*}}(\Delta^{m})) \\ & \stackrel{\cong}{\to} & Mor_{Ch_{\mathbb{K}}}(C \otimes \overline{N_{*}}(\Delta^{m}), X) \\ & \stackrel{\cong}{\to} & Mor_{dgCog^{conil}}(C \otimes \overline{N_{*}}(\Delta^{m}), F^{c}(X)) \end{array}$$

where the first line follows from the universal property of the free-forgetful adjunction, the second line follows from the dualization of the conormalization, and the third line from the free-forgetful adjunction. Let us note that the third line makes sense because the reduced normalization is a coalgebra and the tensor product of coalgebra is defined in chain complexes. We deduce that the functor

$$(-)^{\Delta^m}: F^c(X) \mapsto F^c(X \otimes \overline{N^*}(\Delta^m))$$

is right adjoint to the functor

$$(-)\otimes\Delta^m:C\mapsto C\otimes\overline{N_*}(\Delta^m)$$

on the full subcategory $F^c(Ch_{\mathbb{K}})$ of $dgCog^{conil}$. Consequently, the functor $(-)^{\Delta^{\bullet}}$ defines a simplicial frame on $F^c(Ch_{\mathbb{K}})$ if and only if $(-) \otimes \Delta^{\bullet}$ defines a cosimplicial frame. Since the forgetful functor $dgCog^{conil} \to Ch_{\mathbb{K}}$ defines small colimits,

cofibrations and weak equivalences, the later property follows from the fact that $(-) \otimes \overline{N_*}(\Delta^{\bullet})$ is a cosimplicial frame on $Ch_{\mathbb{K}}$.

Finally, the fact that $(-)^{\Delta^{\bullet}}$ is a simplicial frame on $F^c(Ch_{\mathbb{K}})$ implies that $Res^{\bullet}(C)^{\Delta^{\bullet}}$ is a simplicial frame of $Res^{\bullet}(C)$. Indeed, the morphism

$$\epsilon^* : Res^{\bullet}(C)^{\Delta^0} \to Res^{\bullet}(C)^{\Delta^n}$$

induced by the constant map $\epsilon: \{0, \cdot, n\} \to \{0\}$ defines a weak equivalence in each dimension because $Res^{\bullet}(C)$ is fibrant in each dimension (since it is Reedy fibrant). By definition of weak equivalences in the Reedy model structure of $(dgCog^{conil})^{\Delta}$, this means that ϵ^* is a weak equivalence of $(dgCog^{conil})^{\Delta}$ for every integer n. The simplicial frame construction $Res^{\bullet}(C)^{\Delta^{\bullet}}$ can be extended to an arbitrary simplicial set $K \in sSet$ to give a simplicial-cosimplicial object $Res^{\bullet}(C)^{K}$, and checking the fibration condition in the definition of a simplicial frame amounts to check that for every integer n, the map

$$i^*: Res^{\bullet}(C)^{\Delta^n} \to Res^{\bullet}(C)^{\partial \Delta^n}$$

induced by the inclusion $i^*: \partial \Delta^n \hookrightarrow \Delta^n$ forms a fibration in the base category for any n>0. Here the base category is $(dgCog^{conil})^{\Delta}$, which means that i^* has to be a Reedy fibration of cosimplicial objects. By definition, we thus have to prove that the pullback-corner map

$$(i^*, \mu^*): Res^r(C)^{\Delta^n} \to M_r Res^{\bullet}(C)^{\Delta^n} \times_{M_r Res^{\bullet}(C)^{\partial \Delta^n}} Res^r(C)^{\partial \Delta^n}$$

induced by the matching map $\mu: Res^r(C) \to M_r Res^{\bullet}(C)$ and by i is a fibration of dg coalgebras. The cofree coalgebra functor F^c commutes with limits, so the pullback-corner map above is actually given by

$$(i^*, \mu^*) : F^c(K^r(C) \otimes \overline{N_*}(\Delta^n)) \to F^c\left(M_r K^{\bullet}(C) \otimes \overline{N_*}(\Delta^n) \times_{M_r K^{\bullet}(C) \otimes \overline{N_*}(\partial \Delta^n)} K^r(C) \otimes \overline{N_*}(\partial \Delta^n)\right)$$

which is exactly the image under F^c of the pullback-corner map of complexes induced by i and the matching map of $K^{\bullet}(C)$. Since F^c preserves fibrations between fibrant objects as a right Quillen functor, and since every chain complex is fibrant, the proof boils down to check that the map

$$(i^*, \mu^*): K^r(C) \otimes \overline{N_*}(\Delta^n) \to M_r K^{\bullet}(C) \otimes \overline{N_*}(\Delta^n) \times_{M_r K^{\bullet}(C) \otimes \overline{N_*}(\partial \Delta^n)} K^r(C) \otimes \overline{N_*}(\partial \Delta^n)$$

is a fibration of chain complexes. Dualizing the normalization functor in complexes, this map is, up to isomorphism, given by

$$(i^*,\mu^*): Hom_{dg}(\overline{N^*}(\Delta^n),K^r(C)) \to Hom_{dg}(\overline{N^*}(\Delta^n),M_rK^{\bullet}(C)) \times_{Hom_{dg}(\overline{N^*}(\partial\Delta^n),M_rK^{\bullet}(C))} Hom_{dg}(\overline{N^*}(\partial\Delta^n),K^r(C))$$

. We conclude that this is a fibration by applying the dual pushout-product axiom, which holds true in $Ch_{\mathbb{K}}$.

Proposition 4.8. The coaugmentation $\eta: C \to Tot(Res^{\bullet})$ induced by the counit $\epsilon(C): C \to Res^{0}(C) = Res^{0}(C)^{\Delta^{0}}$ is a quasi-isomorphism of dg coalgebras.

Proof. This proof follows an argument line similar to the one of [28, Lemma B.8]. The totalization of $Res^{\bullet}(C)$ is defined by the end

$$Tot(Res^{\bullet}(C)) = \int_{\underline{n} \in \Delta} F^{c}(K^{n}(C) \otimes \overline{N^{*}}(\Delta^{n})).$$

We consider the following composition of maps

$$\int_{\underline{n}\in\Delta} F^{c}(K^{n}(C)\otimes \overline{N^{*}}(\Delta^{n})) \xrightarrow[(1)]{} \int_{\underline{n}\in\Delta} K^{n}(C)\otimes \overline{N^{*}}(\Delta^{n})$$


$$\overset{\cong}{\underset{(2)}{\longrightarrow}} \overline{N^{*}}K^{\bullet}(C)$$

$$\xrightarrow[(3)]{} \overline{\Omega}(C)$$

where (1) is the canonical projection, (2) is a formal isomorphism and (3) is the projection on maximal simplices in $K^{\bullet}(C)$ (see [26, Appendix C.2.16] for the operadic case) from the reduced conormalized complex of the cosimplicial object $K^{\bullet}(C)$ to the reduced cobar construction $\overline{\Omega}(C)$ of C. The composite (1-3) lifts to a morphism $f: Tot(Res^{\bullet}(C)) \to Bar\overline{\Omega}(C)$, where Bar is the usual bar construction on augmented dg algebras. Using the weight decomposition of the cofree coalgebra functor $F^c = \bigoplus_{s \geq 1} F_s^c$, we apply the argument of [28, Lemma B.8] to get a morphism of spectral sequences associated to the filtration of f induced by this decomposition, such that on the E_0 page, the morphism of graded objects induced by f is given by

$$E^0 f = F^c(\int_{\underline{n} \in \Delta} K^n(C) \otimes \overline{N^*}(\Delta^n) \underbrace{\longrightarrow}_{(2-3)} \overline{\Omega}(C)).$$

The arrow (2) is a formal isomorphism, and the arrow (3) is a quasi-isomorphism (see [21] for a comparison between reduced conormalization and reduced cobar construction in the operadic case), so E^0f is a quasi-isomorphism, hence f is a quasi-isomorphism as well. We conclude that η is a quasi-isomorphism via the commutative triangle



We conclude:

Theorem 4.9. The functor $Tot(Res^{\bullet}(-))$ defines a functorial fibrant resolution on $dqCoq^{conil}$.

We can thus restrict ourselves to prove the desired properties of \mathcal{B}_{E_1} on cosimplicial coalgebras of the form $Res^{\bullet}(C)$.

4.4. **Totalization versus bar construction.** In this section we prove Proposition 4.5.

Lemma 4.10. For any $C \in E_1 - Alg(dgCog^{conil})$, there is an E_1 -algebra structure on $Tot_{Ass^{op}}(Res^{\bullet}(C))$ compatible with its coalgebra structure so that we can chose $Tot_{Ass^{op}}(Res^{\bullet}(C))$ as a model for $Tot_{E_1-Ass^{op}}(Res^{\bullet}(C))$.

Proof. Recall that since $E_1 - Alg(dgCog^{conil})^{\Delta}$ is a Reedy model category, we can define a totalization functor

$$Tot_{E_1 - Ass^{op}}(C^{\bullet}) = \int_{n \in \Delta}^{E_1 - Alg(dgCog^{conil})} (C^n)_{E_1 - Ass^{op}}^{\Delta^n}$$

where $C^{\bullet} \in E_1 - Alg(dgCog^{conil})^{\Delta}$, $(-)_{E_1 - Ass^{op}}^{\bullet}$ is a simplicial frame in $E_1 - Alg(dgCog^{conil})^{\Delta}$, and the limit $\int_{\underline{n} \in \Delta}^{E_1 - Alg(dgCog^{conil})}$ is taken in $E_1 - Alg(dgCog^{conil})$.

The cofree coalgebra functor is defined by the reduced tensor coalgebra $F^c(C) = \bigoplus_{n \geq 1} C^{\otimes n}$ equipped with the deconcatenation product, and comes with a diagonal morphism

$$F^{c}(C \otimes D) = \bigoplus_{n \geq 1} C^{\otimes n} \otimes D^{\otimes n} \to \bigoplus_{m,n \geq 1} C^{\otimes m} \otimes D^{\otimes n} = F^{c}(C) \otimes F^{c}(D)$$

(recall that the tensor product of coalgebras is defined in chain complexes) natural in C and D, so that F^c is an oplax monoidal functor. Moreover, the forgetful functor U is monoidal by construction of the tensor product in $dgCog^{conil}$, so the Quillen adjunction between U and F^c lifts to a Quillen adjunction

$$U: E_1 - Alg(dgCog^{conil}) \rightleftarrows E_1 - Alg(Ch_{\mathbb{K}}): F^c$$

(see [22, Proposition 12.3.3]). This implies that if C is an object of E_1 -Alg($dgCog^{conil}$), then its E_1 -algebra structure is compatible with the cosimplicial constructions $K^{\bullet}(C)$ and $Res^{\bullet}(C)$ so that these cosimplicial coalgebras are actually cosimplicial E_1 -algebras in coalgebras.

Now, recall that the reduced conormalization $\overline{N^*}(\Delta^n)$ is an algebra over the Barratt-Eccles operad BE (see [6]). For any $C \in E_1 - Alg(dgCog^{conil})$, the chain tensor product $K^n(C) \otimes \overline{N^*}(\Delta^n)$ thus forms an $E_1 \otimes_H BE$ -algebra in $Ch_{\mathbb{K}}$, where \otimes_H is the arity-wise tensor product of operads (also called the Hadamard tensor product [51, Section 5.3.2]). Since E_1 , like any operad over a field of characteristic zero, is equipped with an operad morphism $E_1 \to E_1 \otimes_H BE$, the tensor product $K^n(C) \otimes \overline{N^*}(\Delta^n)$ is actually an E_1 -algebra in $Ch_{\mathbb{K}}$. This implies, in turn, that $F^c(K^n(C) \otimes \overline{N^*}(\Delta^n))$ belongs to $E_1 - Alg(dgCog^{conil})$. Moreover, limits in $E_1 - Alg(dgCog^{conil})$ are determined by the forgetful functor, so

$$\int_{\underline{n}\in\Delta}^{E_1-Alg(dgCog^{conil})}F^c(K^n(C)\otimes\overline{N^*}(\Delta^n)) = \int_{\underline{n}\in\Delta}^{dgCog^{conil}}F^c(K^n(C)\otimes\overline{N^*}(\Delta^n))$$
$$= Tot_{Ass^{op}}(Res^{\bullet}(C))$$

where the last line follows from Proposition 4.7, and the totalization of $Res^{\bullet}(C)$ in coalgebras inherits a compatible E_1 -algebra structure. We conclude by noticing that the argument line of the proof of Proposition 4.7 can be literally transposed here: the simplicial-cosimplicial object $F^c(K^{\bullet}(C) \otimes \overline{N^*}(\Delta^{\bullet}))$ defines a simplicial frame of $Res^{\bullet}(C)$ in $E_1 - Alg(dgCog^{conil})^{\Delta}$ because for any $X \in E_1 - Alg(Ch_{\mathbb{K}})$,

the simplicial object $F^c(X \otimes \overline{N^*}(\Delta^{\bullet}))$ defines a simplicial frame of $F^c(X)$ in E_1 – $Alg(dgCog^{conil})$.

Lemma 4.11. For any $C \in E_1 - Alg(dgCog^{conil})$, there is an isomorphism

$$Tot_{Ass^{op}}(Res^{\bullet}(C)) \stackrel{\cong}{\to} N^*Res^{\bullet}(C)$$

 $in E_1 - Alg(dgCog^{conil}).$

Proof. There is an isomorphism

$$\int_{\underline{n}\in\Delta}^{E_1-Alg(Ch_{\mathbb{K}})}K^n(C)\otimes\overline{N^*}(\Delta^n)=\int_{\underline{n}\in\Delta}^{Ch_{\mathbb{K}}}K^n(C)\otimes\overline{N^*}(\Delta^n)\stackrel{\cong}{\to} \overline{N^*}K^{\bullet}(C)$$

in $E_1 - Alg(Ch_{\mathbb{K}})$, inducing an isomorphism

$$F^c(\int_{n\in\Delta}^{Ch_{\mathbb{K}}}K^n(C)\otimes\overline{N^*}(\Delta^n))\stackrel{\cong}{\to} F^c(\overline{N^*}K^\bullet(C))$$

in $E_1 - Alg(dgCog^{conil})$. We have equalities

$$F^c(\int_{\underline{n}\in\Delta}^{Ch_{\mathbb{K}}}K^n(C)\otimes\overline{N^*}(\Delta^n))=\int_{\underline{n}\in\Delta}^{dgCog^{conil}}F^c(K^n(C)\otimes\overline{N^*}(\Delta^n)))$$

because F^c commutes with limits and

$$F^c(\overline{N^*}K^{\bullet}(C)) = \overline{N^*}Res^{\bullet}(C)$$

because $Res^{\bullet}(C) = F^{c}(K^{\bullet}(C))$ and the cosimplicial structure of $Res^{\bullet}(C)$ is induced by the one of $K^{\bullet}(C)$ under F^{c} , hence an isomorphism

$$\int_{\underline{n}\in\Delta}^{dgCog^{conil}}F^c(K^n(C)\otimes\overline{N^*}(\Delta^n)))\overset{\cong}{\to}\overline{N^*}Res^{\bullet}(C).$$

The source of this isomorphism is nothing but $Tot_{Ass^{op}}(Res^{\bullet}(C))$.

Lemma 4.12. For any $C \in E_1 - Alg(dgCog^{conil})$, there is a weak equivalence

$$\mathcal{B}_{E_1}(\overline{N^*}Res^{\bullet}(C)) \stackrel{\sim}{\to} \overline{N^*}\mathcal{B}_{E_1}(Res^{\bullet}(C))$$

in $dqCoq^{conil}$.

Proof. We start by proving that F^c commutes with reflexive coequalizers of E_1 -algebras. Let I be the small category whose associated diagram is given by

$$0 \xrightarrow{} 1$$

and $\{X_i\}_{i\in ob(I)}$ be an *I*-diagram

$$X_0 \xrightarrow{d_0} X_1$$

in $E_1 - Alg(Ch_{\mathbb{K}})$ such that $s_0 \circ d_0 = s_0 \circ d_1 = id_{X_0}$. The colimit $colim_i X_i$ of such a diagram is called a reflexive coequalizer. We have the following series of

isomorphisms in $E_1 - Alg(dgCog^{conil})$:

$$F^{c}(colim_{i}X_{i}) = \bigoplus_{n\geq 1} (colim_{i}X_{i})^{\otimes n}$$

$$\cong \bigoplus_{n\geq 1} colim_{i}(X_{i})^{\otimes n}$$

$$\cong colim_{i} \bigoplus_{n\geq 1} X_{i}^{\otimes n}$$

$$= colim_{i}F^{c}(X_{i}).$$

The first line is by definition of F^c as the reduced tensor coalgebra. Since E_1 is a Hopf operad, $E_1 - Alg(Ch_{\mathbb{K}})$ forms a symmetric monoidal category for the chain tensor product. Moreover, according to [22, Proposition 3.3.1], reflexive coequalizers of E_1 -algebras are also determined in chain complexes, and according to [22, Proposition 1.2.3] the chain tensor power functors preserves reflexive coequalizers, hence the second line. The third line is just the commutation of two colimits of E_1 -algebras, and the last line follows from the fact that colimits of coalgebras are determined by the forgetful functor.

Now, recall from 4.2.1 that \mathcal{B}_{E_1} is constructed as a reflexive coequalizer of E_1 -algebras, so that

$$F^c \circ \mathcal{B}_{E_1}^{\mathbb{K}} = \mathcal{B}_{E_1} \circ F^c.$$

We have

$$\mathcal{B}_{E_1}(\overline{N^*}Res^{\bullet}(C)) = \mathcal{B}_{E_1}(F^c(\overline{N^*}K^{\bullet}(C)))$$
$$= F^c(\mathcal{B}_{E_1}^{\mathbb{K}}(\overline{N^*}K^{\bullet}(C)))$$

by construction of the cosimplicial structures of $K^{\bullet}(C)$ and $Res^{\bullet}(C)$ and the commutation rule above, and

$$\overline{N^*}\mathcal{B}_{E_1}(Res^{\bullet}(C)) = F^c(\overline{N^*}\mathcal{B}_{E_1}^{\mathbb{K}}(K^{\bullet}(C)))$$

by the commutation of F^c with $\overline{N^*}$ and \mathcal{B}_{E_1} as explained above. Consequently, to build a weak equivalence

$$\mathcal{B}_{E_1}(N^*Res^{\bullet}(C)) \stackrel{\sim}{\to} N^*\mathcal{B}_{E_1}(Res^{\bullet}(C))$$

in $dgCog^{conil}$ amounts to build a weak equivalence of cofree coalgebras

$$F^{c}(\mathcal{B}_{E_{1}}^{\mathbb{K}}(\overline{N^{*}}K^{\bullet}(C))) \xrightarrow{\sim} F^{c}(\overline{N^{*}}\mathcal{B}_{E_{1}}^{\mathbb{K}}(K^{\bullet}(C))),$$

thus a quasi-isomorphism of chain complexes

$$\mathcal{B}_{E_1}^{\mathbb{K}}(\overline{N^*}K^{\bullet}(C)) \stackrel{\sim}{\to} \overline{N^*}\mathcal{B}_{E_1}^{\mathbb{K}}(K^{\bullet}(C)).$$

On the one hand, as explained before, the functors U and F^c commute with reflexive coequalizers of E_1 -algebras, in particular with the bar construction $\mathcal{B}_{E_1}^{\mathbb{K}}$, hence the equality

$$\overline{N^*}\mathcal{B}_{E_1}^{\mathbb{K}}(K^{\bullet}(C)) = \overline{N^*}K^{\bullet}(\mathcal{B}_{E_1}^{\mathbb{K}}(C)).$$

By the same argument, the coalgebra structure map $C \to F^c(C)$ of C induces a map $\mathcal{B}_{E_1}^{\mathbb{K}}(C) \to \mathcal{B}_{E_1}^{\mathbb{K}}(F^c(C)) = F^c(\mathcal{B}_{E_1}^{\mathbb{K}}(C))$, so $\mathcal{B}_{E_1}^{\mathbb{K}}(C)$ is a coalgebra and we get the canonical quasi-isomorphism

$$\overline{N^*}K^{\bullet}(\mathcal{B}_{E_1}^{\mathbb{K}}(C)) \stackrel{\sim}{\to} \overline{\Omega}\mathcal{B}_{E_1}^{\mathbb{K}}(C)$$

between the reduced conormalization and the reduced cobar construction. On the other hand, the E_1 -algebra structure of C is compatible with the quasi-isomorphism

$$\overline{N^*}K^{\bullet}(C) \stackrel{\sim}{\to} \overline{\Omega}(C)$$

so that this is actually a quasi-isomorphism of E_1 -algebras. The compatibility of the E_1 -algebra structures with this map follows, once again, from the lax monoidality of the reduced conormalization, and from the lax monoidality of the cobar construction functor of coalgebras via the Milgram map [60]. In turn, this quasi-isomorphism of E_1 -algebras induces a quasi-isomorphism

$$\mathcal{B}_{E_1}^{\mathbb{K}}(\overline{N^*}K^{\bullet}(C)) \stackrel{\sim}{\to} \mathcal{B}_{E_1}^{\mathbb{K}}(\overline{\Omega}(C)).$$

Finally, the equality

$$\mathcal{B}_{E_1}^{\mathbb{K}}(\overline{\Omega}(C)) = \overline{\Omega}(\mathcal{B}_{E_1}^{\mathbb{K}}(C))$$

holds true for any E_1 -algebra in coalgebras C. Indeed, on the one hand, the cobar functor $\overline{\Omega}$ is a left adjoint, so it commutes with colimits of chain complexes. On the other hand, the bar construction $\mathcal{B}_{E_1}^{\mathbb{K}}$ is a reflexive coequalizer of E_1 -algebras, and such reflexive coequalizers are created in chain complexes.

This provides consequently the desired quasi-isomorphism between $\mathcal{B}_{E_1}^{\mathbb{K}}(\overline{N^*}K^{\bullet}(C))$ and $\overline{N^*}\mathcal{B}_{E_1}^{\mathbb{K}}(K^{\bullet}(C))$.

5. Bialgebras versus E_2 -algebras

5.1. (Co)units and (co)augmentations. We refer the reader to [2] for a detailed exposition about associative algebras and coassociative coalgebras in the pointed setting. A pointed complex is a complex X equipped with two maps $e: \mathbb{K} \to X$ and $\epsilon: X \to \mathbb{K}$ such that $\epsilon \circ e = Id_{\mathbb{K}}$. Pointed complexes form a category $Ch_{\mathbb{K}}^{pt}$, with morphisms defined by chain morphisms commuting with these maps. The two functors

$$(-)_-:X\in Ch^{pt}_{\mathbb{K}}\mapsto X_-=ker(\epsilon)\in Ch_{\mathbb{K}}$$

and

$$(-)_+:X\in Ch_{\mathbb{K}}\mapsto X_+=X\oplus \mathbb{K}\in Ch_{\mathbb{K}}^{pt}$$

form an equivalence of categories

$$(-)_+: Ch_{\mathbb{K}} \rightleftarrows Ch_{\mathbb{K}}^{pt}: (-)_-.$$

This equivalence induces on $Ch_{\mathbb{K}}^{pt}$ the structure of a cofibrantly generated symmetric monoidal model category tensored over $Ch_{\mathbb{K}}$, with the following features:

ullet the coproduct (and also product) of two pointed complexes X and Y is given by

$$X \vee Y = (X_- \oplus Y_-)_+;$$

 \bullet the tensor product of two pointed complexes X and Y is given by

$$X \wedge Y = (X_- \otimes Y_-)_+;$$

• the internal hom is given by

$$Hom_{nt}(X,Y) = Hom(X_{-},Y_{-})_{+};$$

ullet the external tensor product of a complex X with a pointed complex Y is given by

$$X \otimes_e Y = X_+ \wedge Y (= (X \otimes Y_-)_+);$$

• the external hom is given by

$$Hom_{dq}(X,Y) = Hom(X_-,Y_-),$$

because of the sequence of equalities

$$Mor_{pt}(X \otimes_e Y, Z) = Mor_{pt}((X \otimes Y_-)_+, Z) \cong Mor(X \otimes Y_-, Z_-)$$

 $\cong Mor(X, Hom(Y_-, Z_-));$

 weak equivalences are quasi-isomorphisms and fibrations are degreewise surjections.

Now let P be an augmented dg operad (i.e with an operad morphism $P \to I$) presented by $P = \mathcal{F}(\mathbb{K}\mu \oplus E)/(R)$, where μ is a generator of arity 2 (a product). We suppose moreover that P is equipped with an operad morphism $P \to Com$, so that \mathbb{K} with its usual commutative associative product is a P-algebra. This includes associative, commutative and Lie algebras (here E is the zero Σ -object), as well as n-Poisson algebras (here E is defined by a generator of degree 1-n in arity 2 giving the shifted Poisson bracket), and E_n -algebras via the sequence of operad morphisms

$$E_1 \hookrightarrow \dots \hookrightarrow E_n \hookrightarrow \dots \hookrightarrow E_\infty \xrightarrow{\sim} Com.$$

Definition 5.1. (1) A P-algebra A is unitary if it is equipped with a chain morphism $e: \mathbb{K} \to A$ satisfying the unitarity relations

$$\mu(e(1_{\mathbb{K}}), -) = \mu(-, e(1_{\mathbb{K}})) = id_A.$$

Unitary P-algebras are uP-algebras, where uP is obtained from P by adding a generator of arity 1 for the unit and adding the unitarity relations to R.

- (2) A P-algebra A is augmented if it is equipped with a P-algebra morphism $\epsilon:A\to\mathbb{K}$, where \mathbb{K} is equipped with the P-algebra structure induced by the morphism $P\to Com$ and the standard commutative algebra structure of \mathbb{K} . Such an algebra A is connected if its augmentation ideal is 0-connected, equivalently if the degree zero part of A is isomorphic to \mathbb{K} .
- (3) A P-algebra A is pointed if it is equipped with a unit e and an augmentation ϵ satisfying

$$\epsilon \circ e = id_{\mathbb{K}}.$$

- (4) A strictly unitary P_{∞} -algebra, or suP_{∞} -algebra, is a P_{∞} -algebra (where P_{∞} is a cofibrant resolution of P) equipped with a strict unit with respect to the arity 2 generator of P_{∞} inducing the product μ in homology. Similarly, we get the notions of augmented P_{∞} -algebra and pointed P_{∞} -algebra.
- (5) Dually, one can define counitary, coaugmented and pointed P-algebras and P_{∞} -algebras.

Remark 5.2. An augmented uP-algebra is an uP-algebra A with a morphism of uP-algebras $\epsilon:A\to\mathbb{K}$. The compatibility of this morphism with the unit $e:\mathbb{K}\to A$ of A and the unit $id_{\mathbb{K}}$ of the uP-algebra \mathbb{K} is exactly the relation $\epsilon\circ e=id_{\mathbb{K}}$, so augmented uP-algebras are exactly pointed P-algebras. Dually, coaugmented uP-coalgebras are exactly pointed P-coalgebras. The same identification holds for suP_{∞} -algebras and suP_{∞} -coalgebras.

Notations. To conclude this section of definitions, let us fix some notations for the next part. When writing our various categories of algebras and coalgebras, we use the superscripts *aug* for augmented algebras, *coaug* for coaugmented coalgebras,

con for connected (co)algebras, 0-con for 0-connected (co)algebras, and pt for pointed (co)algebras.

5.2. **Proofs of Theorem 0.1(2) and Corollary 0.2.** Part (2) of Theorem 0.1 follows from the following lemma:

Lemma 5.3. (1) There is an equivalence of ∞ -categories

$$(-)_+: E_1 - Cog^{conil}(dgCog^{conil}) \rightleftharpoons E_1 - Cog^{conil,pt}(dgCog^{conil}): (-)_-.$$

(2) There is an equivalence of ∞ -categories

$$(-)_+: E_1 - Alg^{0-con}(dgCog^{conil}) \rightleftharpoons E_1 - Alg^{aug,con}(dgCog^{conil}): (-)_-.$$

Proof. Recall that the functors $(-)_+$ and $(-)_-$ defines equivalences of categories

$$(-)_+: E_1 - Alg \rightleftharpoons E_1 - Alg^{pt}: (-)_-$$

between non-unitary A_{∞} -algebras and augmented (strictly) unitary A_{∞} -algebras (see for instance [45]) and dually

$$(-)_+: E_1 - Cog^{conil} \rightleftarrows E_1 - Alg^{conil,pt}: (-)_-$$

between conilpotent non-counitary A_{∞} -coalgebras and conilpotent coaugmented (strictly) counitary A_{∞} -algebras. The functor $(-)_+$ is defined by a direct sum and the functor $(-)_-$ is defined by a cokernel (cokernel of the coaugmentation). Since colimits in conilpotent dg coalgebras are created in chain complexes, these functors lift to the equivalence of categories of part (2). The same argument applies for part (1), combined with the fact that this equivalence restricts to an equivalence between 0-connected algebras and connected augmented algebras. In both cases, the base field $\mathbb K$ gives a well defined pointing because it is an associative and coassociative bialgebra. Moreover, in both cases the functors preserve quasi-isomorphisms by definition, so we conclude by applying Lemma 2.3.

According to Lemma 5.3 and to part (1) of Theorem 0.1, the composite adjunction

$$(\mathcal{B}_{E_1}^{enh,pt}(-)_-)_+:E_1-Alg^{aug,con}(dgCog^{conil})\rightleftarrows E_1-Cog^{conil,pt}(dgCog^{conil}):-(\Omega_{E_1}^{enh,pt}(-)_-)_+$$

defines an equivalence of ∞ -categories.

We now build explicitly the fully faithful ∞ -functor of Corollary 0.2:

Theorem 5.4. There is a fully faithful ∞ -functor

$$\tilde{\Omega}: E_1 - Alg^{aug,con}(dgCog^{conil}) \hookrightarrow E_2 - Alg^{aug}$$

from pointed homotopy associative conilpotent dg bialgebras to augmented dg E_2 -algebras.

Proof. The equivalence with the ∞ -category of nilpotent E_2 -algebras goes through the following sequence of ∞ -functors:

$$E_{1} - Alg^{aug,con}(dgCog^{conil}) \xrightarrow{(\mathcal{B}_{E_{1}}^{enh}(-)_{-})_{+}} E_{1} - Cog^{conil,pt}(dgCog^{conil})$$

$$= suE_{1} \otimes uAss - Cog^{conil,coaug}$$

$$\hookrightarrow uE_{1} \otimes uAss - Cog^{conil,coaug}$$

$$\xrightarrow{\widetilde{\phi}^{*}} uE_{2} - Cog^{conil,coaug} = E_{2} - Cog^{conil,pt}$$

$$\xrightarrow{\widetilde{\phi}^{*}} E_{2} - Cog^{conil}$$

$$\xrightarrow{\widetilde{\phi}^{*}} E_{2} - Alg^{aug}.$$

The equivalence of the first line is the functor $(\mathcal{B}_{E_1}^{enh}(-)_-)_+$ given by Theorem 0.1(2). The second line holds by definition of the Boardman-Vogt tensor product \otimes of operads. Indeed, recall that we have

$$E_1 - Cog^{conil,pt}(dgCog^{conil}) = suE_1 - Cog^{conil,coaug}(dgCog^{conil}).$$

The operad $suE_1 \otimes uAss$ encodes suE_1 -coalgebras in uAss-coalgebras, and the pointing of any coalgebra $A \in E_1 - Cog^{conil,pt}(dgCog^{conil})$ fixes a pointing on A as an object of $dgCog^{conil}$ as well, hence the equality

$$suE_1 - Cog^{conil}(uAss - Cog^{conil}) = suE_1 \otimes uAss - Cog^{conil}$$

restricts to

$$E_1 - Cog^{conil,pt}(dgCog^{conil}) = suE_1 - Cog^{conil,coaug}(dgCog^{conil})$$

= $suE_1 \otimes uAss - Cog^{conil,coaug}$.

The third line is the fully faithful functor induced by the operad morphism

$$uE_1 \otimes uAss \rightarrow suE_1 \otimes uAss$$
,

where $suE_1 \otimes uAss$ parametrizes coalgebras whose E_1 -structure is strictly counitary, and $uE_1 \otimes uAss$ parametrizes coalgebras whose E_1 -structure is counitary up to homotopy in order to use a cofibrant model of uE_1 (noted E_1 in [18]). At the level of the corresponding categories of coalgebras, this embeds coalgebras with a strict counit into coalgebras with a homotopy counit. This is a surjection of operads, which induces consequently a fully faithful ∞ -functor by Corollary 2.5.

The functor ϕ^* is defined as follows. First, the precomposition of any $uE_1 \otimes uAss$ -algebra structure map with the operad morphism

$$\phi: uE_2 \xrightarrow{\sim} uE_1 \otimes uE_1 \xrightarrow{\sim} uE_1 \otimes uAss$$

gives a left Quillen functor

$$\phi^*: uE_1 \otimes uAss - Coq^{conil} \rightarrow uE_2 - Coq^{conil},$$

where the model structures are given by Theorem 1.16 (weak equivalences and cofibrations are quasi-isomorphisms and degreewise injections). According to the results of [54] and [18] this is a composite of weak equivalences of Σ -cofibrant operads, so this functor is actually an equivalence of ∞ -categories. According to [18, Proposition 3.7], we have $uE_1 \otimes uAss - Cog^{conil} = u(E_1 \otimes Ass) - Cog^{conil}$ (the units of two coalgebra structures related by an interchange law coincide). So this is actually an equivalence

$$\phi^*: u(E_1 \otimes Ass) - Cog^{conil} \stackrel{\sim}{\to} uE_2 - Cog^{conil}.$$

This equivalence induces, in turn, an equivalence of the slice ∞ -categories

$$\phi^* : \mathbb{K} \setminus u(E_1 \otimes Ass) - Cog^{conil} \xrightarrow{\sim} \phi^*(\mathbb{K}) \setminus uE_2 - Cog^{conil}.$$

By definition, these slice categories are respectively $\mathbb{K} \setminus u(E_1 \otimes Ass) - Cog^{conil} = uE_1 \otimes uAss - Cog^{conil,coaug}$ and $\phi^*(\mathbb{K}) \setminus uE_2 - Cog^{conil} = uE_2 - Cog^{conil,coaug}$.

The functor $(-)_{-}$ defines an equivalence of categories between pointed E_{2} coalgebras and E_2 -coalgebras which preserves quasi-isomorphisms, hence an equivalence of ∞ -categories by Lemma 2.3.

Finally, the functor $Cobar^{(2)}$ is the twice iterated cobar construction on conilpotent E_2 -coalgebras, which gives the Koszul duality functor associated to E_2 and forms a Quillen equivalence when restricted to conilpotent E_2 -coalgebras and augmented E_2 -algebras, hence an equivalence of ∞ -categories.

The functor $\tilde{\Omega}$ is thus defined, for every pointed homotopy associative conilpotent dg bialgebra B, by

$$\begin{array}{lcl} \tilde{\Omega}(B) & = & Cobar^{(2)}(\phi^*(\mathcal{B}^{enh}_{E_1}(B_-)_+)_-) \\ & = & Cobar^{(2)}\phi^*\mathcal{B}^{enh}_{E_1}(B_-). \end{array}$$

Example 5.5. Our main working example in the last sections of the paper is the symmetric bialgebra Sym(V) over a chain complex V. It is actually a bicomplex, with a homological grading induced by the one of V and a weight grading defined by the symmetric powers. In the remaining part we will consider the total complex associated to this bicomplex, thus we will consider Sym(V) as a dg bialgebra with degree given by the total degree (sum of the homological degree and the weight), and with differential induced by the differential of V. This differential does not change the weight but decreases the homological degree by 1, so it decreases the total degree by 1. There is a decomposition

$$Sym(V) = \mathbb{K} \oplus Sym^{\geq 1}(V)$$

where $Sym^{\geq 1}(V)$ is the part of weight greater or equal to 1, hence the part of total degree greater or equal to 1. The base field $\mathbb{K} = Sym^0(V)$ is the part of total degree 0 in Sym(V), making Sym(V) a pointed conilpotent bialgebra. Hence it makes sense to study $\tilde{\Omega}(Sym(V)) = Cobar^{(2)} \phi^* \mathcal{B}_{E_1}^{enh}(Sym^{\geq 1}(V)).$

5.3. **Proof of Theorem 0.3.** Let us recall the following consequence of Section 2:

Theorem 5.6. Let P_{∞} and Q_{∞} be two properads and X and Y be two complexes. Let $\varphi: P_{\infty} \to End_X$ be a P_{∞} -algebra structure on X and $\psi: P_{\infty} \to End_Y$ be a Q_{∞} -algebra structure on Y. Let us suppose that there is a homotopy equivalence of formal moduli problems.

$$\underline{P_{\infty}\{X\}}^{\varphi} \simeq \underline{Q_{\infty}\{Y\}}^{\psi}.$$

 $\underline{P_\infty\{X\}}^\varphi\simeq\underline{Q_\infty\{Y\}}^\psi.$ Then there exists a zigzag of quasi-isomorphisms

$$Der_{\varphi}(P_{\infty}, End_X) \underbrace{\overset{\sim}{\longleftarrow} \bullet \overset{\sim}{\longrightarrow}}_{L_{\infty}} Der_{\psi}(Q_{\infty}, End_Y)$$

of L_{∞} -algebras.

Proof. By Lurie's equivalence generalized to L_{∞} -algebras (see Section 2.1), a homotopy equivalence of formal moduli problems induces a zigzag of quasi-isomorphisms of their tangent L_{∞} -algebras. In the case of formal moduli problems of algebraic structures, the tangent L_{∞} -algebra can be identified with a derivation complex (see Section 2.2).

To get Theorem 0.3(1), one first applies Theorem 2.16 to the functor $\tilde{\Omega}$. In the sequence of ∞ -functors used to define $\tilde{\Omega}$, there are equivalences plus a functor induced by precomposition with a surjection of operads. This functor is a fully faithful and conservative ∞ -functor, so $\tilde{\Omega}$ gives a fully faithful and conservative ∞ -functor. Moreover, the functor induced by precomposition with an operad morphism extends functorially by definition to algebras in A-modules, and still gives a fully faithful and conservative ∞ -functor (it does not affect the underlying A-module structure of an algebra, and weak equivalences of algebras in A-modules are still the quasi-isomorphisms). The construction of the equivalences in Theorem 5.4 extends readily to algebras in A-modules as well. Indeed, the properties of chain complexes needed to build $\mathcal{B}_{E_1}^{enh}$ and $Cobar^{(2)}$ are the same in A-modules, which form a cofibrantly generated symmetric monoidal dg category with weak equivalences defined by quasi-isomorphisms, and these two functors are by construction compatible with linear extensions of the form $-\otimes_A B$. Second, we have a fully faithful and conservative ∞ -functor

$$E_1 - Alg^{aug,con}(dgCog^{conil}) \hookrightarrow Bialg_{\infty} - Alg$$

induced by the projection $Bialg_{\infty} \to (E_1, Ass^{\vee})$, where (E_1, Ass^{\vee}) is the properad encoding E_1 -algebras in dg coalgebras. So the associated tangent formal moduli problems are equivalent, which concludes the proof, since the formal moduli problem of deformations of $B \in E_1 - Alg^{aug,con}(dgCog^{conil})$ as a homotopy bialgebra is exactly $Bialg_{\infty}\{B\}^{\varphi}$.

The proof of part (2) of Theorem 0.3 is completely similar for the following reasons. The functor $(-)^+$ takes quasi-free properads satisfying the conditions of [57, Corollary 40] to quasi-free properads satisfying the same conditions. By [57, Corollary 40], such properads are cofibrant and by [57, Theorem 42], every properad admits a cofibrant resolution of this form. Since our formal moduli problems are homotopy invariant under the choice of a resolution, we can always choose a resolution of this form, and [57, Proposition 43] ensures that the corresponding deformation complexes are also invariant, up to quasi-isomorphism, under the choice of a resolution. So that all we have to do is to replace props P_{∞} by P_{∞}^+ in the constructions above to transpose our results from P_{∞} -algebras to P_{∞}^+ -algebras. We consequently get, for any $Bialg_{\infty}^+$ -algebra φ^+ : $Bialg_{\infty}^+ \to End_B$ and the corresponding E_2^+ -algebra structure ψ^+ : $E_2^+ \to End_{\Omega B}$, a zigzag of quasi-isomorphisms of L_{∞} -algebras

$$g_{Bialg_{\infty}^{+},B}^{\varphi^{+}} \stackrel{\sim}{\leftarrow} \bullet \stackrel{\sim}{\rightarrow} g_{E_{2}^{+},\tilde{\Omega}B}^{\psi^{+}}.$$

The left-hand complex is identified with the Gerstenhaber-Schack complex $C_{GS}^*(B, B)$ (see [59]). In other words we have a zigzag of quasi-isomorphisms

(5.1)
$$C_{GS}^*(B,B) \stackrel{\sim}{\leftarrow} \bullet \stackrel{\sim}{\rightarrow} g_{E_7^+,\tilde{\Omega}B}^{\psi^+},$$

so it remains to identify the right-hand complex with the E_2 -Hochschild complex $CH_{E_2}^{(\bullet>0)}(\tilde{\Omega}B,\tilde{\Omega}B)$. This is precisely given by Lemma 3.6.

Remark 5.7. As explained in [59], the reason why we need the "plus" construction to recover the Gerstenhaber-Schack complex is that, one the one hand

$$C^*_{GS}(B,B) \cong \prod_{m,n \geq 1} Hom_{dg}(B^{\otimes m},B^{\otimes n})[2-m-n],$$

but on the other hand

$$g_{Bialg_{\infty},B}^{\varphi}\cong\prod_{m,n\geq 1,m+n\geq 3}Hom_{dg}(B^{\otimes m},B^{\otimes n})[2-m-n]$$

is an L_{∞} -algebra with differential also given by the Gerstenhaber-Schack differential, but without the term $Hom_{dq}(B, B)$.

6. Identification of Deformation complexes with higher Hochschild, Gerstenhaber-Schack and Tamarkin complexes of $Pois_n$ -algebras

In this section we identify the deformation complex $g_{E_2^+,\tilde{\Omega}B}^{\psi^+}$ and prove Corollary 0.7. More generally we identify the underlying L_{∞} -structures of several complexes related to the deformation theory of $Pois_n$, E_n and bialgebra structures.

6.1. Tamarkin deformation complexes of $Pois_n$ -algebras. We have already mentioned the higher Hochschild complexes controlling the deformation theory of E_2 -algebras in Section 1.3 and their identification with deformation complexes and tangent complexes (ee Corollary 3.6). We now introduce Tamarkin deformation complexes of a $Pois_n$ -algebra [74] and prove that these complexes do control deformations of (dg-) $Pois_n$ -algebras. Relying on the formality of E_n -operads, we will use these complexes to simplify the computations of the deformation complex of symmetric bialgebras in Section 7. We only need the case n = 2, but the proofs are just as easy for a general n so we do it in this generality.

We denote by $Pois_n$ the operad of $Pois_n$ -algebras and $uPois_n$ the operad of unital $Pois_n$ -algebras.

Let A be a dg $Pois_n$ -algebra, with structure morphism $\psi : Pois_n \to End_A$. We denote by $CH_{Pois_n}^*(A, A)$ its $Pois_n$ -Hochschild cochain complex, also referred to as its $Pois_n$ -deformation complex as defined by Tamarkin [74] and Kontsevich [47]. Following Calaque-Willwacher [7], we note that this complex is given by the suspension

(6.1)
$$CH_{Pois_n}^*(A, A) := Hom_{\Sigma}(uPois_n^*\{n\}, End_A)[-n]$$

of the underlying chain complex of the convolution Lie algebra. Here $(-)^*$ is the linear dual and $\{n\}$ is the operadic n-iterated suspension. The inclusion of $Pois_n$ in $uPois_n$ induces a splitting (as a graded space)

$$(6.2) CH_{Pois_n}^*(A, A) \cong A \oplus Hom_{\Sigma}(Pois_n^*\{n\}, End_A)[-n]$$

and also gives rise to the truncated deformation complex

(6.3)
$$CH_{Pois_n}^{(\bullet>0)}(A,A) = Hom_{\Sigma}(Pois_n^*\{n\}, End_A)[-n]$$

obtained by deleting the "unit part" A, which is more relevant to deformations of $Pois_n$ -algebras², see Lemma 6.3. Note that both complexes are naturally bigraded with respect to the internal grading of A and the "operadic" grading coming from

²as opposed to deformation of categories of modules

 $uPois_n^*$. The notation $CH_{Pois_n}^{(\bullet>0)}(A,A)$ is there to suggest that we are taking the subcomplex with positive weight with respect to the operadic grading.

The suspensions $CH^*_{Pois_n}(A,A)[n]$ and $CH^{(\bullet>0)}_{Pois_n}(A,A)[n]$ have canonical L_{∞} -structures since they are convolution algebras, and $CH^{(\bullet>0)}_{Pois_n}(A,A)[n]$ is canonically a sub L_{∞} -algebra of $CH^*_{Pois_n}(A,A)[n]$. Tamarkin [74] (see also [47, 7]) proved that the complex $CH^*_{Pois_n}(A,A)$ actually inherits a (homotopy) $Pois_{n+1}$ -algebra structure lifting this L_{∞} -structure. Further, by (6.2) we have an exact sequence of cochain complexes

$$(6.4) 0 \longrightarrow CH_{Pois_n}^{(\bullet>0)}(A,A) \longrightarrow CH_{Pois_n}^*(A,A) \longrightarrow A \longrightarrow 0$$

which yields after suspending the exact triangle

$$(6.5) A[n-1] \xrightarrow{\partial_{Pois_n}[n-1]} CH_{Pois_n}^{(\bullet>0)}(A,A)[n] \longrightarrow CH_{Pois_n}^*(A,A)[n].$$

Remark 6.1. The map $\partial_{Pois_n}: A \subset CH^*_{Pois_n}(A,A) \to CH^{(\bullet>0)}_{Pois_n}(A,A)$ is the part of the differential in the cochain complex $CH^*_{Pois_n}(A,A) = A \oplus CH^{(\bullet>0)}_{Pois_n}(A,A)$ which comes from the operadic structure. That is $\partial_{Pois_n}(x) \in Hom(A,A)$ is the map $a \mapsto \pm [x,a]$ where the bracket is the bracket of the $Pois_n$ -algebra. The Jacobi identity for the Lie algebra A[n-1] implies that the sequence (6.5) is a sequence of L_{∞} -algebras.

Remark 6.2. The operad $Pois_n$ is denoted e_n in [7, 74] and the complex $CH^*_{Pois_n}(A, A)$ is simply denoted def(A) in Tamarkin [74]. We prefer to use the notations we have introduced by analogy with (operadic) Hochschild complexes.

The next Lemma compares the L_{∞} -algebra structure of the truncated $Pois_n$ Hochschild complex and the one associated to the derived algebraic group of homotopy automorphisms of a $Pois_n$ -algebra:

Lemma 6.3. Let A be a dg $Pois_n$ -algebra with structure $map \ \psi : Pois_n \to End_A$. There is an equality of dg Lie algebras

$$g_{Pois_n^+,A}^{\psi^+} = CH_{Pois_n}^{(\bullet>0)}(A,A)$$

where the right hand side is the truncated cochain complex of a $Pois_n$ -algebra defined by Tamarkin as above.

Proof. According to the definition of the plus construction $(-)^+$ given in Section 3, we have

$$Pois_{n\infty}^{+} = \Omega(Pois_{n}^{*}\{n\})^{+} = (\mathcal{F}(\overline{Pois_{n}^{*}\{n+1\}}^{+}), \partial^{+})$$

where $Pois_{n\infty}$ is the minimal model of $Pois_n$, $(-)^*$ is the linear dual, $\{n\}$ is the operadic *n*-iterated suspension, Ω is the operadic cobar construction and $\overline{-}$ is the coaugmentation ideal of a coaugmented cooperad. Recall that the collection of generators $\overline{Pois_n^*\{n+1\}}^+$ is given by

$$\overline{Pois_n^*\{n+1\}}^+(1) = \overline{Pois_n^*\{n+1\}}(1) \oplus \mathbb{K}[1] = \overline{Pois_n^*\{n+1\}}(1) \oplus \mathbb{K}d$$

where d is a generator of degree 1 and

$$\overline{Pois_n^*\{n+1\}}^+(r) = \overline{Pois_n^*\{n+1\}}(r)$$

for r > 1. The restriction of the differential ∂^+ on the generators decomposes into $\partial^+ = \partial + \delta$ where ∂ is the differential of the minimal model and δ is defined by

 $\delta(d)=d\otimes d$ and zero when evaluated on the other generators (note that, by the Koszul sign rule and for degree reasons, we have $\delta^2(d)=0$ so we get a differential indeed). Now let $\psi^+: Pois_{n\infty}^+ \to End_A$ be the operad morphism induced by ψ , thus a Maurer-Cartan element of the convolution graded Lie algebra $g_{Pois_n^+,A}$. We twist this Lie algebra by ψ to get a dg Lie algebra $g_{Pois_n^+,A}^{\psi^+}$ with the same Lie bracket and whose differential is defined by

$$\pm (d_A)_* + [\psi, -]$$

where $(-)_*$ denotes the post-composition, d_A is the differential on End_A induced by the differential of A, the \pm sign is defined according to the Koszul sign rule and [-,-] is the convolution Lie bracket. Note here that the Koszul dual cooperad has no internal differential. We refer the reader to [51, Chapter 12] for more details about such convolution Lie algebras. Now let us point out that

$$\overline{Pois_{n}^{*}\{n+1\}}^{+}(1) = \overline{Pois_{n}^{*}\{n+1\}}(1) \oplus \mathbb{K}[1] = (\overline{Pois_{n}^{*}\{n\}}(1) \oplus \mathbb{K})[1],$$

which implies that

$$g_{Pois_n^+A}^{\psi^+} = Hom_{\Sigma}(\overline{Pois_n^*\{n\}} \oplus I, End_A)^{\psi} = Conv(Pois_n^*\{n\}, End_A)$$

where $Conv(Pois_n^*\{n\}, End_A)$ is the convolution Lie algebra of [7, Section 2.2]. This is an equality of dg Lie algebras, because the convolution bracket is defined by the action of the infinitesimal cooperadic coproduct on the coaugmentation ideal, so is the same on both sides.

Remark 6.4. Lemma 6.3 together with Theorem 3.5 implies that the truncated Tamarkin deformation complex $CH^{(\bullet>0)}_{Pois_n}(A,A)$ controls deformations of A into dg $Pois_n$ -algebras, in other words is the tangent Lie algebra of the derived algebraic group $\underbrace{haut_{Pois_n\infty}}(A)$, where $Pois_{n\infty}$ is a cofibrant resolution of $Pois_n$.

The proof of Lemma 6.3 also shows that the deformation complex $g_{Pois_n,A}^{\psi}$ of the formal moduli problem $\underline{Pois_{n_{\infty}}\{A\}}^{\psi}$ is given by the L_{∞} -algebra $CH_{Pois_n}^{(\bullet>1)}(A,A)[n]$, which is the kernel

$$(6.6) CH_{Pois_n}^{(\bullet>1)}(A,A)[n] := \ker\left(CH_{Pois_n}^{(\bullet>0)}(A,A)[n] \twoheadrightarrow Hom(A,A)[n]\right)$$

and is thus a even further truncation of $CH^*_{Pois_n}(A,A)$. The situation is thus similar to what happens in deformation theory of associative algebras.

One can also wonder which deformation problem controls the full complex $CH^*_{Pois_n}(A,A)$. In view of Theorem 6.7 below and classical results on deformation theory of E_{n^-} algebras ([46, 62, 20]), we can conjecture that $CH^*_{Pois_n}(A,A)$ shall control deformations of categories of modules over $Pois_n$ -algebras into $E_{|n-1|}$ -monoidal decategories, with some shift on the linear enrichment of the category when $n \leq 1$ according to the red shift trick [44, 77, 76].

We now compare the deformation complexes of $Pois_n$ -algebras with those of E_n -algebras. Let us fix a formality morphism $\varphi: E_n \stackrel{\sim}{\to} Pois_n \ (n \geq 2)$. This allows to see any dg $Pois_n$ -algebra as an E_n -algebra.

Lemma 6.5. Let A be a dg Pois_n-algebra with structure map $\psi : Pois_n \to End_A$. Then, there is a quasi-isomorphism of dg Lie algebras

$$CH_{Pois_n}^{(\bullet>0)}(A,A)[n] = g_{Pois_n^+,A}^{\psi^+} \stackrel{\sim}{\to} g_{E_n^+,A}^{(\psi\circ\varphi)^+}$$

where the left hand side is the truncated cohomology complex (6.3) of a Pois_n-algebra defined by Tamarkin [74].

Proof. The formality quasi-isomorphism φ induces a quasi-isomorphism of cofibrant resolutions

$$\varphi_{\infty}: E_{n\infty} \stackrel{\sim}{\to} Pois_{n\infty}$$

fitting in a commutative diagram

$$E_{n\infty} = \Omega(E_n^i) \xrightarrow{\sim} Pois_{n\infty} = \Omega(Pois_n^i)$$

$$\downarrow \sim \qquad \qquad \sim \qquad \qquad \wedge \Omega(\pi)$$

$$\Omega \mathcal{B}(E_n) \xrightarrow{\sim} \Omega \mathcal{B}(\varphi) \longrightarrow \Omega \mathcal{B}(Pois_n)$$

where \mathcal{B} is the operadic bar construction, Ω is the operadic cobar construction, $(-)^i$ is the Koszul dual cooperad given by $H^0\mathcal{B}(-)$ (see [51, Chapter 7]), i is the inclusion and $\pi: \mathcal{B}(Pois_n) \to H^0\mathcal{B}(Pois_n)$ is the projection. This construction implies that φ_{∞} is a tangent quasi-isomorphism of dg operads in the sense of [57, Theorem 7], that is, the composite map

$$E_n^{\mathsf{i}}\{1\} \hookrightarrow E_{n\infty} \xrightarrow{\sim} Pois_{n\infty} \twoheadrightarrow Pois_n^{\mathsf{i}}\{1\}$$

is a quasi-isomorphism. Given that, according to Section 1.2.2, the plus construction $(-)^+$ preserves quasi-isomorphisms of dg props, this implies that the construction above works with the plus construction as well and induces a tangent quasi-isomorphism

$$\varphi_{\infty}^+: E_{n\infty}^+ \stackrel{\sim}{\to} Pois_{n\infty}^+,$$

which by [57, Theorem 7] induces the desired quasi-isomorphism of dg Lie algebras

$$(\varphi_{\infty}^+)^*: g_{Pois_n^+, A}^{\psi^+} \xrightarrow{\sim} g_{E_n^+, A}^{(\psi \circ \varphi)^+}.$$

Corollary 6.6. Let A be a $Pois_n$ -algebra with structure map $\psi : Pois_n \to End_A$. The formality map φ induces an equivalence of formal moduli problems

$$\varphi_{\infty}^{*}:\underline{Pois_{n\,\infty}^{+}}^{\psi^{+}}\{A\}\xrightarrow{\simeq}\underline{E_{n\,\infty}^{+}}\{\varphi^{*}A\}^{(\psi\circ\varphi)^{+}}$$

(where φ^*A is the E_n -algebra obtained from the $Pois_n$ -algebra A by restriction of structures along φ).

Further, Lemma (6.5) and Corollary (3.6) combine to give a quasi-isomorphism of L_{∞} -algebras

(6.7)
$$CH_{Pois_n}^{(\bullet>0)}(A,A)[n] \xrightarrow{\simeq} T_A$$

when A is dg- $Pois_n$ -algebra. Here T_A is the tangent complex of A viewed as an E_n -algebra using the chosen formality φ (so strictly speaking it shall be noted T_{φ^*A}).

Recall now that if A is a $Pois_n$ -algebra, then A[n-1] is a dg Lie algebra, in particular an L_{∞} -algebra. We relate its deformation complexes (the full and the truncated) as a $Pois_n$ -algebra with its deformation complexes as an E_n -algebra:

Theorem 6.7. Let A be a Pois_n-algebra $(n \ge 2)$.

(1) The sequence (6.5) fits into a morphism of fiber sequences of L_{∞} -algebras

$$A[n-1] \xrightarrow{} T_A \xrightarrow{} CH_{E_n}^*(A,A)[n]$$

$$\cong \bigwedge^{\simeq} \qquad \qquad \bigwedge^{\simeq} \qquad \qquad \bigwedge^{\simeq} \qquad \qquad \qquad A[n-1] \xrightarrow{} CH_{Pois_n}^{(\bullet>0)}(A,A)[n] \xrightarrow{} CH_{Pois_n}^*(A,A)[n]$$

whose vertical arrows are L_{∞} -quasi-isomorphisms and the middle vertical arrow is (6.7).

(2) The L_{∞} -structure on $CH_{Pois_n}^{(\bullet>0)}(A,A)[n]$ and T_A is the one controlling the derived algebraic group $\underline{haut}_{Pois_n\infty}(A)$ of homotopy automorphisms of A.

Proof. We saw that the equivalence $\varphi_{\infty}: E_{n\infty} \xrightarrow{\sim} Pois_{n\infty}$ induced by the formality morphism φ induces the weak-equivalence $\varphi_{\infty}^+: E_{n\infty}^+ \xrightarrow{\sim} Pois_{n\infty}^+$. Adding a generator for strict units gives us also a weak-equivalence $su\varphi_{\infty}^+: suE_{n\infty}^+ \xrightarrow{\sim} suPois_{n\infty}^+$. Let $\psi_{\infty}: Pois_{n\infty} \to End_A$ be a $Pois_n$ -algebra structure on A (seen as a homotopy $Pois_n$ -algebra structure), and $\psi_{\infty}^+: Pois_{n\infty}^+ \to Pois_{n\infty} \xrightarrow{\psi_{\infty}} End_A$ be the map obtained by first sending the generator δ to 0. In the proof of Lemma 6.5, we saw that φ_{∞}^+ is a tangent quasi-isomorphism, hence inducing an L_{∞} -algebra quasi-isomorphism at the level of the associated deformation complexes. Following readily the proof of Lemma 6.5, we get that $su\varphi_{\infty}^+$ is a tangent quasi-isomorphism as well, thus

$$(\varphi_{\infty}^+)^*: g_{Pois_n,A}^{\psi_{\infty}^+} \xrightarrow{\sim} g_{E_n^+,A}^{(\psi \circ \varphi)^+}$$

and

$$(su\varphi_{\infty}^{+})^{*}:g_{suPois_{n}^{+},A}^{\psi_{\infty}^{+}}\stackrel{\sim}{\longrightarrow}g_{suE_{n}^{+},A}^{(\psi\circ\varphi)^{+}}$$

are both L_{∞} -algebra quasi-isomorphisms fitting in a commutative diagram

$$(6.8) \hspace{1cm} g_{suE_n^+,A}^{(\psi\circ\varphi)^+} \longleftrightarrow g_{E_n^+,A}^{(\psi\circ\varphi)^+} \ .$$

$$(su\varphi_\infty^+)^* \not \simeq \qquad \qquad \simeq \not (\varphi_\infty^+)^*$$

$$g_{suPois_n^+,A}^{\psi_\infty^+} \longleftrightarrow g_{Pois_n^+,A}^{\psi_\infty^+}$$

As in Remark 6.1, the homopy fibers of these inclusions are both A[n-1] equipped with its Lie structure defined by shifting the Poisson bracket of A. We thus get a quasi-isomorphism of homotopy fiber sequences

$$(6.9) g_{suE_{n}^{+},A}^{(\psi\circ\varphi)^{+}} \longleftrightarrow g_{E_{n}^{+},A}^{(\psi\circ\varphi)^{+}} \longleftrightarrow A[n-1] .$$

$$(su\varphi_{\infty}^{+})^{*} \stackrel{\wedge}{\simeq} \simeq \stackrel{\wedge}{\simeq} (\varphi_{\infty}^{+})^{*} \qquad \stackrel{\wedge}{\sim}$$

$$g_{suPois_{n}^{+},A}^{\psi_{\infty}^{+}} \longleftrightarrow g_{Pois_{n}^{+},A}^{\psi_{\infty}^{+}} \longleftrightarrow A[n-1]$$

Following readily the proof of Lemma 6.3, we see that $g_{suPois_n,A}^{\psi_{\infty}^+} = CH_{Pois_n}^*(A,A)[n]$: the generator of arity zero in $suPois_n^+$ (defining the unit) is of degree 0, hence of degree n in $\overline{Pois_n^*\{n\}}$. In the associated convolution Lie algebra $g_{suPois_n,A}^{\psi_{\infty}^+}$, it gives

an additional term A[n] in operadic degree 0 which corresponds to the part of weight 0 in $CH^*_{Pois_n}(A,A)[n]$. The inclusion $g^{\psi^+_\infty}_{Pois_n^+,A} \hookrightarrow g^{\psi^+_\infty}_{suPois_n^+,A}$ is thus nothing but the inclusion $CH^{(\bullet>0)}_{Pois_n}(A,A)[n] \hookrightarrow CH^*_{Pois_n}(A,A)[n]$. This means that we have a quasi-isomorphism of homotopy fiber sequences

$$(6.10) g_{suE_{n}^{+},A}^{(\psi\circ\varphi)^{+}} \longleftarrow g_{E_{n}^{+},A}^{(\psi\circ\varphi)^{+}} \longleftarrow A[n-1] .$$

$$(su\varphi_{\infty}^{+})^{*} \stackrel{}{ } \simeq \qquad \simeq \stackrel{}{ } (\varphi_{\infty}^{+})^{*} \qquad \stackrel{}{ } \sim \stackrel{}{ }$$

$$CH_{Pois_{n}}^{*}(A,A)[n] \longleftarrow CH_{Pois_{n}}^{(\bullet>0)}(A,A)[n] \longleftarrow A[n-1]$$

Now we want to prove that the full Poisson complex $CH_{Pois_n}^*(A, A)$ is quasiisomorphic to the full E_n -Hochschild complex $CH_{E_n}^*(A, A)$. For this, we use the fact (proved later in Lemma 7.7) that the formality morphism

$$\varphi: E_n \stackrel{\sim}{\to} Pois_n$$

induces a natural quasi-isomorphism

$$Bar^{(n)} \circ \varphi^* \stackrel{\sim}{\to} Bar^{(n)}_{Pois_n},$$

where Bar^n is the operadic bar construction for E_n -algebras, $Bar_{Pois_n}^{(n)}$ is the one for $Pois_n$ -algebras and φ^* is the restriction of structures from $Pois_n$ -algebras to E_n -algebras. So, for a given n-Poisson algebra A, we have a quasi-isomorphism

$$Bar^{(n)}(\varphi^*A) \stackrel{\sim}{\to} Bar^{(n)}_{Pois_n}(A)$$

inducing a quasi-isomorphism

$$Hom_{dg}(Bar_{Pois_n}^{(n)}(A), A) \stackrel{\sim}{\to} Hom_{dg}(Bar^{(n)}(\varphi^*A), A)$$

between the complexes computing respectively the operadic $Pois_n$ -cohomology of A and the operadic E_n -cohomology of φ^*A (see [51, Section 12.4] for the general construction of operadic cohomology). By [51, Theorem 12.4.5], the operadic complex of an algebra over an operad is quasi-isomorphic to its André-Quillen complex so we get a quasi-isomorphism of André-Quillen complexes

$$C_{AQ}^{Pois_n}(A,A) \xrightarrow{\sim} C_{AQ}^{E_n}(\varphi^*A,\varphi^*A).$$

By definition, the André-Quillen complex $C_{AQ}^{E_n}(\varphi^*A, \varphi^*A)$ is the right derived functor, applied to φ^*A , of the hom functor $Hom_A^{E_n}(-, \varphi^*A)$ in operadic A-modules over E_n , that is, exactly $CH_{E_n}^*(\varphi^*A, \varphi^*A)$. Similarly, we have $C_{AQ}^{Pois_n}(A, A) = CH_{Pois_n}^*(A, A)$. Tensoring the map above by $\mathbb{K}[n]$, we get a quasi-isomorphism

$$CH_{Pois_n}^*(A,A)[n] \xrightarrow{\sim} CH_{E_n}^*(A,A)[n].$$

The equality $g_{suPois_n^+,A}^{\psi_{\infty}^+} = CH_{Pois_n}^*(A,A)[n]$ gives to $CH_{Pois_n}^*(A,A)[n]$ an L_{∞} -algebra structure that we can transfer to $CH_{E_n}^*(A,A)[n]$ by the quasi-isomorphism above to make it a quasi-isomorphism of L_{∞} -algebras.

We now compare this structure with the one given by the solution to the Deligne conjecture. The E_n -Hochschild cohomology has a natural (and unital) E_{n+1} -structure (see [54, 20, 36]) given by the $suE_1 \otimes suE_n$ -algebra structure³ on the

³The centralizer is canonically endowed with a structure of E_1 -algebras in E_n -algebras. By Dunn additivity theorem [54], this is equivalent to an E_{n+1} -structure.

 E_n -centralizer $\mathfrak{z}_{suE_n}(\varphi^*(A)) := \mathfrak{z}_{suE_n}(id_{\varphi^*(A)})$ of A, seen as an E_n -algebra using our chosen formality map. By [20], the upper horizontal sequence of chain complexes in claim (1) of the theorem is a natural sequence of L_∞ -algebras. Here, by the theory of formal moduli problems [52] and deformation theory of E_n -algebras [54, 20], $CH_{E_n}^*(A,A)[n]$ is endowed with the underlying L_∞ -algebra structure characterized by the commutative formal moduli problem defined by

(6.11)
$$Map_{E_{n+1}-Alg^{aug}}(\mathcal{D}^{(n+1)} \circ I^{(n+1)}(-), \mathfrak{z}_{suE_n}(\varphi^*(A)))$$

where $I^{(n+1)}: cdga^{aug} \to E_n^{aug}$ is the canonical functor sending a cdga to the E_n -algebra with trivial higher products, and $\mathcal{D}^{(n+1)}$ is the appropriate Koszul duality ∞ -functor as used in [52]. Note that $\mathcal{D}^{(n+1)}$ can be identified with the centralizer of the augmentation $\mathfrak{z}_{suE_n}(R \to \mathbb{K})$, hence with the dual of the n+1-fold bar construction, see [52, 36, 3].

Let us denote $\mathfrak{z}_{suPois_n}(A)$ the centralizer of the identity map $A \to A$ in the ∞ -category of $Pois_n$ -algebras; it is a (unital) E_1 -algebra in $Pois_n$ -algebras. Similarly as above, the formality map φ induces an equivalence of E_1 -algebras in E_n -algebras (hence of unital E_{n+1} -algebras):

(6.12)
$$\varphi^*(\mathfrak{z}_{suPois_n}(A)) \cong \mathfrak{z}_{suE_n}(\varphi^*(A)).$$

By [74] (also see [7, 36]), the $E_1 \otimes Pois_n$ -algebra structure of $\mathfrak{z}_{suPois_n}(A)$ is represented by an explicit compatible⁴ dg algebra structure on the dg $Pois_n$ -coalgebra $Bar_{Pois_n}^{(n)}(\mathfrak{z}_{suPois_n}(A)))$ whose differential encodes its canonical $Pois_n\infty$ -algebra structure⁵

We wish to rewrite the formal moduli problem (6.11). To do so, following [54, Sections 2.3 and 3.3], we note that we have an adjunction of ∞ -categories C_{CE}^* : $dgLie \rightleftharpoons dgArt_{\mathbb{K}}^{aug}: \mathcal{D}_{Lie}$, where C_{CE}^* is the ∞ -functor induced by the Chevalley-Eilenberg cochain algebra functor, and \mathcal{D}_{Lie} is an equivalence onto its essential image in dgLie (with inverse given by C_{CE}^*). Furthermore, by [3, Corollary 4.2.2], there is an equivalence

$$(6.13) \mathcal{D}^{(n+1)} \circ U_{n+1} (\mathcal{D}_{Lie}(R)) \cong C_{CE}^* (\mathcal{D}_{Lie}(R))$$

where U_{n+1} is the the E_{n+1} -envelopping algebra functor; in particular we have an adjunction $U_{n+1}: E_{n+1} - Alg^{aug} \hookrightarrow dgLie: [n]$. Since $\mathcal{D}^{(n+1)}$ is self-dual, for an artinian augmented cdga R, we get from (6.13) and [52, Theorem 4.4.5], an equivalence

$$U_{n+1}(\mathcal{D}_{Lie}(R)) \cong \mathcal{D}^{(n+1)} \circ C_{CE}^*(\mathcal{D}_{Lie}(R)).$$

 $^{^4{\}rm the}$ compatibility means that we have an $E_1\otimes Pois_{n\infty}\text{-algebra}$ structure

⁵ Note that the (non-unital) centralizer and is corresponding dg $Pois_n$ -coalgebra are denoted def(A)[-n] and $Hom^{Id}(A,A)$ in [74]

Combining all the above we get an equivalence of formal moduli problems given, for every artinian augmented $\operatorname{cdga} R$, by

$$(6.14) \quad Map_{E_{n+1}-Alg^{aug}} \left(\mathcal{D}^{(n+1)} \circ I^{(n+1)}(R), \mathfrak{z}_{suE_n}(\varphi^*(A)) \right)$$

$$\cong Map_{E_{n+1}-Alg^{aug}} \left(\mathcal{D}^{(n+1)} \circ C_{CE}^*(\mathcal{D}_{Lie}(R)), \varphi^*(\mathfrak{z}_{suPois_n}(A)) \right)$$

$$\cong Map_{E_{n+1}-Alg^{aug}} \left(U_{n+1}(\mathcal{D}_{Lie}(R)), \varphi^*(\mathfrak{z}_{suPois_n}(A)) \right)$$

$$\cong Map_{dgLie} \left(\mathcal{D}_{Lie}(R), \mathfrak{z}_{suPois_n}(A)[n] \right).$$

This identifies the L_{∞} -algebra structure underlying $CH_{E_n}^*(A,A) \cong \mathfrak{z}_{suE_n}(\varphi^*(A))$ with the one obtained by n-fold desuspension of $\mathfrak{z}_{suPois_n}(A)$. It remains to prove that this structure on the center coincides with the one on $CH_{Pois_n}^*(A,A)[n] = g_{suPois_n}^{\psi_{\infty}^+}(A)$; so far we have proved that they are quasi-isomorphic as chain complexes. Tamarkin [74, Section 5] has proved that $CH_{Pois_n}^*(A,A)[n]$ has the underlying L_{∞} -algebra structure of a certain $Pois_{n+1\infty}$ -algebra structure on $\mathfrak{z}_{suPois_n}(A) \cong CH_{Pois_n}^*(A,A)[n]$. This $Pois_{n+1\infty}$ -algebra structure is given by the differential on $Bar_{Pois_{n+1}}^{(n+1)}(\mathfrak{z}_{suPois_n}(A))$ and we use the later notation (or simply $CH_{Pois_n}^*(A,A)[n]$) to refer to this precise homotopy $Pois_{n+1}$ -algebra structure on $\mathfrak{z}_{suPois_n}(A)$, in order to distinguish it from the one given by the centralizer. Further Tamarkin showed that this structure induces a cocommutative Hopf bialgebra structure on $Bar_{Pois_n}^{(n)}(\mathfrak{z}_{suPois_n}(A))$, an additional compatible degree n Lie cobracket on its primitive part and finally an isomorphism of dg $Pois_{n+1}$ -coalgebras:

$$(6.15) Bar_{Pois_{n+1}}^{(n+1)}(\mathfrak{z}_{suPois_n}(A)) \cong C_*^{CE}(Prim(Bar_{Pois_n}^{(n)}(\mathfrak{z}_{suPois_n}(A))))$$

where C_*^{CE} is the standard Chevalley-Eilenberg commutative coalgebra functor (whose associated derived functor computes $\mathbb{K} \otimes_{U(-)}^{\mathbb{L}} \mathbb{K}$).

As above, Tamarkin L_{∞} -structure on $CH^*_{Pois_n}(A,A)[n]$ is characterized by the following equivalence of moduli functors

$$(6.16) \quad Map_{dgLie}(\mathcal{D}_{Lie}(R), CH^*_{Pois_n}(A, A)[n])$$

$$\cong Map_{Pois_{n+1}-Alg^{aug}}(\mathcal{D}^{(n+1)}_{Pois_{n+1}} \circ I^{(n+1)}(R), CH^*_{Pois_n}(A, A)))$$

$$\cong Map_{Pois_{n+1}-Cog}(I_{(n+1)}(R'), Bar^{(n+1)}_{Pois_{n+1}}(\mathfrak{z}_{suPois_n}(A)))$$

where $I_{(d)}: Com - Cog \rightarrow Pois_d - Cog$ is the canonical functor and R' is the (linear) dual of the artinian algebra R. The last equivalence follows from the barcobar equivalence and the definition of $Pois_{n+1\infty}$ -morphisms.

Recall that the centralizer $\mathfrak{z}_{suPois_n}(A)$ is canonically an E_1 -algebra in $Pois_{n\infty}$ algebras and this structure is given by a compatible algebra structure on $Bar_{Pois_n}^{(n)}(\mathfrak{z}_{suPois_n}(A))$. In particular it is canonically isomorphic as a cocommutative Hopf algebra to the
envelopping algebra of its primitive elements $Prim(Bar_{Pois_n}^{(n)}(\mathfrak{z}_{suPois_n}(A)))$. Also
if C is a dg cocommutative coalgebra, then its cobar construction $coBar^{(1)}(C)$ is
canonically a cocommutative Hopf algebra. Using moreover the natural equivalence $Bar^{(1)} \circ U(-) \cong C_*^{CE}(-)$ of dg cocommutative coalgebras and the adjunction

between $Bar^{(1)}$ and $coBar^{(1)}$, we get an equivalence of moduli problems

$$(6.17) \quad Map_{dgLie}\left(\mathcal{D}_{Lie}(R), \mathfrak{z}_{suPois_n}(A)[n]\right)$$

$$\cong Map_{Pois_{n+1}-Alg^{aug}}\left(\mathcal{D}_{Pois_{n+1}}^{(n+1)} \circ I^{(n+1)}(R), \mathfrak{z}_{suPois_n}(A)\right)$$

$$\cong Map_{E_1 \otimes Pois_n - Cog}\left(I_{(n)} \circ coBar^{(1)}(R'), Bar_{Pois_n}^{(n)}(\mathfrak{z}_{suPois_n}(A))\right)$$

$$\cong Map_{Pois_{n+1}-Cog}\left(I_{(n+1)}(R'), Bar^{(1)} \circ Bar_{Pois_n}^{(n)}(\mathfrak{z}_{suPois_n}(A))\right)$$

$$\cong Map_{Pois_{n+1}-Cog}\left(I_{(n+1)}(R'), C_*^{CE}(Prim(Bar_{Pois_n}^{(n)}(\mathfrak{z}_{suPois_n}(A))))\right).$$

Therefore the isomorphism (6.15) implies that the two moduli functors (6.16) and (6.17) are equivalent. Consequently, the two L_{∞} -algebras that they represent are equivalent as well by Lurie's equivalence theorem, which yields the right vertical equivalence of L_{∞} -algebras in Claim (1).

Moreover, by Corollary 3.6 we have an equivalence of L_{∞} -algebras $g_{E_n^+,A}^{(\psi\circ\varphi)^+} \stackrel{\sim}{\to} T_A$ and we finally get the commutative diagram of L_{∞} -algebras

Hence both lines have quasi-isomorphic fibers as well which are A[n-1] according to (6.5) and [20].

Remark 6.8. Claim (2) can be proved directly by using the equivalence $(\varphi_{\infty}^+)^*$: $\underline{Pois_{n_{\infty}}}^+\{A\} \stackrel{\sim}{\to} \underline{E_{n_{\infty}}}^+\{\varphi^*A\}$ of Corollary 6.6. Indeed, applying Theorem 3.5 gives us directly a quasi-isomorphism of L_{∞} -algebras

$$CH_{Pois}^{*>0}(A,A)[n] = g_{Pois_n^+,A}^{\psi^+} \stackrel{\sim}{\to} T_{\varphi^*A}.$$

Remark 6.9. In particular, we also get out of Theorem 6.7 that the L_{∞} -algebra $g_{Pois_n,A}^{\psi}$ controlling the moduli space $\underline{Pois_{n_{\infty}}}\{A\}$ is thus isomorphic to a sub- L_{∞} -algebra of T_A .

Remark 6.10. Let $\tilde{C}_{(d)}: Pois_d - Cog \to Com - Cog$ be the right adjoint of the canonical functor $I_{(d)}: Com - Cog \to Pois_d - Cog$. The moduli functor (6.17) encoding the L_{∞} -structure on $CH_{Pois}(A,A)[n]$ is then equivalent to

$$\begin{split} Map_{Pois_{n+1}-Alg^{aug}} \Big(\mathcal{D}_{Pois_{n+1}}^{(n+1)} \circ I^{(n+1)}(R), CH_{Pois_{n}}^{*}(A,A)) \Big) \\ &\cong Map_{Pois_{n+1}-Cog} \Big(I_{(n+1)}(R'), Bar^{(1)} \circ Bar_{Pois_{n}}^{(n)}(\mathfrak{z}_{suPois_{n}}(A)) \Big) \\ &\cong Map_{E_{1} \otimes Com-Cog} \Big(coBar^{(1)}(R'), \tilde{C}_{(n)}(Bar_{Pois_{n}}^{(n)}(\mathfrak{z}_{suPois_{n}}(A))) \Big) \\ &\cong Map_{E_{1} \otimes Com-Cog} \Big(coBar^{(1)}(R'), U(\mathfrak{z}_{suPois_{n}}(A)[n]) \Big). \end{split}$$

wher the last line is [74, Corollary 4.7]. By (6.16), this is also equivalent to

$$Map_{Pois_{n+1}-Cog}\Big(I^{(n+1)}(R'), Bar_{Pois_{n+1}}^{(n+1)}(\mathfrak{z}_{suPois_n}(A))\Big)$$


$$\cong Map_{Com-Cog}\Big(R', \tilde{C}_{(n+1)}(Bar_{Pois_{n+1}}^{(n+1)}(\mathfrak{z}_{suPois_n}(A)))\Big).$$

Let $\eta: B \to k$ be an augmented unital cdga and $\overline{B} := \ker(\eta)$ be its augmentation ideal, so that $B \cong k \oplus \overline{B}$. Both B and \overline{B} are cdgas, hence (non-unital) E_n -algebras for any n. In particular, B and \overline{B} are canonically E_n -B-modules as well as E_n - \overline{B} -modules. Let us denote $\iota: \overline{B} \to B$ the canonical inclusion.

Lemma 6.11. The canonical maps

$$CH_{E_n}^*(B,B) \overset{\iota^*}{\to} CH_{E_n}^*(\overline{B},B), \quad CH_{E_n}^*(B,\overline{B}) \overset{\iota^*}{\to} CH_{E_n}^*(\overline{B},\overline{B})$$

are quasi-isomorphisms and the following diagram



is a commutative diagram of E_{n+1} -algebras.

Proof. The E_n -Hochschild cohomology of a cdga B with value in a bimodule M can be computed using Pirashvili type higher Hochschild cochains (see [36]):

$$CH_{E_n}^*(B,M) \cong CH^{S_{\bullet}^n}(B,M) \cong Hom_B(B^{\otimes S_{\bullet}^n},M)$$

for any simplicial set model S^n_{\bullet} of the *n*-sphere. Being obtained as a cosimplicial cochain complex, the later one is naturally quasi-isomorphic to its non-degenerate cochains, which amounts to quotienting by the submodule generated by the unit (see [35]). Hence, denoting * the base-point of S^n_{\bullet} , since B is augmented, we have a quasi-isomorphism

$$q_*: Hom_B(B^{\otimes S^n_{\bullet}}, M) \stackrel{\simeq}{\to} Hom_B(B \otimes \overline{B}^{\otimes S^n_{\bullet} \setminus \{*\}}, M) \cong CH^*_{E_n}(\overline{B}, M).$$

Since $q_* \circ \iota^*$ is the identity, we obtain that ι^* is a quasi-isomorphism as well, hence the first claim by taking M = B or \overline{B} . The commutativity of the diagram follows from the same identification with higher Hochschild cochains and a straightforward computation.

Since all modules structures are induced by maps of cdgas, the same identification with higher Hochschild cochains implies that all complexes in the diagram above are canonically E_{n+1} -algebras, this structure being functorial with respect to maps of E_n -algebras, see [36]; this also follows from [20] using moduli problems.

From Lemma 6.11, we get a canonical (zigzag of) cochain maps

(6.18)
$$\iota: CH_{E_n}^*(\overline{B}, \overline{B}) \simeq CH_{E_n}^*(B, \overline{B}) \xrightarrow{\iota_*} CH_{E_n}^*(B, B).$$

Note that we endow $CH_{E_n}^*(B, \overline{B})$ with its E_{n+1} -algebra structure given by transfer along $\iota^*: CH_{E_n}^*(B, B) \stackrel{\iota^*}{\to} CH_{E_n}^*(\overline{B}, B)$.

Proposition 6.12. Assume $n \geq 2$ and B is an augmented cdga essentially of finite type. The canonical map (6.18) $\iota : CH_{E_n}^*(\overline{B}, \overline{B}) \to CH_{E_n}^*(B, B)$ is a (non-unital) E_{n+1} -algebra homomorphism. Further, we have a cofiber sequence of cochain complexes

$$CH_{E_n}^*(\overline{B}, \overline{B}) \xrightarrow{\iota} CH_{E_n}^*(B, B) \xrightarrow{\pi_*} CH_{E_n}^*(B, \mathbb{K}).$$

Proof. Taking a cofibrant resolution of B as a cdga, we may assume B = (Sym(W), d) is an augmented cofibrant cdga and each W has finite dimensional cohomology; in particular $d(W) \subset Sym^{\geq 1}(W)$ and $\overline{B} = Sym^{\geq 1}(W)$. Being cdgas, these algebras are canonically $Pois_n$ -algebras, and this structure is compatible with the induced E_n -algebra structure. By [7], [36, Corollary 6.39]⁶, we have, for any cofibrant cdga A, an equivalence of E_{n+1} -algebras

(6.19)
$$CH_{E_n}^*(A,A) \simeq \widehat{Sym}_A(Der(A,A)[-2])$$

where the shifted Lie bracket is given by the Lie bracket on derivations and the Leibniz rule. Here Der(A,A) is the cochain complex of derivations and the E_{n+1} -algebra structure on the right hand side of (6.19) is induced by formality from its canonical $Pois_{n+1}$ -structure. By Theorem 6.7, we can replace $CH_{E_n}^*(A,A)$ by $CH_{Pois_n}^*(A,A) = Hom_{\Sigma}(uPois_n^*\{n\}, End_A)[-n]$ in the above equivalence.

Since B = (Sym(W), d), we have an isomorphism of complexes:

$$\iota^* : Der(\overline{B}, B) \cong Der(B, B)$$

as well as canonical quasi-isomorphisms

$$(W)^* \otimes B \stackrel{\simeq}{\to} Der(B,B), \quad (W)^* \otimes \overline{B} \stackrel{\simeq}{\to} Der(\overline{B},\overline{B})$$

identifying the canonical map $\iota_*: Der(\overline{B}, \overline{B}) \hookrightarrow Der(B, B)$ (extending a derivation on \overline{B} to a derivation on B) with the map $id_{(W)^*} \otimes \iota$. The equivalence (6.19) is given by the map

$$\widehat{Sym}_A(Der(A,A)[-n] \subset Hom_{\Sigma}(uCom^*\{n\},End_A)[-n] \hookrightarrow Hom_{\Sigma}(uPois_n^*\{n\},End_A)[-n]$$

where the right map is induced by the canonical operad morphism $Com \to Pois_n$. Hence we have a commutative diagram (6.20)

$$\widehat{Sym}\big((W)^*\big)\otimes Sym^{\geq 1}\big(W\big)\overset{\cong}{\longrightarrow} \widehat{Sym}_B\big(Der(\overline{B},\overline{B})[-n]\big)\overset{\cong}{\longrightarrow} CH_{E_n}^*(\overline{B},\overline{B})$$

$$\downarrow \iota$$

$$\widehat{Sym}\big((W)^*\big)\otimes Sym\big(W\big)\overset{\cong}{\longrightarrow} \widehat{Sym}_B\big(Der(B,B)[-n]\big)\overset{\cong}{\longrightarrow} CH_{E_n}^*(B,B).$$

Since the left vertical map is a map of dg $Pois_{n+1}$ -algebras and the horizontal maps are equivalences of E_{n+1} -algebras, the first claim follows. The second claim follows from the fact that the left hand side is a cofiber sequence with cofiber $\widehat{Sym}((W)^*) \otimes k$.

 $^{^6}$ beware that one uses an homological grading in loc. cit. while we are using a cohomological grading

6.2. Gerstenhaber-Schack cochain complexes for bialgebras and proof of Corollary 0.7. We now precisely relate Gerstenhaber-Schack complexes [29] and E_2 -Hochschild cohomology. As for $Pois_n$ and E_n -algebras, there are several more or less truncated complexes one can consider and encounter in the literature⁷, each related to different deformation problems. Let us fix some notations. What we call the Gerstenhaber-Schack complex is the total complex of a bicomplex, defined by

(6.21)
$$C_{GS}^*(B,B) \cong \prod_{m,n>1} Hom_{dg}(B^{\otimes m}, B^{\otimes n})[-m-n].$$

The horizontal differential is defined, for every n, by the Hochschild differential associated to the Hochschild complex of B seen as an associative algebra with coefficients in the B-bimodule $B^{\otimes n}$. The vertical differential is defined, for every m, by the co-Hochschild differential associated to the co-Hochschild complex of B seen as a coassociative coalgebra with coefficients in the B-bicomodule $B^{\otimes m}$. The compatibility between these differentials, which gives us a well defined bicomplex, follows from the distributive law relating the product and the coproduct of the bialgebra B (see [29, 59] for details). This is the complex relevant for Etingof-Kazdhan quantization in Section 8 because

Proposition 6.13 ([59]). The Gerstenhaber-Schack complex is quasi-isomorphic to the deformation complex of dg-bialgebras (up to isomorphisms):

$$C_{GS}^*(B,B) \cong g_{Bialg_{\infty},B}^{\varphi^+}.$$

Note here that in the definition of $C^*_{GS}(B,B)$ we use truncated versions of Hochschild and co-Hochschild complexes. We can also use their full versions and define an extended Gerstenhaber-Schack complex

(6.22)
$$\tilde{C}_{GS}^*(B,B) \cong \prod_{m \geq 0, n \geq 1} Hom_{dg}(B^{\otimes m}, B^{\otimes n})[-m-n],$$

which sits inside the fully extended Gerstenhaber-Schack complex

$$C_{GS}^{full}(B,B) \cong \prod_{m,n>0} Hom_{dg}(B^{\otimes m},B^{\otimes n})[-m-n].$$

The differentials are defined as for $C_{GS}^*(B,B)$, which is naturally a (non-split) subcomplex of each of the preceding ones ([29]).

Corollary 6.14. Let B be a conilpotent dg-bialgebra. There is a commutative diagram of L_{∞} -algebras

whose vertical arrows are L_{∞} -quasi-isomorphisms and the middle vertical arrow is given by Lemma 3.6 and the quasi-isomorphism (5.1). The horizontal arrows are further fiber sequences.

⁷beware that, unfortunately, the terminology, notations or degree shifting are not consistent in the literature.

Proof. We endow ΩB with its E_2 -algebra structure given by Corollary 0.2. The upper horizontal sequence is then given by the main result of [20]. The middle vertical equivalence is assertion (2) in Theorem 0.6. A proof similar to the one of Theorem 6.7 gives a commutative diagram of L_{∞} -algebras

, which induces a morphism of fiber sequences, and an identification of $g_{suBialg_{\infty}^{+},B}^{\varphi^{+}}$ with $C_{GS}^{*}(B,B)[2]$ similar to the one of $g_{suPois_{2}^{+},A}^{(\psi)^{+}}$ with $CH_{Pois_{2}}^{*}(A,A)$. The upper line has been identified with $T_{\tilde{\Omega}B} \longrightarrow CH_{E_{2}}^{*}(\tilde{\Omega}B,\tilde{\Omega}B)[2]$ in the proof of Theorem 6.7. The diagram being a diagram of fiber sequences of L_{∞} -algebras, we get that the fibers are also isomorphic as L_{∞} -algebras.

Let B be a conilpotent dg-bialgebra. Then ΩB is an E_2 -algebra by Corollary 0.2 and the sequence of L_{∞} -algebras

$$\tilde{\Omega}B[1] \longrightarrow T_{\tilde{\Omega}B} \longrightarrow CH_{E_2}^*(\tilde{\Omega}B, \tilde{\Omega}B)[2]$$

lits to a sequence of (non-unital) E_3 -algebras (after shifting it down by 2) by the solution to (Kontsevich) higher Deligne conjecture, and more precisely the main result of Francis in [20] (also see [54, 36] for the relationship with Pirashvili Hochschild cohomology and derived centralizers).

Finally we improved Corollary 6.14 into

Corollary 6.15 (Gerstenhaber-Schack conjecture). (1) The E_3 -algebra structures of $CH_{E_2}^{(\bullet>0)}(\tilde{\Omega}B,\tilde{\Omega}B)\cong T_{\tilde{\Omega}B}[-2]$ and $CH_{E_2}^*(\tilde{\Omega}B,\tilde{\Omega}B)$ induce E_3 -algebra structures on $C_{GS}^*(B,B)$ and $\tilde{C}_{GS}(B,B)$) such that the following diagram

is a commutative diagram of non-unital E_3 -algebras with vertical arrows being equivalences.

(2) The E_3 -algebra structure on $C_{GS}^*(B,B)$ is a refinement of its L_{∞} -algebra structure controlling the deformation theory of the bialgebra B.

Proof. The diagram and the existence of the lifts of the E_3 -structures are obtained by transfer of the structure of the first line [20] along the equivalences given by Corollary 6.14. We further know that the underlying L_{∞} -algebra structure of $T_{\tilde{\Omega}(B)}$ is equivalent to the one of the deformation complexes of Lie bialgebra $g_{Bialg_{\infty}^+,B}^{\varphi^+} \cong C_{GS}^*(B,B)$ by Theorem 0.6. Claim (2) follows.

Remark 6.16. We conjecture that the underlying L_{∞} -algebra structure of the full Gerstenhaber-Schack complex $C_{GS}^{full}(B,B)$ controls deformations as a monoidal dg category of the dg category of representations of B.

Remark 6.17 (Relevance of the choice of E_3 -lift). In view of Theorem 6.7, we can also endow the Gerstenhaber-Schack complexes $C_{GS}^*(B,B)$) and $C_{GS}^{full}(B,B)$) with E_3 -algebra structures given by the ones on Tamarkin deformation complexes $CH_{Pois_n}^{\bullet>0}(-,-)$, $CH_{Pois_n}^*(-,-)$ ([74, 7]), which carry the same underlying L_{∞} -structure. In view of Theorem 0.6, our choice of solutions to the Gerstehaber-Schack conjecture seems more natural.

With respect to applications to quantization of Lie bialgebras, this choice is however not important: any E_3 -algebra structure inducing the correct L_∞ -structure will be good enough to compute the deformation complex of bialgebras; the additional datum of the E_3 -structure is in fact used to simplify computations of deformation complexes by adding more rigidity, and gives in quantization the independence from the choice of an associator. In our case of interest for Etingof-Kazdhan quantization in § 8, we are in fact in a case where both E_3 -structures are the same. We actually believe that the diagram of Theorem 6.7 is in fact a commutative diagram of E_3 -algebras in general so that the aforementioned two E_3 -structures are always the same.

As a consequence, the Gerstenhaber-Schack complex $C_{GS}^*(B,B)$ inherits a homotopy associative multiplication (that is an E_1 -algebra structure obtained through the canonical map $E_1 \to E_3$ of operads). There is a standard dg associative multiplication on the Gerstenhaber-Schack complex given by the cup-product [29], which is a model for the Yoneda extension product [71]. On the other hand, following Corollary 6.15, the E_3 -algebra structure on $C_{GS}^*(B,B)$ is given by its identification with the derived center $\mathbb{R}Hom_{\tilde{\Omega}(B)}^{E_2}(\tilde{\Omega}(B),\tilde{\Omega}(B))$ of $\tilde{\Omega}(B)$. By definition, its E_1 -algebra structure is given by composition of E_2 - $\tilde{\Omega}(B)$ -modules endomorphisms which is the same as derived homomorphisms of left modules over factorization homology $\int_{S^1} \tilde{\Omega}(B)$ ([20, 36]); that is, it models the Yoneda extension product for E_2 - $\tilde{\Omega}(B)$ -modules. Then, from Theorem 0.6, one can deduce:

Proposition 6.18. The E_1 -structure induced by Corollary 6.15 on $C_{GS}^*(B,B)$ is equivalent to the E_1 -algebra structure induced by the cup-product.

7. The E_3 -formality Theorem

7.1. The Gerstenhaber-Schack complexes of Sym(V). This section is devoted to the proof of our E_3 -formality theorem (Theorem 0.5). Corollary 0.4 tells us two facts. First, we can choose the higher Hoschchild complex $CH_{E_2}^{(\bullet>0)}(\tilde{\Omega}B,\tilde{\Omega}B)$ as a model for the deformation complex Def(B) of a dg bialgebra B. By (Kontsevich) higher Deligne conjecture [20, 54, 36], the L_{∞} -structure controlling these deformations has a lift to an E_3 -algebra structure (and an E_3 -moduli problem). Then, the Gerstenhaber-Schack cohomology groups $H_{GS}^*(B,B)$ of B inherits a 3-Poisson algebra structure, that is, an algebra over the homology $H_*(E_3) \cong Pois_3$ of the little 3-disks operad, so that $H^*Def(B) \cong H^*(T_{\tilde{\Omega}(B)})$ and $H_{GS}^*(B,B)$ are isomorphic as 3-Poisson algebras.

The formality of the little 3-disks operad then gives an E_3 -algebra structure on $H_{GS}^*(B,B)$. The main question of interest here is to know whether $C_{GS}^*(B,B)$ and

 $H_{GS}^*(B,B)$ are then quasi-isomorphic as E_3 -algebras. We cannot expect such a result to holds true in full generality, however, for our purposes it will be sufficient to prove it in the case where B = Sym(V) is the symmetric bialgebra over a \mathbb{Z} -graded cochain complex V with cohomology of finite dimension in each degree. The general strategy is very similar to the famous obstruction theoretic approach used by Tamarkin to prove the E_2 -formality of the Lie algebra of polyvector fields on an affine space ([39],[72], [73]).

Now let (V,d) be a (\mathbb{Z} -graded) cochain complex with cohomology of finite dimension in each degree. Its symmetric algebra Sym(V) equipped with the induced differential (extending $d: V^{\bullet} \to V^{\bullet+1} \subset Sym(V)$ into a derivation) is a conilpotent dg bialgebra (see Example 5.5) so that we can apply Theorem 0.6.

Furthermore, there is a natural (strict) dg $Pois_3$ -algebra structure on (possibly completed) symmetric powers of $(V \oplus (V)^*)[-1]$ mimicking the Poisson structure of vector fields.

Definition 7.1. Let V be a cochain complex. We define a dg $Pois_3$ -algebra structure on

$$\widehat{Sym}(V[-1] \oplus (V)^*[-1]) \cong \prod_{m,n \ge 1} Sym^m(V[-1]) \otimes Sym^n((V)^*[-1])$$

the (fully completed) symmetric algebra (on $(V[-1] \oplus (V)^*[-1])$), with the usual differential given by the extension as a derivation of the one of V, and the degree -2 Poisson bracket induced by the evaluation pairing $ev: (V^{\bullet})^* \otimes V \to \mathbb{K}$, $[\chi, v] := ev(\chi \otimes v) = \chi(v)$ and the Leibniz rule:

$$[a \cdot b, c] = a \cdot [b, c] + (-1)^{|a||b|} b \cdot [a, c].$$

The subalgebra

$$\widehat{Sym}\big((V)^*[-1]\big)\otimes Sym(V[-1])\cong \prod_{m\geq 1}\bigoplus_{n\geq 1}Sym^m(V[-1])\otimes Sym^n((V)^*[-1])$$

is a dg sub- $Pois_3$ -algebra and such also are the sub-complexes

$$\widehat{Sym}^{\geq 1}\big((V)^*[-1]\big)\otimes Sym(V[-1]), \qquad \widehat{Sym}\big((V)^*[-1]\big)\otimes Sym^{\geq 1}(V[-1]).$$

Note that if $V^1=0$, then the completed symmetric algebra is just the usual symmetric algebra.

Theorem 7.2. Let (V, d) be a (\mathbb{Z} -graded) cochain complex with finite dimensional cohomology in each degre.

(1) The Gertenhaber-Schack cohomology of Sym(V) is given by

$$\begin{split} H^*_{GS}(Sym(V),Sym(V)) &\cong \prod_{m\geq 1} \big(\bigoplus_{n\geq 1} Sym^n(H^*(V)[-1]) \otimes Sym^m(H^*(V)^*[-1])\big) \\ &\cong \widehat{Sym}(H^*(V)^*[-1]) \otimes Sym(H^*(V)[-1]) \end{split}$$

as a Pois₃-algebra. Here, the algebra structure is the one of the symmetric algebra and the the degree -2 Poisson bracket is induced by the evaluation pairing $ev: H^*(V)^* \otimes H^*(V) \to \mathbb{K}$ as above.

(2) The Gertenhaber-Schack complex $C_{GS}^*(Sym(V), Sym(V))$ is equivalent as an E_3 -algebra to $\widehat{Sym}(V[-1] \oplus (V)^*[-1])$, where the later is equipped with

the E_3 -algebra structure induced by the chosen formality morphism $E_3 \xrightarrow{\sim} Pois_3$. Furthermore, there is a commutative diagram of E_3 -algebras

$$C^*_{GS}(Sym(V),Sym(V)) \xrightarrow{\simeq} Sym^{\geq 1}(V[-1]) \otimes \widehat{Sym}^{\geq 1}(V)^*[-1])$$

$$\widehat{C}^*_{GS}(Sym(V),Sym(V)) \xrightarrow{\simeq} Sym(V[-1]) \otimes \widehat{Sym}^{\geq 1}((V)^*[-1])$$

$$C^{full}_{GS}(Sym(V),Sym(V)) \xrightarrow{\simeq} Sym(V[-1]) \otimes \widehat{Sym}((V)^*[-1])$$

whose horizontal arrows are natural (with respect to V) equivalences of E_3 -algebras and the vertical arrows are the canonical inclusions.

(3) The three Gertenhaber-Schack complexes are formal as E_3 -algebras (in particular, there is an equivalence of E_3 -algebras between $C_{GS}^*(Sym(V), Sym(V))$ and $H_{GS}^*(Sym(V), Sym(V))$).

The theorem also holds with $T_{\tilde{\Omega}Sym(V)}$ and $CH_{E_2}(\tilde{\Omega}Sym(V), \tilde{\Omega}Sym(V))$ instead of $C_{GS}^*(Sym(V), Sym(V))$ and $\tilde{C}_{GS}^*(Sym(V), Sym(V))$ since the E_3 -structure on the later one is induced by transfer from the first one along the weak equivalence given by Theorem 0.6.

Proof of Theorem 0.8. Since Def(Sym(V)) is precisely the (shifted) Gerstenhaber-Schack complex with its underlying L_{∞} -structure, Theorem 0.6 shows that it suffices to prove that $CH_{E_2}^*(\tilde{\Omega}Sym(V),\tilde{\Omega}Sym(V))$ is formal as an L_{∞} -algebra. The previous Theorem 7.2 (in the case where V has trivial differential) shows the stronger statement that the later is actually formal as an E_3 -algebra, which concludes the proof.

Theorem 7.2 is a consequence of Proposition 7.4 and Proposition 7.3 below.

Proposition 7.3. There is a commutative diagram of E_3 -algebras

$$T_{Sym(V[-1])}[-2] \xrightarrow{\hspace*{2cm}} CH_{E_2}^*((Sym(V[-1]),(Sym(V[-1]))))$$

$$\uparrow \simeq \qquad \qquad \uparrow \simeq \qquad \qquad \uparrow \simeq$$

$$Sym(V[-1]) \otimes \widehat{Sym}^{\geq 1}((V)^*[-1]) \xrightarrow{\hspace*{2cm}} Sym(V[-1]) \otimes \widehat{Sym}((V)^*[-1])$$

whose vertical arrows are quasi-isomorphisms (and the upper map is the canonical one from Theorem 6.7).

Proof. By [7], [36, Corollary 6.39]⁸, we have, for a Sullivan algebra A (that is a cofibrant cdga), an equivalence of E_3 -algebras

(7.1)
$$CH_{E_2}^*(A,A) \simeq \widehat{Sym}_A(Der(A,A)[-2])$$

where the shifted Lie bracket is given by the Lie bracket on derivations and the Leibniz rule. Since the differential on Sym(V[-1]) is the unique derivation induced by the desuspension of $d:V^{\bullet}\to V^{\bullet+1}$, it is in particular a Sullivan algebra so

 $^{^{8}}$ beware that one uses an homological grading in loc. cit. while we are using a cohomological grading

that we can apply this result to A = Sym(V[-1]). For any cochain complex W with cohomology of finite dimension in each degree, we have an isomorphism of cochain complexes $Der(Sym(W), Sym(W)) \cong (W)^* \otimes Sym(W)$, and, under this equivalence, the Lie bracket of derivations is induced by the pairing $W \otimes (W)^* \to \mathbb{K}$. Thus, for A = Sym(V[-1]), the right hand side of the equivalence (7.1) is precisely $Sym(V[-1]) \otimes \widehat{Sym}^{\geq 1}((V)^*[-1])$ as an E_3 -algebra. Thanks to Theorem 6.7 (since our E_3 -algebra structures are induced by the one on Tamarkin complexes, we are in fact mainly using the sequence (6.4)), we can identify $T_{Sym(V[-1])}$ with the kernel of the canonical projection

$$Sym(V[-1]) \otimes \widehat{Sym}((V)^*[-1]) \rightarrow Sym(V[-1])$$

and the equivalences as well as the diagram follow.

Proof of Theorem 7.2. Recall that the E_3 -algebra structure on $\tilde{C}_{GS}(Sym(V), Sym(V))$ is obtained from the one on $CH_{E_2}^*(\tilde{\Omega}Sym(V), \tilde{\Omega}Sym(V))$ by transfer thanks to Theorem 0.6. Then, from Proposition 7.4 we obtain weak equivalences of E_3 -algebras

$$\begin{split} C_{GS}(Sym(V),Sym(V)) &\simeq T_{\tilde{\Omega}Sym(V)} \cong T_{Sym^{\geq 1}(V[-1])}, \\ \tilde{C}_{GS}(Sym(V),Sym(V)) &\simeq CH_{E_2}^*(\tilde{\Omega}Sym(V),\tilde{\Omega}Sym(V)) \\ &\cong CH_{E_2}^*(Sym^{\geq 1}(V[-1]),Sym^{\geq 1}(V[-1])). \end{split}$$

We are going to use Proposition 7.3 and Proposition 6.12 to compute these complexes. Indeed, we have a canonical augmentation $Sym(V[-1])) \cong \mathbb{K} \oplus Sym^{\geq 1}(V[-1])$ and these propositions as well as diagram (6.20) yields the equivalence of E_3 -algebras

$$Sym(V[-1]) \otimes \widehat{Sym}^{\geq 1}((V)^*[-1]) \xrightarrow{\simeq} \tilde{C}_{GS}(Sym(V), Sym(V)).$$

The commutative left square of the diagram of claim (2) follows similarly. Further, by Proposition 6.12, the right upper map of the diagram fits into a diagram of cofiber sequences

$$CH_{E_2}^*(Sym^{\geq 1}(V[-1])) \xrightarrow{\iota} CH_{E_2}^*(Sym(V[-1])) \xrightarrow{\pi_*} CH_{E_2}^*(Sym(V[-1]), \mathbb{K})$$

$$\cong \uparrow \qquad \qquad \cong \uparrow \qquad \qquad \cong \uparrow$$

$$Sym(V[-1]) \otimes \widehat{Sym}^{\geq 1}((V)^*[-1]) \hookrightarrow Sym(V[-1]) \otimes \widehat{Sym}((V)^*[-1]) \xrightarrow{\pi_*} \widehat{Sym}((V)^*[-1])$$

$$\cong \downarrow \qquad \qquad \cong \downarrow \qquad \qquad \cong \downarrow$$

$$\tilde{C}_{GS}(Sym(V), Sym(V)) \hookrightarrow C_{GS}^{full}(Sym(V), Sym(V)). \xrightarrow{\pi_*} C_{GS}^{full}(Sym(V), \mathbb{K}).$$

Hence the commutativity of the right hand square. This proves assertion (2).

Assertion (1) follows from assertion (2) by passing to the cohomology groups, while assertion (3) also follows from (2) by choosing any quasi-isomorphism $H^*(V) \stackrel{\sim}{\to} V$ since we work over a characteristic zero field.

7.2. Enhanced Cobar functor on Sym(V). The main goal of this section is to compute the value of our enhanced cobar functor $\tilde{\Omega}: E_1 - Alg(dgCog^{conil}) \to E_2 - Alg^{aug}$ on the commutative and cocommutative dg bialgebra Sym(V), where (V,d) is a cochain complex with cohomology of finite dimension in each degree:

Proposition 7.4. There is a quasi-isomorphism of E_2 -algebras $\tilde{\Omega}(Sym(V)) \stackrel{\sim}{\to} Sym^{\geq 1}(V[-1])$.

We first start by exhibiting a cofibrant resolution of Sym(V) in $E_1-Alg(dgCog^{conil})$.

Lemma 7.5. There is a cofibrant resolution $Cobar(Sym(V[1])) \rightarrow Sym(V)$ of Sym(V) in $E_1 - Alg(dgCog^{conil})$.

Proof. This is dual to the usual Hochschild-Kostant-Rosenberg theorem for symmetric algebras: the composition $p:Sym(V[1])[-1] \to V \hookrightarrow Sym(V)$ of the canonical projection with the canonical inclusion, yields the map of dg algebras $\pi:T(Sym(V[1])[-1]) \to Sym(V)$, which is further a map of coalgebras with respect to the shuffle coproduct on the source. Then Cobar(Sym(V[1])) is the semifree dg algebra obtained from T(Sym(V[1])[-1]) by adding the extra-differential $\delta(x_1 \otimes x_n) := \sum_{i=1}^n \pm x_1 \otimes \cdots \otimes x_i^{(1)} \otimes x_i^{(2)} \otimes \cdots \otimes x_n \in T^{n+1}(Sym(V[1])[-1])$, using the coalgebra structure of Sym(V[1]). Since the latter is cofree cocommutative on the shift of V, we get that $p \circ \delta = 0$. Hence $\pi:Cobar(Sym(V[1])) \to Sym(V)$ is indeed a map in $E_1 - Alg(dgCog^{conil})$. This is a quasi-isomorphism by the same argument (replacing algebras by coalgebras) as for the bar construction of a symmetric algebra. Indeed, Cobar(Sym(V[1])) is the derived cotensor product of \mathbb{K} and Sym(V[1]) as Sym(V[1])-bicomodules and we have a resolution $Sym(V[1]) \xrightarrow{\Delta} Sym(V[1]) \otimes \bigoplus Sym^n(V) \otimes Sym(V[1])$, where the target is the tensor product of cocommutative coalgebras equipped with the extra differential $\delta(x \otimes f \otimes y) = x^{(1)} \otimes p(x^{(2)} \cdot f \otimes y \pm x^{\otimes} f \cdot p(y^{(1)}) \otimes y^{(2)}$.


In fact one can also directly apply the usual HKR theorem which gives a quasi-isomorphism of Hopf algebras $Sym(V[1]) \to Bar(Sym(V))$ (given by the unique dg coalgebra map extending the linear map $Sym(V[1]) \stackrel{p}{\to} V[1] \hookrightarrow Sym(V)[1]$) and then conclude by the counit of the bar-cobar adjunction.

Let

$$\phi^*\mathcal{B}^{enh}_{E_1}(-)_-: E_1 - Alg^{con,aug}(dgCog^{conil}) \to E_2 - Cog^{conil}$$

be the equivalence given by Theorem 0.1 and Section 5.2. We wish to evaluate $\phi^*\mathcal{B}_{E_1}^{enh}(-)_-$ on Sym(V).

Note that we have a commutative diagram of $(\infty$ -)operads



which in turn induces a commutative diagram

where obl is induced by forgetting the dg-coalgebra structure and the non-labbeled arrows are given by the standard restriction functors between E_m -algebras.

The differential d on a cochain complex (V, d) extends canonically to the coartinian cofree cocommutative coalgebra Sym(V), giving it a canonical dg-cocommutative coalgebra structure and thus $E_2 - Cog^{conil}$ structure by restriction.

Lemma 7.6. One has an equivalence $\phi^*\mathcal{B}^{enh}_{E_1}(Sym(V))_- \cong Sym^{\geq 1}(V[1])$ in $E_2 - Coq^{conil}$.

Proof. Applying the argument line of the proof of Lemma 7.5 to the augmentation ideal $Sym^{\geq 1}(V)$ of Sym(V), we have

$$Sym^{\geq 1}(V) \simeq Cobar(Sym^{\geq 1}(V[1]))$$

in $E_1 - Alg(dgCog^{conil})$.

Applying the functor $\mathcal{B}_{E_1}^{enh}$ gives an equivalence

$$\mathcal{B}^{enh}_{E_1}Sym^{\geq 1}(V)\simeq \mathcal{B}^{enh}_{E_1}Cobar(Sym^{\geq 1}(V[1]))$$

in $E_1 - Cog^{conil}(dgCog^{conil})$, hence

$$\phi^*\mathcal{B}^{enh}_{E_1}Sym^{\geq 1}(V)\simeq \phi^*\mathcal{B}^{enh}_{E_1}Cobar(Sym^{\geq 1}(V[1]))$$

in $E_2 - Cog^{conil}$. It thus remains to prove that $\mathcal{B}_{E_1}^{enh}Cobar(Sym^{\geq 1}(V[1])) \simeq Sym^{\geq 1}(V[1])$ in $E_1 - Cog^{conil}(dgCog^{conil})$.

Recall that the Koszul duality of E_1 -operads provides us a Quillen equivalence

$$Cobar: E_1 - Cog^{conil} \rightleftharpoons E_1 - Alg^{aug}: Bar$$

where Bar is weakly equivalent to the derived functor of indecomposables. Let us note that the unit and counit of this adjunction are actually always weak equivalences, because every object of $E_1 - Cog^{conil}$ is cofibrant (cofibrations of coalgebras are determined in chain complexes) and every object of $E_1 - Alg^{aug}$ is fibrant (fibrations of algebras are determined in chain complexes). The fact that E_1 is a Hopf operad gives a distributive law between E_1 and Ass allowing to lift these functors as follows

$$E_{1} - Alg^{aug}(dgCog^{conil})$$

$$Cobar \longrightarrow E_{1} - Cog^{conil}(dgCog^{conil}) \longrightarrow E_{1} - Cog^{conil} \longrightarrow E_{1} - Alg^{aug}$$

where oblv is the forgeful functor, and

$$E_{1} - Cog^{conil}(dgCog^{conil})$$

$$E_{1} - Alg^{aug}(dgCog^{conil}) \xrightarrow{oblv} E_{1} - Alg^{aug} \xrightarrow{Bar} E_{1} - Cog^{conil}$$

This gives us a new adjunction

$$Cobar: E_1 - Cog^{conil}(dgCog^{conil}) \rightleftharpoons E_1 - Alg^{aug}(dgCog^{conil}): Bar$$

whose unit and counit are still weak equivalences (recall that weak equivalences here are all determined in cochain complexes). This functor Cobar is the one of Lemma 7.5 and the functor Bar is actually weakly equivalent to $\mathcal{B}_{E_1}^{enh}$ (they are

both equivalent to the derived functor of indecomposables), which means that we have a natural quasi-isomorphism

$$Id \stackrel{\sim}{\to} \mathcal{B}_{E_1}^{enh} \circ Cobar,$$

giving in particular

$$Sym^{\geq 1}(V[1]) \stackrel{\sim}{\to} \mathcal{B}^{enh}_{E_1}Cobar(Sym^{\geq 1}(V[1]))$$

in
$$E_1 - Cog^{conil}(dgCog^{conil})$$
.

The last step in the construction of the functor Ω of Corollary 0.2 is the Koszul duality equivalence $Cobar^{(n)}: E_n - Cog^{conil} \to E_n - Alg^{aug}$ in the case n=2. We wish to prove that $Cobar^{(n)}(Sym^{\geq 1}(W)) \cong Sym^{\geq 1}(W[-n])$. A first step is

We wish to prove that $Cobar^{(n)}(Sym^{\geq 1}(W)) \cong Sym^{\geq 1}(W[-n])$. A first step is to prove that, by formality of E_n -operads, it is enough to prove this result with the operad $Pois_n$ (instead of E_n), given that $Sym^{\geq 1}(W)$ can be seen as a $Pois_n$ -coalgebra. For this, we use several features of the theory of right modules over operads as thoroughly studied in [22].

Given any operad R equipped with an operad morphism $E_1 \to R$, one can extend the bar construction from E_1 -algebras to R-algebras: this extended bar construction is the functor $S_R(B_R, -): R - Alg \to Ch_{\mathbb{K}}$ associated to a certain cofibrant quasi-free right R-module B_R . Moreover, any weak equivalence of operads $\varphi: R \xrightarrow{\sim} S$ induces a weak equivalence of right R-modules $B_R \xrightarrow{\sim} \varphi^* B_S$, where $\varphi^* B_S$ is B_S equipped with the right R-module structure induced by φ . In the case of E_n -operads, it turns out that this extended bar construction is given by the iterated bar construction $Bar^{(n)}$ [25] (right adjoint to $Cobar^{(n)}$).

Let us fix a formality morphism $\varphi: E_n \xrightarrow{\sim} Pois_n$ and denote respectively $Bar_{Pois_n}^{(n)}$ and $Cobar_{Pois_n}^{(n)}$ the Koszul duality Bar and Cobar construction for $Pois_n$ -algebras.

Lemma 7.7. Let $n \geq 2$. One has natural equivalences

$$Bar^{(n)} \circ \varphi^* \sim Bar^{(n)}_{Pois_n},$$

where φ^* is the restriction of structures from Pois_n-algebras to E_n -algebras and

$$Cobar^{(n)} \circ \varphi^* \sim Cobar^{(n)}_{Pois_n}$$

where φ^* is the restriction of structures from $Pois_n$ -coalgebras to E_n -coalgebras.

Proof. Given the formality morphism $\varphi: E_n \xrightarrow{\sim} Pois_n$, according to [22, Theorem 7.2.2], there is a natural isomorphism

$$S_{E_n}(B_{E_n}, -) \circ \varphi^* \cong S_{Pois_n}(\varphi_! B_{E_n}, -)$$

where φ^* is the restriction of structures fitting in the adjunction

$$\varphi_! : Pois_n - Alg \rightleftharpoons E_n - Alg : \varphi^*$$

and φ_1 is the extension of structures fitting in the adjunction

$$\varphi_! : Pois_n - Mod \rightleftharpoons E_n - Mod : \varphi^*$$

between right $Pois_n$ -modules and right E_n -modules. Since φ is a weak equivalence, by [22, Theorem 16.B] the adjunction above between right modules is a Quillen equivalence, so the weak equivalence of right E_n -modules $B_{E_n} \stackrel{\sim}{\to} \varphi^* B_{Pois_n}$ corresponds by adjunction to a weak equivalence of right $Pois_n$ -modules $\varphi_! B_{E_n} \stackrel{\sim}{\to} B_{Pois_n}$. Moreover, since $\varphi_!$ is a left Quillen functor, the right module $\varphi_! B_{E_n}$ is still

cofibrant, so this is a weak equivalence of cofibrant right $Pois_n$ -modules. By [22, Theorem 15.1.A], it induces consequently a natural weak equivalence

$$S_{E_n}(B_{E_n}, -) \circ \varphi^* \cong S_{Pois_n}(\varphi_! B_{E_n}, -) \xrightarrow{\sim} S_{Pois_n}(B_{Pois_n}, -),$$

hence

$$Bar^{(n)} \circ \varphi^* \sim Bar^{(n)}_{Pois_n}$$

where φ^* is the restriction of structures. This proves the first claim. Since $Bar^{(n)}$ fits in a Quillen equivalence with $Cobar^{(n)}$ (for both operads) as a left adjoint, the first claim also implies that

$$Cobar^{(n)} \circ \varphi^* \sim Cobar^{(n)}_{Pois_n}$$

as well, where φ^* is now the restriction of structures from $Pois_n$ -coalgebras to E_n -coalgebras.

We now compute $Cobar_{Pois_n}^{(n)}$ on $Sym^{\geq 1}(W)$. Recall that to a adg Lie coalgebra $(\mathfrak{g}, \delta, d)$, we can associate a $Pois_n$ -coalgebra defined by $Sym^{\geq 1}(\mathfrak{g}[n-1])$ with cocommutative cobracket given by the one of $Sym^{\geq 1}(-) = Cocom$ (the free cocommutative conilpotent coalgebra functor) and Lie cobracket induced by the one of \mathfrak{g} and the Leibniz rule. Note that $\mathfrak{g} \mapsto Sym^{\geq 1}(\mathfrak{g}[1-n])$ is the right adjoint of the canonical functor $Pois_n - Coalg \to Lie - Coalg$ given by $P \mapsto P[1-n]$. The following Lemma is rather standard

Lemma 7.8. One has

$$Cobar_{Pois_n}^{(n)}(Sym^{\geq 1}(\mathfrak{g}[n-1])) \cong \left(Com(\mathfrak{g}[-1]), d_{CE}\right)$$

where $Cobar_{Pois_n}^{(n)}(-)$ is the E_n -Koszul duality functor $Pois_n - Cog^{conil} \to Pois_n - Alg^{aug}$ and the right hand side is the part of positive weight of the Chevalley-Eilenberg algebra of the dg-Lie coalgebra \mathfrak{g} .

Proof. By Koszul duality of the $Pois_n$ -operad, the functor $Cobar_{Pois_n}^{(n)}: dg - Pois_n - Cog^{conil} \rightarrow dg - Pois_n - Alg^{aug}$ on C is given by the free $Pois_n^{(n)}$ -algebra on C endowed with the differential induced by its $Pois_n$ -coalgebra structure. Recall that $Pois_n^! = \Lambda^n Pois_n$. Hence, one has an equivalence

$$Cobar^{(n)}(C) = Com(Lie(C[-1])[1-n])$$

see [74, 33, 39]. We are interested in the case where $C = Sym^{\geq 1}(\mathfrak{g}[n-1])$ which is cofree as a conilpotent cocommutative coalgebra (in cochain complexes). It follows that the Harrison cochain complex $Har_*(Sym^{\geq 1}(\mathfrak{g}[n-1])) = Lie(Sym^{\geq 1}(\mathfrak{g}[n-1]))$ of $Sym^{\geq 1}(\mathfrak{g}[n-1])$ is quasi-isomorphic to $(\mathfrak{g}[n-1])[-1]$. Hence we have a quasi-isomorphism of cochain complexes

$$Cobar^{(n)}(Sym(\mathfrak{g}[n-1])) = Com\left(Lie(Sym^{\geq 1}(\mathfrak{g}[n-1])[-1])[1-n]\right) \simeq Com(\mathfrak{g}[-1])$$

where the right hand side is identified with the part of positive weight in the Chevalley-Eilenberg algebra of the Lie coalgebra \mathfrak{g} . (see [72, 33, 39] for detailled computation of this dg- $Pois_n$ -algebra cohomology of the free $Pois_n$ -algebra generated by a dq-Lie algebra).

We finally compute $Cobar^{(n)}$ on $Sym^{\geq 1}(W)$.

Lemma 7.9. Let $n \geq 2$. One has $Cobar^{(n)}(Sym^{\geq 1}(W)) \cong Sym^{\geq 1}(W[-n])$ where $Cobar^{(n)}$ is the E_n -Koszul duality functor $E_n - Cog^{conil} \rightarrow E_n - Alg^{aug}$.

Proof. By Lemma 7.7, we conclude that proving the result for the E_n -cobar construction $Cobar^{(n)}(Sym^{\geq 1}(W))$ on $Sym^{\geq 1}(W)$ considered as an E_n -coalgebra boils down to check it for the $Pois_n$ -cobar construction $Cobar^{(n)}_{Pois_n}(-)$ on $Sym^{\geq 1}(W)$ considered as a $Pois_n$ -coalgebra with trivial cobracket. Applying the previous Lemma 7.8 to the case $\mathfrak{g} = W[1-n]$ equipped with zero cobracket (or simply redoing the proof of the Lemma in this simpler case), we get

$$Cobar^{(n)}(Sym^{\geq 1}(W)) = Com(Lie(Sym^{\geq 1}(W)[-1])[1-n]) \simeq Com(W[-n])$$

where the right hand side is identified with the part of positive weight of the Chevalley-Eilenberg algebra of a Lie coalgebra with a null cobracket, since $Sym^{\geq 1}(W)$ has a null cobracket. Hence it is formal and we get that $Cobar^{(n)}(Sym^{\geq 1}(W)) \simeq Sym^{\geq 1}(W[-n])$.

Proof of Proposition 7.4. By definition of the functor $\tilde{\Omega}$ (Section 5.2), it is the composition of functors $Cobar^{(2)} \circ \phi^* \mathcal{B}_{E_1}^{enh}(-)_-$. Hence, one has

$$\begin{split} \tilde{\Omega}(Sym(V)) & \simeq & Cobar^{(n)} \left(\phi^* \mathcal{B}^{enh}_{E_1}(Sym^{\geq 1}(V)) \right) \\ & \simeq & Cobar^{(n)} \left(Sym^{\geq 1}(V[1]) \right) \quad \text{by Lemma 7.6} \\ & \simeq & Sym^{\geq 1}(V[1][-2]) = Sym^{\geq 1}(V[-1]) \end{split}$$

by Lemma 7.9 in the case n=2.

8. Etingof-Kazdhan deformation quantization

We now apply our computations of the L_{∞} -structure of the Gerstenhaber-Schack complex to quantization of Lie bialgebras and prove Corollary 0.9.

If V is a finite dimensional vector space, by [58, Corollary 5.1], the Maurer-Cartan elements of the graded Lie algebra

$$\prod_{m\geq 1} \Big(\bigoplus_{n\geq 1} Sym^n(V[-1]) \otimes Sym^m((V)^*[-1])\Big)[2]$$

(for the Lie bracket induced by the Poisson bracket of Definition 7.1) are exactly the prop morphisms $BiLie_{\infty}^{+} \to End_{V}$, that is, the homotopy Lie bialgebra structures on V. Here BiLie is the prop governing (dg-)Lie bialgebras structures on (dg-)vector spaces, $BiLie_{\infty}$ is its standard cofibrant resolution, and $BiLie_{\infty}^{+}$ is the result of the + construction (Section 3) on the later one.

By Theorem 7.2, this Lie algebra is the one of $H^*_{GS}(Sym(V), Sym(V))$ and is quasi-isomorphic to the deformation complex $C^*_{GS}(Sym(V), Sym(V))$. This is the first step to derive the Etingof-Kazdhan quantization theorem from the formality theorem

Next, let $\varphi^+: Bialg^+_{\infty} \to End^{Poly}_{Sym(V)}$ be the map canonically associated to the (cocommutative and commutative) bialgebra structure on Sym(V). The Etingof-Kazdhan quantization theorem assert that any Maurer-Cartan element in the above Lie algebra gives rise functorially to a formal deformation of ψ^+ , in other words a point of the moduli space $Bialg^+_{\infty}\{Sym(V)\}^{\varphi^+}(k[[\hbar]])$. These observations (a dg extension of the quantization of Lie bialgebra was established in the Appendix of [34]) can be made for a dg vector space V as well.

According to Merkulov [59], there is an explicit completed dg prop $(DefQ^+, d^+)$ such that, for any cochain complex (V, d_V) , the representations of $(DefQ^+, d^+)$ into (V, d_V) are in bijection with polydifferential representations of homotopy bialgebra structures on the cochain algebra $(Sym(V), d_V)$. In other words

$$(8.1) Mor_{Prop}(DefQ^+, End_V) \cong Mor_{Prop}(Bialg_{\infty}^+, End_{Sym(V)}^{Poly}).$$

Polydifferential representations are explained in [58, 59]; they amount to replace the endomorphism prop $End_{Sym(V)}$ with its polydifferential endomorphism $End_{Sym(V)}^{Poly}$ which is the dg-sub-prop of $End_{Sym(V)}$ spanned by those multilinear maps that are (normalized) polydifferential operators on Sym(V). Merkulov [59] has proved that the natural map

(8.2)
$$Mor_{Prop}(DefQ^+, End_V) \xrightarrow{\simeq} Mor_{Prop}(Bialg_{\infty}^+, End_{Sym(V)}).$$
 is a quasi-isomorphism.

Using the above analysis, the formality Theorem 0.8 implies the following quantization theorem. Following the previous notation, we have the map $\varphi^+: Bialg_\infty^+ \to End_{Sym(V)}$ giving the dg bialgebra structure of Sym(V) for a cochain complex V. Further, we also have $0^+: BiLie_\infty^+ \to End_V$ the (trivial) map corresponding to the trivial (= null) Lie bialgebra structure on V.

Theorem 8.1. Let (V, d) be a (\mathbb{Z} -graded) cochain complex with finite dimensional cohomology in each degree. There is a weak-equivalence of formal moduli problems

$$Bialg_{\infty}^{+} \{Sym(V)\}^{\varphi^{+}} \simeq BiLie_{\infty}^{+} \{V\}^{0^{+}}$$

In particular, given an artinian cdga R, any R-deformation of the trivial Lie bialgebra structure on V (in other words a dg-bialgebra structure on $V \otimes R$) gives rise to a (essentially unique isomorphism class of) R-deformation of Sym(V).

Proof. By Theorem 7.2 and Theorem 0.6, we have an equivalence of deformation complexes

$$\begin{split} Def(Sym(V)) &\simeq g_{Bialg^+_{\infty},Sym(V)}^{\varphi^+} &\simeq C_{GS}^*(Sym(V),Sym(V)) \\ &\simeq Sym^{\geq 1}(V[-1]) \otimes \widehat{Sym}^{\geq 1}(V)^*[-1]) \\ &\simeq g_{BiLie^+_{\infty},V}^{0^+} \end{split}$$

where the last equivalence is from [58, Corollary 5.1]. Then the result follows from Theorem 2.10 or Lurie classification of (commutative) formal moduli problem [52].

The theorem also implies that formal deformations of the null Lie bialgebra structure quantize into formal deformation s of the symmetric bialgebra on V, hence Corollary 0.9.

Proof of Corollary 0.9. The first claim of the corollary is equivalent to the existence of a natural (in V) weak-equivalence

$$(8.3) \qquad \underline{Bialg_{\infty}^{+}\{Sym(V)\}^{\varphi^{+}}(k[[\hbar]])} \simeq \underline{BiLie_{\infty}^{+}\{V\}^{0^{+}}(k[[\hbar]])}$$

 $^{^{9}}$ completion for the filtrations induced by the genus and the number of vertices

where, for any formal moduli problem \mathcal{F} , we denote

$$\mathcal{F}(k[[\hbar]]) := \lim_{i} \mathcal{F}(k[\hbar]/h^{i}).$$

By [78, Corollaire 2.11] (or [52]), there is an natural weak-equivalence

(8.4)
$$\mathcal{F}(k[[\hbar]]) \simeq Map(k[-1], g_{\mathcal{F}})$$

where $g_{\mathcal{F}}$ is the L_{∞} -algebra associated to the formal moduli problem \mathcal{F} and k[-1] is the abelian one dimensional Lie algebra concentrated in degree 1.

The equivalence (8.4) now implies the first claim (8.3) in virtue of Theorem 8.1 (and Theorem 5.6).

For claim (2), we note that the above proof relies essentially on the formality Theorem 7.2 for the Gerstenhaber-Schack L_{∞} -algebra of Sym(V). This theorem relies on the formality of the E_n -operad for $n \geq 3$, which does not rely on the choice of a Drinfeld associator.

Our proof somehow goes in the converse way as Merkulov [59] proof of a dg prop quasi-isomorphism $EK: (DefQ^+, d^+) \to (\widehat{BiLie}, 0)$, which relies in loc. cit. on the existence of the Etingof-Kazdhan quantization functor for dg Lie bialgebras. Our proof establishes first this equivalence, from which the above statement actually follows. Applied to an ordinary finite dimensional vector space, our result gives a new proof of Etingof-Kazhdan deformation quantization. Indeed, homotopy Lie bialgebra structures $BiLie_{\infty} \to End_V$ are in this case strict Lie bialgebra structures: since the endomorphism prop is concentrated in degree 0 and prop morphisms preserve the degree, the generators of positive degree in $BiLie_{\infty}$ are sent to zero. In particular, the 1-cycles whose images under the differential of $BiLie_{\infty}$ are the 0-boundaries inducing the Jacobi, co-Jacobi and Drinfeld's compatibility relations in homology, are sent to 0. Since dg prop morphisms commute with the differentials, this means that these 0-boundaries are also sent to 0. That is, the two generators of degree 0 in $BiLie_{\infty}$ are sent to a bracket and a cobracket satisfying the Jacobi, co-Jacobi and Drinfeld's compatibility relations, i.e. a strict Lie bialgebra structure. This is a particular incarnation of the general fact that any P_{∞} -algebra structure on a vector space is actually a strict one. Non trivial homotopy algebra structures appear only in the differential graded setting. By the same argument, homotopy bialgebra structures $Bialg_{\infty} \to End_{Sym(V)}$ are only the ones factoring through Bialg.

Moreover, it further applies to any \mathbb{Z} -graded dg Lie bialgebra g with cohomology of finite total dimension as well as homotopy Lie bialgebra structures on g.

References

- [1] M. Anel, Champs de modules des catégories linéaires et abéliennes, PhD thesis, Université Toulouse III Paul Sabatier (2006).
- [2] M. Anel, A. Joyal, Sweedler Theory for (co)algebras and the bar-cobar constructions, arXiv:1309.6952.
- [3] D. Ayala, J. Francis, Poincaré/Koszul duality, preprint arXiv:1409.2478.
- [4] C. Barwick, D. M. Kan, Relative categories: another model for the homotopy theory of homotopy theories, Indagationes Mathematicae 23 (2012), 42-68.
- [5] H.-J. Baues, The cobar construction as a Hopf algebra, Invent. Math. 132 (1998), no. 3, 467–489.
- [6] C. Berger, B. Fresse, Combinatorial operad actions on cochains, Math. Proc. Cambridge Philos. Soc. 137 (2004), 135–174.

- [7] D. Calaque, T. Willwacher, Triviality of the higher Formality Theorem, Proc. Amer. Math. Soc. 143 (2015), no. 12, 5181–5193.
- [8] D. Calaque, T. Pantev, B. Toen, M. Vaquié, G. Vezzosi, Shifted Poisson structures and deformation quantization, preprint arXiv:1506.03699.
- [9] V. G. Drinfeld, On quasitriangular quasi-Hopf algebras and on a group that is closely connected with Gal(Q/Q, Algebra i Analiz 2 (1990), 149-181. English translation in Leningrad Math. J. 2 (1991), 829-860.
- [10] V. G. Drinfeld, Quantum groups, in Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), 798–820, Amer. Math. Soc., Providence, RI.
- [11] W. Dwyer, D. Kan, Function complexes in homotopical algebra, Topology 19 (1980), 427-440.
- [12] B. Enriquez, On Quantization Functors of Lie Bialgebras, Acta Appl. Math. 73 (2002), Issue 1-2, 133-140.
- [13] B. Enriquez, P. Etingof, On the invertibility of quantization functors, J. Algebra 289 (2005), no. 2, 321-345.
- [14] B. Enriquez, G. Halbout, Quantization of quasi-Lie bialgebras, J. Amer. Math. Soc. 23 (2010), 611–653.
- [15] B. Enriquez, G. Halbout, Quantization of coboundary Lie bialgebras, Ann. of Math. (2) 171 (2010), 1267–1345.
- [16] P. Etingof, D. Kazhdan, Quantization of Lie bialgebras I, Selecta Math. (N. S.) 2 (1996), no. 1, 1-41.
- [17] P. Etingof, D. Kazdhan, Quantization of Lie bialgebras. II, III, Selecta Math. (N.S.) 4 (1998), no. 2, 213–231, 233–269.
- [18] Z. Fiedorowicz, R. M. Vogt, An Additivity Theorem for the interchange of E_n structures, Adv. Math. 273 (2015), 421-484.
- [19] J. Francis, D. Gaitsgory, Chiral Koszul duality, Selecta Math. New Ser. 18 (2012), 27-87.
- [20] J. Francis, The tangent complex and Hochschild cohomology of E_n-rings, Compositio Mathematica 149 (2013), no. 3, 430-480.
- [21] B. Fresse, Koszul duality of operads and homology of partition posets, Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K-theory, 115–215, Contemp. Math. 346 (2004).
- [22] B. Fresse, Modules over operads and functors, Lecture Notes in Mathematics 1967, Springer-Verlag (2009).
- [23] B. Fresse, Props in model categories and homotopy invariance of structures, Georgian Math. J. 17 (2010), pp. 79-160.
- [24] B. Fresse, Koszul duality of En-operads, Selecta Math. (N.S.) 17 (2011), 363-434.
- [25] B. Fresse, Iterated bar complexes of E-infinity algebras and homology theories, Alg. Geom. Topol. 11 (2011), pp. 747-838.
- [26] B. Fresse, Homotopy of operads and Grothendieck-Teichmuller groups, book in progress available at http://math.univ-lille1.fr/~fresse/OperadHomotopyBook/
- [27] B. Fresse, The cotriple resolution of differential graded algebras, to appear in Proc. Amer. Math. Soc.
- [28] B. Fresse, T. Willwacher, The intrinsic formality of E_n-operads, preprint arXiv:1503.08699.
- [29] M. Gerstenhaber, S. D. Schack, Algebras, bialgebras, quantum groups, and algebraic deformations, Deformation theory and quantum groups with applications to mathematical physics (Amherst, MA, 1990), 51-92, Contemp. Math., 134, Amer. Math. Soc., Providence, RI, 1992.
- [30] E. Getzler, J. D. S. Jones Operads, homotopy algebra and iterated integrals for double loop spaces, preprint arXiv:hep-th/9403055 (1994).
- [31] E. Getzler, P. Goerss, A model category structure for differential graded coalgebras, preprint (1999).
- [32] J. Giansiracusa, P. Salvatore, Formality of the framed little 2-discs operad and semidirect products, Homotopy theory of function spaces and related topics, Contemp. Math. 519 (2010), 115–121.
- [33] G. Ginot, Homologie et modèle minimal des algèbres de Gerstenhaber, Annales Mathématiques Blaise Pascal 11 (2004) no. 1, 95-127.
- [34] G. Ginot, G. Halbout, A Formality Theorem for Poisson Manifolds, Lett. Math. Phys. 66 (2003), 37-64.
- [35] G. Ginot, T. Tradler, M. Zeinalian, A Chen model for mapping spaces and the surface product, Ann. Sc. de l'Éc. Norm. Sup., 4e série, t. 43 (2010), p. 811-881.

- [36] G. Ginot, T. Tradler, M. Zeinalian, Higher Hochschild cohomology of E-infinity algebras, Brane topology and centralizers of E-n algebra maps, preprint arXiv:1205.7056.
- [37] K. Hess, B. Shipley, The homotopy theory of coalgebras over a comonad, Proc. London Math. Soc. (3) 108 (2014), 484–516.
- [38] V. Hinich, DG coalgebras as formal stacks, J. Pure Appl. Algebra 162 (2001), 209-250.
- [39] V. Hinich, Tamarkin's proof of Kontsevich formality theorem, Forum Math. 15 (2003), 591–614.
- [40] V. Hinich, D. Lemberg, Formality theorem and bialgebra deformations, preprint arXiv:1410.2132
- [41] Mark Hovey, Model categories, Mathematical Surveys and Monographs volume 63, AMS (1999).
- [42] Philip S. Hirschhorn, Model categories and their localizations, Mathematical Surveys and Monographs volume 99, AMS (2003).
- [43] T. Kadeishvili, On the cobar construction of a bialgebra, Homology Homotopy Appl. 7 (2005), 109–122.
- [44] A. Kapustin, Topological field theory, higher categories, and their applications, in Proceedings of the International Congress of Mathematicians. Volume III, Hindustan Book Agency, New Delhi (2010), 2021–2043.
- [45] B. Keller, A-infinity algebras, modules and functor categories, in Trends in representation theory of algebras and related topics, Contemp. Math. 406 (2006), 67-93.
- [46] B. Keller and W. Lowen, On Hochschild cohomology and Morita deformations, Int. Math. Res. Not. IMRN 2009, no. 17, 3221–3235.
- [47] M. Kontsevich, Operads and motives in deformation quantization, Moshé Flato (1937-1998), Lett. Math. Phys. 48 (1999), no. 1, 35-72.
- [48] M. Kontsevich, Deformation quantization of Poisson manifolds, Lett. Math. Phys. 66 (2003), no. 3, 157-216.
- [49] M. Kontsevich and Y. Soibelman, Notes on A_{∞} -algebras, A_{∞} -categories and non-commutative geometry, in *Homological mirror symmetry*, 153–219, Lecture Notes in Phys., 757, Springer, Berlin.
- [50] P. Lambrechts, and I. Volic, Formality of the little N-disks operad, Mem. Amer. Math. Soc. 230 (1079), 2014.
- [51] J-L. Loday, B. Vallette, Algebraic Operads, Grundlehren der mathematischen Wissenschaften, Volume 346, Springer-Verlag (2012).
- [52] J. Lurie, Derived Algebraic Geometry X, available at http://www.math.harvard.edu/~lurie/
- [53] J. Lurie, Higher topos theory, Annals of Mathematics Studies 170, Princeton University Press, Princeton, NJ, 2009.
- [54] J. Lurie, Higher Algebra, book available at http://www.math.harvard.edu/~lurie/
- [55] J. P. May, The geometry of iterated loop spaces, Springer, Berlin, 1972.
- [56] S. Merkulov, B. Vallette, Deformation theory of representation of prop(erad)s I, J. für die reine und angewandte Math. (Crelles Journal), Issue 634 (2009), 51-106.
- [57] S. Merkulov, B. Vallette, Deformation theory of representation of prop(erad)s II, J. für die reine und angewandte Math. (Crelles Journal), Issue 636 (2009), 125-174.
- [58] S. Merkulov, Prop profile of Poisson geometry, Comm. Math. Phys. 262 (2006), 117–135.
- [59] S. Merkulov, Formality theorem for quantization of Lie bialgebras, Lett. Math. Phys. 106 (2016), no. 2, 169–195.
- [60] R. J. Milgram, Iterated loop spaces, Ann. of Math. (2) 84 (1966), 386–403.
- [61] F. Muro, Moduli spaces of algebras over nonsymmetric operads, Algebr. Geom. Topol. 14 (2014) 1489–1539.
- [62] Pr A. Preygel, Thom-Sebastiani and Duality for Matrix Factorizations, and Results on the Higher Structures of the Hochschild Invariants, Thesis (Ph.D.), M.I.T. 2012
- [63] J. P. Pridham, Unifying derived deformation theories, Adv. Math. 224 (2010), 772–826.
- [64] C. W. Rezk, Spaces of algebra structures and cohomology of operads, Thesis, MIT, 1996.
- [65] C. W. Rezk, A model for the homotopy theory of homotopy theory, Trans. Amer. Math. Soc. 353 (2001), 973–1007.
- [66] P. Salvatore, N. Wahl, Framed discs operads and Batalin-Vilkovisky algebras, Quart. J. Math. 54 (2003), 213-231.
- [67] P. Severa, Formality of the chain operad of framed little disks, Lett. Math. Phys. 93 (2010), 29–35.

- [68] S. Schwede, B. Shipley, Equivalences of monoidal model categories, Algebr. Geom. Topol. 3 (2003), 287–334.
- [69] B. Shoikhet, Tetramodules over a bialgebra form a 2-fold monoidal category, Appl. Categ. Structures 21 (2013), no. 3, 291–309.
- [70] D. Sullivan, Infinitesimal computations in topology, Inst. Hautes Études Sci. Publ. Math. 47 (1977), 269–331.
- [71] R. Taillefer, Cohomology theories of Hopf bimodules and cup-product, Algebr. Represent. Theory 7 (2004), no. 5, 471–490.
- [72] D. Tamarkin, Another proof of M. Kontsevich formality theorem, preprint arXiv:math/9803025, 1998.
- [73] D. Tamarkin. Formality of chain operad of little discs, Lett. Math. Phys. 66 (2003), 65–72.
- [74] D. Tamarkin, Deformation complex of a d-algebra is a (d+1)-algebra, preprint arXiv:math/0010072.
- [75] B. Toën, G. Vezzosi, Homotopical algebraic geometry II. Geometric stacks and applications, Mem. Amer. Math. Soc. 193 (2008), no. 902, x+224 pp.
- [76] B. Toën, Derived algebraic geometry, EMS Surv. Math. Sci. 1 (2014), no. 2, 153–240.
- [77] B. Toën, Derived Algebraic Geometry and Deformation Quantization, ICM lecture.
- [78] B. . Toën, *Problèmes de modules formels*, Séminaire BOURBAKI 68ème année, 2015-2016, no 1111
- [79] B. Vallette, A Koszul duality for props, Trans. Amer. Math. Soc. 359 (2007), 4865-4943.
- [80] S. Yalin, The homotopy theory of bialgebras over pairs of operads, J. Pure Appl. Algebra 218(2014), 973–991.
- [81] S. Yalin, Maurer-Cartan spaces of filtered L_{∞} -algebras, J. Homotopy Relat. Struct. (2015), published online, DOI 10.1007/s40062-015-0108-9.
- [82] S. Yalin, Moduli stacks of algebraic structures and deformation theory, to appear in J. Noncommut. Geom.
- [83] S. Yalin, Function spaces and classifying spaces of algebras over a prop, to appear in Algebr. Geom. Topol.